



## 2 Construction of Hypergeometric Solutions

### 2.1 Discrete Riccati Equation and Its Linearization

The  $q$ -Painlevé equations admit particular solutions characterized by discrete Riccati equations for special values of parameters. Reduction to discrete Riccati equation has been already done for all the  $q$ -Painlevé equations[3, 6, 7, 8]. We also note that such special situations have clear geometrical meaning, as was discussed in [1]. The basic idea for constructing hypergeometric solution is as follows: we linearize the discrete Riccati equations to yield second order linear  $q$ -difference equations. We then identify them with the three-term relation of an appropriate basic hypergeometric series.

Let us explain this procedure in detail. Suppose we have a discrete Riccati equation of the form

$$\bar{z} = \frac{Az + B}{Cz + D}, \quad (2)$$

where  $z = z(t)$  and  $\bar{z} = z(qt)$ . We also use the notation  $\underline{z} = z(t/q)$ , and so forth. Moreover, the coefficients  $A, B, C$  and  $D$  are functions of  $t$ . First let us put an ansatz

$$z = \frac{F}{G}. \quad (3)$$

Then the discrete Riccati equation is linearized to

$$\frac{\bar{F}}{H} = AF + BG, \quad \frac{\bar{G}}{H} = CF + DG, \quad (4)$$

where  $H$  is an arbitrary decoupling factor. Eliminating  $G$  from eq.(4) we have for  $F$  the three-term relation

$$\bar{F} + c_1 F + c_2 \underline{F} = 0, \quad c_1 = -\frac{H}{B}(A\underline{B} + B\underline{D}), \quad c_2 = \frac{B}{B}H\underline{H}(A\underline{D} - \underline{B}C). \quad (5)$$

The three-term relation for a basic hypergeometric series often takes the form

$$V_1(\bar{\Phi} - \Phi) + V_2\Phi + V_3(\underline{\Phi} - \Phi) = 0, \quad (6)$$

where the coefficients  $V_1, V_2$  and  $V_3$  are factorized into binomials involving the independent variable and parameters. Comparing eqs.(5) with (6), we have

$$\frac{V_2}{V_1} = 1 + c_1 + c_2, \quad \frac{V_3}{V_2} = c_2. \quad (7)$$

We look for the decoupling factor  $H$  so that these quantities factorize. We then identify the three-term relation with that for appropriate hypergeometric function. This is done by trial and error with the aid of computer algebra, but it is not practically difficult since we already know the hypergeometric function and its three-term relation to appear for each  $q$ -Painlevé equation.

**Step 1.** Find the decoupling factor  $H$  such that

$$\frac{V_2}{V_1} = 1 + c_1 + c_2, \quad \frac{V_3}{V_2} = c_2, \quad (8)$$

factorize. Then identify the three-term relation

$$V_1(\bar{F} - F) + V_2F + V_3(\underline{F} - F) = 0, \quad (9)$$

with that for an appropriate hypergeometric function.

Similarly, we have for  $G$  the three-term relation

$$\bar{G} + \tilde{d}_1 G + \tilde{d}_2 \underline{G} = 0, \quad \tilde{d}_1 = -\frac{H}{C}(D\underline{C} + C\underline{A}), \quad \tilde{d}_2 = \frac{C}{C}H\underline{H}(A\underline{D} - \underline{B}C). \quad (10)$$

However, usually  $1 + \tilde{d}_1 + \tilde{d}_2$  does not factorize for  $H$  obtained above. Replacing  $G$  with  $\kappa G$  ( $\bar{\kappa} = k\kappa$ ), we have

$$z = \frac{1}{\kappa} \frac{F}{G}, \quad (11)$$

$$\bar{G} + d_1 G + d_2 \underline{G} = 0, \quad d_1 = -\frac{H}{\underline{kC}}(D\underline{C} + C\underline{A}), \quad d_2 = \frac{C}{\underline{C}} \frac{HH}{\underline{k}k}(AD - \underline{BC}). \quad (12)$$

We then look for  $k$  so that  $1 + d_1 + d_2$  factorizes, and identify eq.(12) with the three-term relation of the same hypergeometric function as  $F$  with different parameters. Putting  $H/k = K$ , this is equivalent to the following procedure:

**Step 2.** *In the three-term relation*

$$\bar{G} + d_1 G + d_2 \underline{G} = 0, \quad d_1 = -\frac{K}{\underline{C}}(D\underline{C} + C\underline{A}), \quad d_2 = \frac{C}{\underline{C}} \frac{KK}{\underline{C}}(AD - \underline{BC}), \quad (13)$$

find decoupling factor  $K$  so that

$$\frac{U_2}{U_1} = 1 + d_1 + d_2, \quad \frac{U_3}{U_2} = d_2, \quad (14)$$

factorize. Then identify the three-term relation with that for an appropriate hypergeometric function. Now we have

$$z = \frac{1}{\kappa} \frac{\theta_1 \Phi}{\theta_2 \Psi}, \quad \frac{H}{K} = k, \quad \frac{\bar{\kappa}}{\kappa} = k, \quad (15)$$

where  $\Phi$  and  $\Psi$  are some hypergeometric functions, and  $\theta_i$  ( $i = 1, 2$ ) are constants (gauge factors).

Finally we determine the gauge factors  $\theta_1$  and  $\theta_2$ :

**Step 3.** *Compare the linear relations,*

$$\frac{\theta_1 \bar{\Phi}}{H} = A\theta_1 \Phi + \kappa B\theta_2 \Psi, \quad \frac{\bar{\kappa} \theta_2 \bar{\Psi}}{H} = C\theta_1 \Phi + \kappa D\theta_2 \Psi, \quad (16)$$

with contiguity relations of the hypergeometric functions to determine  $\theta_1$  and  $\theta_2$ .

## 2.2 An Example: Case of $E_7^{(1)}$

In this section we demonstrate the construction of the hypergeometric solution to the  $q$ -Painlevé equation of type  $E_7^{(1)}$  as an example, following the procedure in the previous section.

Before proceeding, let us first summarize the definition and terminology of the basic hypergeometric series [9]. The basic hypergeometric series  ${}_r\varphi_s$  is given by

$${}_r\varphi_s \left( \begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; q, z \right) = \sum_{n=0}^{\infty} \frac{(a_1; q)_n \cdots (a_r; q)_n}{(b_1; q)_n \cdots (b_s; q)_n (q; q)_n} [(-1)^n q^{\binom{n}{2}}]^{1+s-r} z^n, \quad (17)$$

$$(a; q)_n = (1-a)(1-qa) \cdots (1-q^{n-1}a).$$

The basic hypergeometric series  ${}_{r+1}\varphi_r$  is called *balanced*<sup>1</sup> if the condition

$$qa_1 a_2 \cdots a_{r+1} = b_1 b_2 \cdots b_r, \quad z = q, \quad (18)$$

is satisfied, and is called *very-well-poised* if the condition

$$qa_1 = a_2 b_1 = \cdots = a_{r+1} b_r, \quad a_2 = qa_1^{\frac{1}{2}}, \quad a_3 = -qa_1^{\frac{1}{2}}, \quad (19)$$

<sup>1</sup>The “balanced  ${}_3\varphi_2$ ” in the diagram (1) is due to the convention that was used in [10].

is satisfied. A very-well-poised hypergeometric series  ${}_{r+1}\varphi_r$  is denoted as  ${}_{r+1}W_r$ :

$${}_{r+1}W_r(a_1; a_4, \dots, a_{r+1}; q, z) = {}_r\varphi_s \left( \begin{matrix} a_1, qa_1^{\frac{1}{2}}, -qa_1^{\frac{1}{2}}, a_4, \dots, a_{r+1} \\ a_1^{\frac{1}{2}}, -a_1^{\frac{1}{2}}, qa_1/a_4, \dots, qa_1/a_{r+1} \end{matrix} ; q, z \right). \quad (20)$$

Now the  $q$ -Painlevé equation of type  $E_7^{(1)}$  is given by [2, 3]

$$\begin{aligned} \frac{(\overline{fg} - \overline{it})(fg - t^2)}{(\overline{fg} - 1)(fg - 1)} &= \frac{(f - b_1t)(f - b_2t)(f - b_3t)(f - b_4t)}{(f - b_5)(f - b_6)(f - b_7)(f - b_8)}, \\ \frac{(fg - t^2)(\underline{fg} - t\underline{t})}{(fg - 1)(\underline{fg} - 1)} &= \frac{\left(g - \frac{t}{b_1}\right)\left(g - \frac{t}{b_2}\right)\left(g - \frac{t}{b_3}\right)\left(g - \frac{t}{b_4}\right)}{\left(g - \frac{1}{b_5}\right)\left(g - \frac{1}{b_6}\right)\left(g - \frac{1}{b_7}\right)\left(g - \frac{1}{b_8}\right)}, \end{aligned} \quad (21)$$

where  $t$  is the independent variable and  $b_i$  ( $i = 1, \dots, 8$ ) are parameters satisfying

$$\overline{t} = qt, \quad b_1b_2b_3b_4 = q, \quad b_5b_6b_7b_8 = 1. \quad (22)$$

**Proposition 2.1** [3, 6] *In case of  $b_1b_3 = b_5b_7$ , eq.(21) admits a specialization to the discrete Riccati equation,*

$$\overline{g} = \frac{(\overline{it} - 1)f + t\{-(b_6 + b_8)\overline{t} + (b_2 + b_4)\}}{\{-(b_6 + b_8) + (b_2 + b_4)t\}f + b_6b_8(1 - \overline{it})}, \quad (23)$$

$$f = \frac{(t^2 - 1)b_5b_7g + t\{(b_1 + b_3) - (b_5 + b_7)t\}}{\{t(b_1 + b_3) - (b_5 + b_7)\}g + (1 - t^2)}. \quad (24)$$

As was pointed out in [1], in the cases of  $E_{6,7,8}^{(1)}$  the variables  $f$  and  $g$  are not expressed by ratio of hypergeometric functions. We choose the variable as<sup>2</sup>

$$z = \frac{g - t/b_1}{g - 1/b_5}. \quad (25)$$

Then the discrete Riccati equation (23) and (24) is rewritten as

$$\overline{z} = \frac{Az + B}{Cz + D},$$

with

$$\begin{aligned} A &= b_1b_5(-b_3 + b_5t) \\ &\times \left[ b_4b_6b_8q^2t^3 + (b_1b_4b_5 - b_4^2b_5 - b_1b_4b_6 - b_1b_4b_8 - b_5b_6b_8q)qt^2 \right. \\ &\quad \left. + (b_1b_4^2 + b_4b_5b_6q + b_4b_5b_8q + b_1b_6b_8q - b_4b_6b_8q)t - b_1b_4b_5 \right], \end{aligned} \quad (26)$$

$$B = -(b_1 - b_4)b_5^2(b_1b_4 - qb_6b_8)t(b_5 - b_3t)(-1 + qt^2), \quad (27)$$

$$C = -b_1^2b_4(b_5 - b_6)(b_5 - b_8)(b_3 - b_5t)(-1 + qt^2), \quad (28)$$

$$\begin{aligned} D &= b_1b_5(b_5 - b_3t) \\ &\times \left[ -b_1b_4b_5qt^3 + (b_1b_4^2 + b_4b_5b_6q + b_4b_5b_8q + b_1b_6b_8q - b_4b_6b_8q)t^2 \right. \\ &\quad \left. + (b_1b_4b_5 - b_4^2b_5 - b_1b_4b_6 - b_1b_4b_8 - b_5b_6b_8q)t + b_4b_6b_8 \right]. \end{aligned} \quad (29)$$

<sup>2</sup>This variable  $z$  should be understood as a ratio of  $\tau$  functions.

**Step 1.** We choose the decoupling factor  $H$  as

$$H = \frac{1}{qb_1b_5(b_5t - b_3)(b_{1t} - b_5)(b_{4t} - b_6)(b_{4t} - b_8)}. \quad (30)$$

Then we have for  $F$  the three-term relation

$$\begin{aligned} V_1(\overline{F} - F) + V_2F + V_3(\underline{F} - F) &= 0, \\ \frac{V_2}{V_1} &= \frac{b_5(b_3 - b_4)(b_1b_4 - qb_6b_8)(-1 + t)(1 + t)(-1 + qt^2)}{q(b_5 - b_{1t})(-b_6 + b_{4t})(-b_8 + b_{4t})(-b_3 + b_5t)}, \\ \frac{V_3}{V_1} &= \frac{(b_5 - b_3t)(-b_1 + b_5t)(-b_4 + b_6t)(-b_4 + b_8t)(-1 + qt^2)}{(b_5 - b_{1t})(-b_6 + b_{4t})(-b_8 + b_{4t})(-b_3 + b_5t)(-q + t^2)}. \end{aligned} \quad (31)$$

The three-term relation for the very-well-poised basic hypergeometric series

$$\Phi = {}_8W_7 \left( a_0; a_1, a_2, a_3, a_4, a_5; q, \frac{q^2a_0^2}{a_1a_2a_3a_4a_5} \right), \quad (32)$$

is given by [11]

$$\begin{aligned} U_1(\overline{\Phi} - \Phi) + U_2\Phi + U_3(\underline{\Phi} - \Phi) &= 0, \quad \overline{\Phi} = \Phi|_{a_2 \rightarrow qa_2, a_3 \rightarrow a_3/q}, \quad \underline{\Phi} = \Phi|_{a_2 \rightarrow a_2/q, a_3 \rightarrow qa_3}, \\ U_1 &= \frac{(1 - a_2) \left(1 - \frac{a_0}{a_2}\right) \left(1 - \frac{qa_0}{a_2}\right) \left(1 - \frac{qa_0}{a_1a_3}\right) \left(1 - \frac{qa_0}{a_3a_4}\right) \left(1 - \frac{qa_0}{a_3a_5}\right)}{a_3 \left(1 - \frac{a_2}{a_3}\right) \left(1 - \frac{qa_2}{a_3}\right)}, \quad U_3 = U_1|_{a_2 \leftrightarrow a_3}, \\ U_2 &= \frac{qa_0^2}{a_1a_2a_3a_4a_5} \left(1 - \frac{qa_0}{a_2a_3}\right) (1 - a_1)(1 - a_4)(1 - a_5). \end{aligned} \quad (33)$$

Comparing eqs.(31) with (33), we identify  $F$  with  ${}_8W_7$  as

$$F \propto {}_8W_7 \left( \frac{b_1b_8}{b_3b_5}, \frac{qb_8}{b_5}, \frac{b_{1t}}{b_5}, \frac{b_1}{b_{5t}}, \frac{b_2}{b_3}, \frac{b_4}{b_3}, \frac{b_5}{b_6}; q, \frac{b_5}{b_6} \right). \quad (34)$$

**Step 2.** We choose the decoupling factor  $K$  as

$$K = \frac{1}{b_1b_5(qb_5t - b_3)(b_{1t} - b_5)(b_{4t} - b_6)(b_{4t} - b_8)}. \quad (35)$$

Then we have for  $G$  the three-term relation

$$\begin{aligned} X_1(\overline{G} - G) + X_2F + X_3(\underline{G} - G) &= 0, \\ \frac{X_2}{X_1} &= \frac{b_5(b_3 - b_4)(b_1b_4 - b_6b_8)(-1 + t)(1 + t)(-1 + qt^2)}{(b_5 - b_{1t})(-b_6 + b_{4t})(-b_8 + b_{4t})(-b_3 + qb_5t)}, \\ \frac{X_3}{X_1} &= \frac{(qb_5 - b_3t)(-b_1 + b_5t)(-b_4 + b_6t)(-b_4 + b_8t)(-1 + qt^2)}{(b_5 - b_{1t})(-b_6 + b_{4t})(-b_8 + b_{4t})(-b_3 + qb_5t)(-q + t^2)}. \end{aligned} \quad (36)$$

Comparing eqs.(36) with (33), we have

$$G \propto {}_8W_7 \left( \frac{b_1b_8}{b_3b_5}, \frac{b_8}{b_5}, \frac{b_{1t}}{b_5}, \frac{b_1}{b_{5t}}, \frac{b_2}{b_3}, \frac{b_4}{b_3}; q, \frac{qb_5}{b_6} \right). \quad (37)$$

Moreover, from  $k = H/K = (1 - b_3/qb_5t)/(1 - b_3/b_5t)$ , we have  $\kappa = 1 - b_3/b_5t$ . Therefore we obtain

$$z \propto \frac{1}{1 - \frac{b_3}{b_5t}} \frac{{}_8W_7 \left( a_0; qa_1, a_2, a_3, a_4, a_5; q, \frac{qa_0^2}{a_1a_2a_3a_4a_5} \right)}{{}_8W_7 \left( a_0, a_1, a_2, a_3, a_4, a_5; q, \frac{q^2a_0^2}{a_1a_2a_3a_4a_5} \right)}, \quad (38)$$

with

$$a_0 = \frac{b_1 b_8}{b_3 b_5}, \quad a_1 = \frac{b_8}{b_5}, \quad a_2 = \frac{b_1 t}{b_5}, \quad a_3 = \frac{b_1}{b_5 t}, \quad a_4 = \frac{b_2}{b_3}, \quad a_5 = \frac{b_4}{b_3}, \quad \frac{q^2 a_0^2}{a_1 a_2 a_3 a_4 a_5} = \frac{q b_5}{b_6}. \quad (39)$$

**Step 3.** Let us put

$$F = \theta(qa_1, a_2, a_3) \Phi(qa_1, a_2, a_3), \quad G = \theta(a_1, a_2, a_3) \Phi(a_1, a_2, a_3), \quad (40)$$

where  $\theta(a_1, a_2, a_3)$  is a gauge factor to be determined. Here, we have omitted the dependence of  $a_0, a_4$  and  $a_5$ , since they are not relevant to the calculation. Then linear equations (16) yield

$$\frac{1}{\kappa H B} \frac{\theta(qa_1, qa_2, a_3/q)}{\theta(a_1, a_2, a_3)} \Phi(qa_1, qa_2, a_3/q) = \frac{A}{\kappa B} \frac{\theta(qa_1, a_2, a_3)}{\theta(a_1, a_2, a_3)} \Phi(qa_1, a_2, a_3) + \Phi(a_1, a_2, a_3), \quad (41)$$

and

$$\frac{\bar{\kappa}}{\kappa H D} \frac{\theta(a_1, qa_2, a_3/q)}{\theta(a_1, a_2, a_3)} \Phi(a_1, qa_2, a_3/q) = \frac{C}{\kappa D} \frac{\theta(qa_1, a_2, a_3)}{\theta(a_1, a_2, a_3)} \Phi(qa_1, a_2, a_3) + \Phi(a_1, a_2, a_3), \quad (42)$$

respectively. Now, we have the contiguity relations for  $\Phi = {}_8W_7$  [11]

$$\frac{a_1 \left(1 - \frac{a_0 q}{a_1 a_3}\right) \left(1 - \frac{a_0 q}{a_1 a_4}\right) \left(1 - \frac{a_0 q}{a_1 a_5}\right)}{1 - \frac{a_0 q}{a_1}} \Phi(a_1/q, a_2, a_3) - (a_1 \leftrightarrow a_2) \quad (43)$$

$$= (a_1 - a_2) \left(1 - \frac{a_0^2 q^2}{a_1 a_2 a_3 a_4 a_5}\right) \Phi(a_1, a_2, a_3),$$

$$(a_2 - 1) \left(1 - \frac{a_0}{a_2}\right) \Phi(a_1/q, qa_2, a_3) + \left(1 - \frac{a_1}{q}\right) \left(1 - \frac{a_0 q}{a_1}\right) \Phi(a_1, a_2, a_3) \quad (44)$$

$$= \left(a_2 - \frac{a_1}{q}\right) \left(1 - \frac{a_0 q}{a_1 a_2}\right) \Phi(a_1/q, a_2, a_3).$$

We denote eqs.(43) and (44) as CR1[ $a_1, a_2, a_3$ ] and CR2[ $a_1, a_2, a_3$ ], respectively. Moreover, note that the relations CR1[ $a_1, a_2, a_3$ ] and CR2[ $a_1, a_2, a_3$ ] hold for any permutation of  $a_1, a_2$  and  $a_3$ , since these parameters are on equal footing in  $\Phi = {}_8W_7$ .

Now we eliminate  $\Phi(a_1, a_2, a_3/q)$  from CR1[ $a_1, a_3, a_2$ ] and CR2[ $a_3, a_2, a_1$ ]. Shifting  $a_1$  to  $qa_1$ , we have a linear relation among  $\Phi(a_1, qa_2, a_3/q)$ ,  $\Phi(a_1, a_2, a_3)$  and  $\Phi(a_1/q, a_2, a_3)$ , which should coincide with eq. (41). Similarly, eliminating  $\Phi(qa_1, a_2, a_3/q)$  from CR1[ $qa_1, a_3, a_2$ ] and CR2[ $a_3, a_1, a_2$ ], we have a linear relation among  $\Phi(qa_1, a_2, a_3)$ ,  $\Phi(a_1, a_2, a_3)$  and  $\Phi(a_1, a_2, a_3/q)$ . Elimination further  $\Phi(a_1, a_2, a_3/q)$  from this relation and CR2[ $a_3, a_2, a_1$ ] yields a linear relation among  $\Phi(a_1, qa_2, a_3/q)$ ,  $\Phi(qa_1, a_2, a_3)$  and  $\Phi(a_1, a_2, a_3)$ , which should coincide with eq.(42). From these calculations, we find that  $\theta(a_1, a_2, a_3)$  should satisfy

$$\frac{\theta(a_1, qa_2, a_3/q)}{\theta(a_1, a_2, a_3)} = \frac{\left(1 - \frac{a_2}{a_0}\right) \left(1 - \frac{a_3 a_5}{a_0 q}\right)}{\left(1 - \frac{a_2 a_5}{a_0}\right) \left(1 - \frac{a_3}{a_0 q}\right)} = \frac{\left(1 - \frac{b_3 t}{b_8}\right) \left(1 - \frac{b_4}{b_8 q t}\right)}{\left(1 - \frac{b_4 t}{b_8}\right) \left(1 - \frac{b_3}{b_8 q t}\right)}, \quad (45)$$

$$\frac{\theta(qa_1, a_2, a_3)}{\theta(a_1, a_2, a_3)} = 1 - \frac{a_1}{a_0} = 1 - \frac{b_3}{b_1},$$

$$\frac{\theta(qa_1, qa_2, a_3/q)}{\theta(a_1, a_2, a_3)} = \frac{\theta(a_1, qa_2, a_3/q)}{\theta(a_1, a_2, a_3)} \times \frac{\theta(qa_1, a_2, a_3)}{\theta(a_1, a_2, a_3)},$$

which yield

$$\theta(a_1, a_2, a_3) = \frac{\left(\frac{a_2}{a_0 q}, \frac{a_3 a_5}{a_0 q}, \frac{a_1}{a_0 q}\right)_\infty}{\left(\frac{a_2 a_5}{a q}, \frac{a_3}{a_0 q}\right)_\infty} = \frac{\left(\frac{b_3 t}{q b_8}, \frac{b_4}{q b_8 t}, \frac{b_3}{q b_1}\right)_\infty}{\left(\frac{b_4 t}{q b_8}, \frac{b_3}{q b_8 t}\right)_\infty}. \quad (46)$$

Therefore we arrive at the final result

$$\begin{aligned}
z &= \frac{1}{1 - \frac{b_3}{b_5 t}} \frac{\theta(qa_1, a_2, a_3)}{\theta(a_1, a_2, a_3)} \frac{\Phi(qa_1, a_2, a_3)}{\Phi(a_1, a_2, a_3)} \\
&= \frac{1 - \frac{b_3}{b_1}}{1 - \frac{b_3}{b_5 t}} \frac{{}_8W_7\left(a_0; qa_1, a_2, a_3, a_4, a_5; q, \frac{qa_0^2}{a_1 a_2 a_3 a_4 a_5}\right)}{{}_8W_7\left(a_0, a_1, a_2, a_3, a_4, a_5; q, \frac{q^2 a_0^2}{a_1 a_2 a_3 a_4 a_5}\right)}.
\end{aligned} \tag{47}$$

### 3 Hypergeometric Solutions

Hypergeometric solutions to other  $q$ -Painlevé equations can be constructed by the same procedure as that was demonstrated in the previous section. Instead of describing full procedure, we give a list of equations, solutions and the other data that are necessary for construction of solutions. We note that the case of  $D_5^{(1)}$  is omitted as mentioned in the introduction.

#### 3.1 Case of $E_8^{(1)}$

##### 3.1.1 Equation and Solution

(1)  $q$ -Painlevé equation [3, 6, 12]

$$\begin{aligned}
\frac{(\overline{g}st - f)(gst - f) - (\overline{s}^2 t^2 - 1)(s^2 t^2 - 1)}{\left(\frac{\overline{g}}{st} - f\right)\left(\frac{g}{st} - f\right) - \left(1 - \frac{1}{\overline{s}^2 t^2}\right)\left(1 - \frac{1}{s^2 t^2}\right)} &= \frac{P(f, t, m_1, \dots, m_7)}{P(f, t^{-1}, m_7, \dots, m_1)}, \\
\frac{(f \underline{st} - g)(fst - g) - (s^2 \underline{t}^2 - 1)(s^2 t^2 - 1)}{\left(\frac{f}{\underline{st}} - g\right)\left(\frac{f}{st} - g\right) - \left(1 - \frac{1}{s^2 \underline{t}^2}\right)\left(1 - \frac{1}{s^2 t^2}\right)} &= \frac{P(g, s, m_7, \dots, m_1)}{P(g, s^{-1}, m_1, \dots, m_7)},
\end{aligned} \tag{48}$$

where

$$\begin{aligned}
P(f, t, m_1, \dots, m_7) &= f^4 - m_1 t f^3 + (m_2 t^2 - 3 - t^8) f^2 \\
&\quad + (m_7 t^7 - m_3 t^3 + 2m_1 t) f + (t^8 - m_6 t^6 + m_4 t^4 - m_2 t^2 + 1),
\end{aligned} \tag{49}$$

and  $m_k$  ( $k = 1, 2, \dots, 7$ ) are the elementary symmetric functions of  $k$ -th degree in  $b_i$  ( $i = 1, 2, \dots, 8$ ) with

$$b_1 b_2 \cdots b_8 = 1. \tag{50}$$

Moreover,

$$\overline{t} = qt, \quad t = q^{\frac{1}{2}} s. \tag{51}$$

(2) Constraint on parameters [6]

$$qb_1 b_3 b_5 b_7 = 1, \quad b_2 b_4 b_6 b_8 = q. \tag{52}$$

(3) Hypergeometric solution

$$z = \frac{g - \left(\frac{s}{b_1} + \frac{b_1}{s}\right)}{g - \left(\frac{s}{b_8} + \frac{b_8}{s}\right)} = \lambda \frac{\Phi(q^4 a_0; a_1, q^2 a_2, \dots, q^2 a_7)}{\Phi(a_0; a_1, \dots, a_7)}, \tag{53}$$

where  $\Phi$  is defined in terms of the balanced  ${}_{10}W_9$  series by

$$\begin{aligned} \Phi(a_0; a_1, \dots, a_7) &= {}_{10}W_9(a_0; a_1, \dots, a_7; q^2, q^2) \\ &+ \frac{\left(q^2 a_0, \frac{a_7}{a_0}; q^2\right)_\infty}{\left(\frac{a_0}{a_7}, \frac{q^2 a_7^2}{a_0}; q^2\right)_\infty} \prod_{k=1}^6 \frac{\left(a_k, \frac{q^2 a_7}{a_k}; q^2\right)_\infty}{\left(\frac{q^2 a_0}{a_k}, \frac{a_k a_7}{a_0}; q^2\right)_\infty} {}_{10}W_9\left(\frac{a_7^2}{a_0}; \frac{a_1 a_7}{a_0}, \dots, \frac{a_6 a_7}{a_0}, a_7; q^2, q^2\right). \end{aligned} \quad (54)$$

Here,  $a_i$  ( $i = 0, 1, \dots, 7$ ) and  $\lambda$  are given by

$$\begin{aligned} a_0 &= \frac{1}{qb_1 b_2 b_8^2}, \quad a_1 = \frac{q^2}{b_2 b_8 t^2}, \quad a_2 = \frac{s^2}{b_2 b_8}, \\ a_i &= \frac{b_i}{b_8} \quad (i = 3, 5, 7), \quad a_i = \frac{b_i}{b_1} \quad (i = 4, 6), \end{aligned} \quad (55)$$

and

$$\lambda = \frac{b_1 b_4 b_6}{b_8 s^2} \frac{\left(1 - \frac{b_4 b_6}{b_1 b_8}\right) \left(1 - q^2 \frac{b_4 b_6}{b_1 b_8}\right) (1 - b_3 b_5 t^2) (1 - b_3 b_7 t^2) (1 - b_5 b_7 t^2) \prod_{i=2,4,6} \left(1 - \frac{b_i}{b_1}\right)}{\left(1 - \frac{s^2}{b_1 b_8}\right) \left(1 - \frac{q^2 s^2}{b_1 b_8}\right) \left(1 - \frac{b_4}{b_8}\right) \left(1 - \frac{b_6}{b_8}\right) \left(1 - \frac{q}{b_1 b_8 s^2}\right) \prod_{i=3,5,7} \left(1 - \frac{b_4 b_6}{b_1 b_i}\right)}, \quad (56)$$

respectively.

### 3.1.2 Data

(1) Riccati equation

$$\begin{vmatrix} 1 & f & g & fg \\ 1 & f_1 & g_1 & f_1 g_1 \\ 1 & f_3 & g_3 & f_3 g_3 \\ 1 & f_5 & g_5 & f_5 g_5 \end{vmatrix} = 0, \quad \begin{vmatrix} 1 & f & \bar{g} & f\bar{g} \\ 1 & f_8 & \bar{g}_8 & f_8 \bar{g}_8 \\ 1 & f_6 & \bar{g}_6 & f_6 \bar{g}_6 \\ 1 & f_4 & \bar{g}_4 & f_4 \bar{g}_4 \end{vmatrix} = 0, \quad (57)$$

where

$$f_i = b_i t + \frac{1}{b_i t}, \quad g_i = \frac{s}{b_i} + \frac{b_i}{s}. \quad (58)$$

The Riccati equation for  $z = \frac{g - g_1}{g - g_8}$  is given by

$$\bar{z} = \frac{Az + B}{Cz + D}, \quad (59)$$

$$\begin{aligned} B &= -f_{35} g_{13} g_{15} d'_{1468}, \\ C &= f_{46} \bar{g}_{48} \bar{g}_{68} d_{1358}, \\ D &= -f_{35} f_{46} f_{18} g_{13} g_{15} \bar{g}_{48} \bar{g}_{68}, \\ \Delta &= AD - BC = f_{13} f_{35} f_{15} f_{46} f_{48} f_{68} g_{13} g_{15} g_{35} g_{18} \bar{g}_{46} \bar{g}_{48} \bar{g}_{68} \bar{g}_{18}, \end{aligned} \quad (60)$$

where  $f_{ij} = f_i - f_j$  and

$$d_{1358} = \begin{vmatrix} 1 & f_1 & g_1 & f_1 g_1 \\ 1 & f_3 & g_3 & f_3 g_3 \\ 1 & f_5 & g_5 & f_5 g_5 \\ 1 & f_8 & g_8 & f_8 g_8 \end{vmatrix}, \quad d'_{1468} = \begin{vmatrix} 1 & f_1 & \bar{g}_1 & f_1 \bar{g}_1 \\ 1 & f_4 & \bar{g}_4 & f_4 \bar{g}_4 \\ 1 & f_6 & \bar{g}_6 & f_6 \bar{g}_6 \\ 1 & f_8 & \bar{g}_8 & f_8 \bar{g}_8 \end{vmatrix}, \quad (61)$$

respectively.



(2) Three-term and contiguity relations for hypergeometric function [13]

(a) Three-term relation

$$U_1(\bar{\Phi} - \Phi) + U_2\Phi + U_3(\underline{\Phi} - \Phi) = 0, \quad (62)$$

where

$$U_1 = \frac{a_1(1-a_2)\left(1-\frac{a_0}{a_2}\right)\left(1-\frac{q^2a_0}{a_2}\right)}{\left(1-\frac{q^2a_2}{a_1}\right)\prod_{j=3}^7\left(1-\frac{q^2a_0}{a_1a_j}\right)}, \quad (63)$$

$$U_2 = -(a_1 - a_2)\left(1 - \frac{q^2a_0}{a_1a_2}\right)\prod_{j=3}^7(1 - a_j), \quad U_3 = U_1|_{a_1 \leftrightarrow a_2},$$

$$\bar{\Phi} = \Phi(a_0; a_1/q^2, q^2a_2, a_3, \dots, a_7), \quad \underline{\Phi} = \Phi(a_0; q^2a_1, a_2/q^2, a_3, \dots, a_7), \quad (64)$$

and  $\Phi$  is defined by eq.(54).

(b) Contiguity relations

$$\begin{aligned} & \Phi(a_0; a_1/q^2, q^2a_2, a_3, \dots, a_7) - \Phi(a_0; a_1, a_2, a_3, \dots, a_7) \\ &= V_1 \Phi(q^4a_0^2; a_1, q^2a_2, \dots, q^2a_7), \end{aligned} \quad (65)$$

$$\begin{aligned} & V_2 \Phi(q^4a_0^2; a_1, q^2a_2, q^2a_3, \dots, q^2a_7) - V_3 \Phi(q^4a_0^2; q^2a_1, a_2, q^2a_3, \dots, q^2a_7) \\ &= V_4 \Phi(a_0; a_1, a_2, a_3, \dots, a_7), \end{aligned} \quad (66)$$

where

$$V_1 = \frac{\frac{q^2a_0}{a_2}\left(1-\frac{q^2a_2}{a_1}\right)\left(1-\frac{a_1a_2}{q^2a_0}\right)(1-q^2a_0)(1-q^4a_0)\prod_{j=3}^7(1-a_j)}{\left(1-\frac{q^2a_0}{a_1}\right)\left(1-\frac{q^4a_0}{a_1}\right)\left(1-\frac{a_0}{a_2}\right)\left(1-\frac{q^2a_0}{a_2}\right)\prod_{j=3}^7\left(1-\frac{q^2a_0}{a_j}\right)},$$

$$V_2 = \frac{a_1^2(1-a_2)\prod_{j=3}^7\left(1-\frac{q^2a_0}{a_1a_j}\right)}{\left(1-\frac{q^2a_0}{a_1}\right)\left(1-\frac{q^4a_0}{a_1}\right)}, \quad V_3 = V_2|_{a_1 \leftrightarrow a_2},$$

$$V_4 = \frac{a_1\left(1-\frac{a_2}{a_1}\right)\prod_{j=3}^7\left(1-\frac{q^2a_0}{a_j}\right)}{(1-q^2a_0)(1-q^4a_0)}.$$

(3) Decoupling factors

$$H = \frac{D}{\Delta} = -\frac{f_{18}}{f_{13}f_{15}f_{48}f_{68}g_{35}g_{18}\bar{g}_{46}\bar{g}_{18}}, \quad K = \frac{1}{D} = -\frac{1}{f_{35}f_{46}f_{18}g_{13}g_{15}\bar{g}_{48}\bar{g}_{68}}, \quad (68)$$

so that

$$k = \frac{H}{K} = \frac{D^2}{\Delta} = \frac{f_{35}f_{46}f_{18}^2g_{13}g_{15}\bar{g}_{48}\bar{g}_{68}}{f_{13}f_{15}f_{48}f_{68}g_{35}\bar{g}_{46}\bar{g}_{18}}. \quad (69)$$

(4) Identification

$$z = \frac{1}{\kappa} \frac{F}{G}, \quad F \propto \Phi(q^4a_0; a_1, q^2a_2, \dots, q^2a_7), \quad G \propto \Phi(a_0; a_1, \dots, a_7), \quad \bar{\kappa} = \kappa\kappa, \quad (70)$$

where  $a_i$  ( $i = 0, \dots, 7$ ) are given by eq.(55).

- (5) Gauge factors  
Putting

$$\begin{aligned} F &= \theta(q^4 a_0; a_1, q^2 a_2, \dots, q^2 a_7) \Phi(q^4 a_0; a_1, q^2 a_2, \dots, q^2 a_7), \\ G &= \theta(a_0; a_1 \dots, a_7) \Phi(a_0; a_1 \dots, a_7), \end{aligned} \quad (71)$$

we have:

$$\frac{\theta(a_0; a_1/q^2, a_2 q^2, \dots, a_7)}{\theta(a_0; a_1 \dots, a_7)} = 1, \quad \frac{1}{\kappa} \frac{\theta(q^4 a_0; a_1, q^2 a_2, \dots, q^2 a_7)}{\theta(a_0; a_1 \dots, a_7)} = \lambda. \quad (72)$$

## 3.2 Case of $E_6^{(1)}$

### 3.2.1 Equation and Solution

- (1)  $q$ -Painlevé equation[3, 14, 15]

$$\begin{aligned} (f\bar{g} - 1)(fg - 1) &= \bar{t}\bar{t} \frac{(f - b_1 t)(f - b_2 t)(f - b_3 t)(f - b_4 t)}{(f - b_5 t) \left(f - \frac{t}{b_5}\right)}, \\ (fg - 1)(\underline{f}g - 1) &= t^2 \frac{\left(g - \frac{1}{b_1}\right) \left(g - \frac{1}{b_2}\right) \left(g - \frac{1}{b_3}\right) \left(g - \frac{1}{b_4}\right)}{(g - b_6 t) \left(g - \frac{t}{b_6}\right)}, \end{aligned} \quad (73)$$

$$\bar{t} = qt, \quad b_1 b_2 b_3 b_4 = 1.$$

- (2) Constraint on parameters [3, 15]

$$b_1 b_2 = b_5 b_6. \quad (74)$$

- (3) Hypergeometric solution

$$z = \frac{g - \frac{1}{b_1}}{g - t b_6} = \frac{1 - \frac{b_3}{b_1}}{1 - \frac{b_1 b_2 b_3 t}{b_5}} \frac{\Phi(qa, b, c, d, e)}{\Phi(a, b, qc, d, e)}, \quad (75)$$

where  $\Phi$  is the balanced  ${}_3\phi_2$  series defined by

$$\Phi(a, b, c, d, e) = {}_3\phi_2 \left( \begin{matrix} a, b, c \\ d, e \end{matrix}; q, \frac{de}{abc} \right), \quad (76)$$

with

$$a = \frac{b_3 b_5}{t}, \quad b = \frac{b_3}{b_2}, \quad c = b_1^2 b_2 b_3, \quad d = q \frac{b_3 b_5^2}{b_2}, \quad e = q b_1 b_2 b_3^2. \quad (77)$$

### 3.2.2 Data

- (1) Riccati equation [3, 15]

$$\bar{g} = \frac{1 + \frac{b_5 \bar{t}}{b_1 b_2} (f - b_1 - b_3)}{f - \bar{t} b_5}, \quad f = \frac{1 + b_6 t (b_3 b_4 g - b_3 - b_4)}{g - t b_6}. \quad (78)$$

The Riccati equation for

$$z = \frac{g - 1/b_1}{g - t b_6}, \quad (79)$$

is given by,

$$\bar{z} = \frac{Az + B}{Cz + D},$$

$$\begin{aligned} A &= b_1 b_5 (b_3 b_5 - t)(b_5 - b_1 b_2 b_3 t)(-b_2 + b_5 q t), \\ B &= -b_2^2 t (b_1 - b_3)(-1 + b_1^2 b_2 b_3)(-b_2 + b_5 q t), \\ C &= q t b_1 (b_1 b_2 - b_5)(b_1 b_2 + b_5)(b_3 b_5 - t)(-b_5 + b_1 b_2 b_3 t), \\ D &= -b_5 \left[ b_1 b_2 b_3 b_5^2 + (-b_1^3 b_2^2 b_3 b_5 - b_1^3 b_2^2 b_3 b_5 q - b_2 b_3 b_5^3 q) t \right. \\ &\quad \left. + (b_1^3 b_2^2 - b_1^2 b_2^2 b_3 + b_1^4 b_2^3 b_3^2 - b_1 b_5^2 + b_3 b_5^2 + b_1^3 b_2 b_3 b_5^2 + b_1^2 b_2^2 b_3 b_5^2 - b_1^2 b_2 b_3^2 b_5^2 + b_1^2 b_2^2 b_3 b_5^2 q) q t^2 \right. \\ &\quad \left. - b_1^4 b_2^3 b_3 b_5 q^2 t^3 \right]. \end{aligned} \quad (80)$$

(2) Three-term and contiguity relations for hypergeometric function [10]

(a) Three-term relation

$$V_1 (\bar{\Phi} - \Phi) + V_2 \Phi + V_3 (\Phi - \underline{\Phi}) = 0, \quad (81)$$

where

$$\begin{aligned} V_1 &= \left(1 - \frac{q}{z}\right)(1 - a), \quad V_2 = (1 - b)(1 - c), \quad V_3 = \frac{a}{z} \left(1 - \frac{d}{a}\right) \left(1 - \frac{e}{a}\right), \\ \Phi &= \Phi(a, b, c, d, e) = {}_3\phi_2 \left( \begin{matrix} a, b, c \\ d, e \end{matrix}; q; z \right), \quad z = \frac{de}{abc}, \quad \bar{\Phi} = \Phi|_{a \rightarrow qa}, \quad \underline{\Phi} = \Phi|_{a \rightarrow a/q}. \end{aligned} \quad (82)$$

(b) Contiguity relations

$$(a - c)\Phi(a, b, c, d, e) + (1 - a)\Phi(qa, b, c, d, e) - (1 - c)\Phi(a, b, qc, d, e) = 0, \quad (83)$$

$$\begin{aligned} (a - c)(de - abc)\Phi(a, b, c, d, e) + bc(d - a)(e - a)\Phi(a/q, b, c, d, e) \\ - ab(d - c)(e - c)\Phi(a, b, c/q, d, e) = 0. \end{aligned} \quad (84)$$

(3) Decoupling factors

$$\begin{aligned} H &= \frac{1}{b_1 b_3 b_5 (-b_5 + b_1^2 b_2 t)(b_5 - b_1 b_2 b_3 t)(b_2 - q b_5 t)}, \\ K &= \frac{1}{b_1 b_3 b_5 (-b_5 + b_1^2 b_2 t)(b_5 - q b_1 b_2 b_3 t)(b_2 - q b_5 t)}, \\ k &= \frac{H}{K} = \frac{b_5 - q b_1 b_2 b_3 t}{b_5 - b_1 b_2 b_3 t}, \quad \kappa = 1 - \frac{b_1 b_2 b_3 t}{b_5}. \end{aligned} \quad (85)$$

(4) Identification

$$z = \frac{1}{\kappa} \frac{F}{G}, \quad F \propto \Phi(qa, b, c, d, e), \quad G \propto \Phi(a, b, qc, d, e), \quad (86)$$

where  $a, \dots, e$  are given by eq.(77).

(5) Gauge factors

Putting

$$F = \theta(qa, b, c, d, e)\Phi(qa, b, c, d, e), \quad G = \theta(a, b, qc, d, e)\Phi(a, b, qc, d, e), \quad (87)$$

we have:

$$\frac{\theta(a, b, c, d, e)}{\theta(a, b, qc, d, e)} = \frac{\theta(qa, b, c, d, e)}{\theta(a, b, qc, d, e)} = 1 - \frac{b_3}{b_1}, \quad \frac{\theta(a/q, b, qc, d, e)}{\theta(a, b, qc, d, e)} = 1. \quad (88)$$

### 3.3 Case of $A_4^{(1)}$

#### 3.3.1 Equation and Solution

(1)  $q$ -Painlevé equation [3]

$$\begin{aligned}\bar{g}g &= \frac{\left(f + \frac{a_1}{t}\right)\left(f + \frac{1}{a_1t}\right)}{1 + a_3f}, \\ f\bar{f} &= \frac{\left(g + \frac{a_2}{s}\right)\left(g + \frac{1}{a_2s}\right)}{1 + g/a_3}, \\ \bar{t} &= qt, \quad t = q^{\frac{1}{2}}s.\end{aligned}\tag{89}$$

(2) Constraint on parameters [3]

$$a_1a_2a_3^2 = q^{-\frac{1}{2}}.\tag{90}$$

(3) Hypergeometric solution

$$g = -\frac{1}{a_1a_3^2t} \frac{\Phi(\alpha_1, \alpha_2, z)}{\Phi(\alpha_1, q\alpha_2, z)}, \quad f = \frac{1}{a_3} \left(1 - \frac{1}{a_2^2}\right) \frac{\Phi(q\alpha_1, q\alpha_2, z)}{\Phi(\alpha_1, q\alpha_2, z)},\tag{91}$$

where  $\Phi$  is the  ${}_2\phi_1$  series defined by

$$\Phi(\alpha_1, \alpha_2, z) = {}_2\phi_1\left(\begin{matrix} \alpha_1, \alpha_2 \\ 0 \end{matrix}; q, z\right),\tag{92}$$

with

$$\alpha_1 = \frac{1}{a_2^2}, \quad \alpha_2 = a_1^2, \quad z = \frac{t}{a_1a_3}.\tag{93}$$

Note that the solution is also expressible in terms of  ${}_1\phi_1$  series by using the formula [16],

$${}_2\phi_1\left(\begin{matrix} a, b \\ 0 \end{matrix}; q, z\right) = \frac{(bz; q)_\infty}{(z; q)_\infty} {}_1\phi_1\left(\begin{matrix} b \\ bz \end{matrix}; q, az\right).\tag{94}$$

#### 3.3.2 Data

(1) Riccati equation [3]

$$\bar{g} = \frac{a_3^2g + \frac{1-a_1^2}{a_1t}}{-a_3g + \left(\frac{1}{a_3^2} - \frac{1}{a_1a_3t}\right)}, \quad g = -\frac{f + \frac{1}{a_1t}}{a_3^2}.\tag{95}$$

(2) Three-term and contiguity relations for hypergeometric function

(a) Three-term relation

$$\frac{\alpha_1\alpha_2}{q}z(\bar{\Phi} - \Phi) + \frac{z}{q}(1 - \alpha_1)(1 - \alpha_2)\Phi + \left(\frac{z}{q} - 1\right)(\underline{\Phi} - \Phi) = 0,\tag{96}$$

where

$$\Phi(\alpha_1, \alpha_2, z) = {}_2\phi_1\left(\begin{matrix} \alpha_1, \alpha_2 \\ 0 \end{matrix}; q, z\right), \quad \bar{\Phi} = \Phi|_{z \rightarrow qz}, \quad \underline{\Phi} = \Phi|_{z \rightarrow z/q}.\tag{97}$$

(b) Contiguity relations

$$\Phi(\alpha_1, \alpha_2, z) - \alpha_2 \Phi(\alpha_1, \alpha_2, qz) = (1 - \alpha_2) \Phi(\alpha_1, q\alpha_2, z), \quad (98)$$

$$\Phi(\alpha_1, \alpha_2/q, z) - \Phi(\alpha_1, \alpha_2, z) = -\frac{\alpha_2 z}{q} (1 - \alpha_1) \Phi(q\alpha_1, \alpha_2, z), \quad (99)$$

$$\alpha_1 \alpha_2 z \Phi(\alpha_1, q\alpha_2, qz) = \Phi - (1 - \alpha_2 z) \Phi(\alpha_1, q\alpha_2, z). \quad (100)$$

(3) Decoupling factors

$$H = \frac{1}{a_1^2 a_3^2}, \quad K = \frac{1}{q a_1^2 a_3^2}, \quad k = \frac{H}{K} = q, \quad \kappa = t. \quad (101)$$

(4) Identification

$$f = \frac{1}{\kappa} \frac{F}{G}, \quad F \propto \Phi(\alpha_1, \alpha_2, z), \quad G \propto \Phi(\alpha_1, q\alpha_2, z), \quad (102)$$

with parameters given in eq.(93).

(5) Gauge factors

Putting

$$F = \theta(\alpha_1, \alpha_2, z) \Phi(\alpha_1, \alpha_2, z), \quad G = \theta(\alpha_1, q\alpha_2, z) \Phi(\alpha_1, q\alpha_2, z), \quad (103)$$

we have:

$$\frac{\theta(\alpha_1, \alpha_2, qz)}{\theta(\alpha_1, \alpha_2, z)} = 1, \quad \frac{\theta(\alpha_1, q\alpha_2, z)}{\theta(\alpha_1, \alpha_2, z)} = -a_1 a_3^2. \quad (104)$$

### 3.4 Case of $(A_2 + A_1)^{(1)}$

#### 3.4.1 Equation and Solution

(1)  $q$ -Painlevé equation [2, 3, 7, 8, 17]

$$\bar{g} g f = b_0 \frac{1 + a_0 t f}{a_0 t + f}, \quad g f \bar{f} = b_0 \frac{\frac{a_1}{t} + g}{1 + \frac{a_1}{t} g}, \quad \bar{t} = qt. \quad (105)$$

Eq. (105) admits two different specializations for hypergeometric solutions: (a) specialization of  $b_i$  (parameter of  $A_1$ ), (b) specialization of  $a_i$  (parameter of  $A_2$ ). See also [7, 8] for details.

(2) Constraint on parameters

(a)

$$b_0 = q. \quad (106)$$

(b)

$$a_0 a_1 = q. \quad (107)$$

(3) Hypergeometric solution

(a)

$$g = -\frac{a_1}{t} \left( 1 - \frac{q^2}{a_0^2 a_1^2} \right) \frac{\Phi(b, z)}{\Phi(q^2 b, z)}, \quad f = \frac{q^2 t}{a_0 a_1^2} \frac{1}{1 - \frac{q^2}{a_0^2 a_1^2}} \frac{\Phi(q^2 b, q^2 z)}{\Phi(b, z)}, \quad (108)$$

where

$$\Phi = \Phi(b, z) = {}_1\varphi_1 \left( \begin{matrix} 0 \\ b \end{matrix}; q^2, z \right), \quad (109)$$

with

$$b = q^2 / a_0^2 a_1^2, \quad z = q^2 t^2 / a_1^2. \quad (110)$$

(b)

$$g = \frac{b_0}{a_0 t} \frac{\Psi(a, z)}{\Psi(a, q^2 z)}, \quad f = -a_0 t \frac{\Psi(a, q^2 z)}{\Psi(a, z)}, \quad (111)$$

where

$$\Psi = \Psi(a, z) = {}_1\varphi_1 \left( \begin{matrix} a \\ 0 \end{matrix}; q^2, z \right), \quad (112)$$

with

$$a = a_0^2 t^2, \quad z = q/b_0. \quad (113)$$

### 3.4.2 Data

(1) Riccati equation

(a)

$$\bar{f} = \frac{(a_0^2 a_1^2 / q - a_0^2 q t^2) f - a_0 q t}{a_0 t f + 1}, \quad g = -a_0 a_1 \frac{1}{a_0 t + f}. \quad (114)$$

(b)

$$\bar{g} = -\frac{g - a_0 b_0 t}{a_0 t g - b_0}, \quad f g = -b_0. \quad (115)$$

(2) Three-term and contiguity relations for hypergeometric function

(a) (i) Three-term relation

$$\frac{b}{z} (\Phi(b, q^2 z) - \Phi(b, z)) + \Phi(b, z) + \frac{q^2}{z} (\Phi(b, z/q^2) - \Phi) = 0, \quad (116)$$

where

$$\Phi(b, z) = {}_1\varphi_1 \left( \begin{matrix} 0 \\ b \end{matrix}; q^2, z \right). \quad (117)$$

(ii) Contiguity relations

$$\Phi(b, z) - \frac{b}{q^2} \Phi(b, q^2 z) = \left( 1 - \frac{b}{q^2} \right) \Phi(b/q^2, z), \quad (118)$$

$$\Phi(b, z) - \Phi(b, z/q^2) = \frac{z/q^2}{1-b} \Phi(q^2 b, z). \quad (119)$$

(b) (i) Three-term relation

$$(1-a) \frac{z}{q^2} (\Psi(q^2 a, z) - \Psi(a, z)) - \frac{a z}{q^2} \Psi(a, z) + (\Psi(a/q^2, z) - \Psi(a, z)) = 0, \quad (120)$$

where

$$\Psi(a, z) = {}_1\varphi_1 \left( \begin{matrix} a \\ 0 \end{matrix}; q^2, z \right). \quad (121)$$

(ii) Contiguity relations

$$\Psi(a, z) - a \Psi(a, q^2 z) = (1-a) \Psi(q^2 a, z), \quad (122)$$

$$\Psi(a, z) - \Psi(q, z/q^2) = (1-a) \frac{z}{q^2} \Psi(q^2 a, z). \quad (123)$$

(3) Decoupling factors

(a)

$$H = \frac{1}{q}, \quad K = 1, \quad k = \frac{H}{K} = \frac{1}{q}, \quad \kappa = \frac{a_1}{qt}. \quad (124)$$

(b)

$$H = -\frac{1}{1-a_0^2 t^2}, \quad K = \frac{1}{q} \frac{1}{1-a_0^2 t^2}, \quad k = \frac{H}{K} = q, \quad \kappa = a_0 t. \quad (125)$$

(4) Identification

(a)

$$f = \frac{1}{\kappa} \frac{F}{G}, \quad F \propto \Phi(q^2 b, q^2 z), \quad G \propto \Phi(b, z), \quad (126)$$

with parameters given in eq.(110).

(b)

$$g = \frac{1}{\kappa} \frac{F}{G}, \quad F \propto \Psi(a, z), \quad G \propto \Psi(a, q^2 z), \quad (127)$$

with parameters given in eq.(113).

(5) Gauge factors

(a) Putting

$$F = \theta(q^2 b, q^2 z) \Phi(q^2 b, q^2 z), \quad G = \theta(b, z) \Phi(b, z), \quad (128)$$

we have:

$$\frac{\theta(b, q^2 z)}{\theta(b, z)} = 1, \quad \frac{\theta(q^2 b, z)}{\theta(b, z)} = \frac{\theta(q^2 b, q^2 z)}{\theta(b, z)} = \frac{q}{a_0 a_1 \left(1 - \frac{q^2}{a_0^2 a_1^2}\right)}. \quad (129)$$

(b) Putting

$$F = \theta(a, z) \Psi(a, z), \quad G = \theta(a, q^2 z) \Psi(a, q^2 z), \quad (130)$$

we have:

$$\frac{\theta(q^2 a, z)}{\theta(a, z)} = 1, \quad \frac{\theta(a, q^2 z)}{\theta(a, z)} = \frac{\theta(q^2 a, q^2 z)}{\theta(a, z)} = \frac{1}{b_0}. \quad (131)$$

### 3.5 Case of $(A_1 + A_1')^{(1)}$

#### 3.5.1 Equation and Solution

(1)  $q$ -Painlevé equation [2, 3, 18]

$$(\bar{f}f - 1)(f\underline{f} - 1) = \frac{at^2 f}{f+t}, \quad \bar{t} = qt. \quad (132)$$

(2) Constraint on parameters

$$a = q. \quad (133)$$

(3) Hypergeometric solution

$$f = \frac{\Phi(qt)}{\Phi(t)}, \quad \Phi = {}_1\phi_1 \left( \begin{matrix} 0 \\ -q \end{matrix} ; q, -qt \right). \quad (134)$$

### 3.5.2 Data

(1) Riccati equation

$$\bar{f} = \frac{1}{f} - qt. \quad (135)$$

(2) Three-term relation

$$\Phi(qt) + t\Phi(t) = \Phi(t/q). \quad (136)$$

(3) Identification

$$f = \frac{F}{G}, \quad F = \Phi(qt), \quad G = \Phi(t). \quad (137)$$

We note that there is no need to introduce decoupling and gauge factors.

**Acknowledgment** One of the author(K.K.) would like to thank Professor Nalini Joshi for hospitality during his stay in University of Sydney.

## References

- [1] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada, *Hypergeometric solutions to the  $q$ -Painlevé equations*, Int. Math. Res. Not. **2004** 2497-2521.
- [2] H. Sakai, *Rational surfaces associated with affine root systems and geometry of the Painlevé equations*, Commun. Math. Phys. **220** (2001) 165–229.
- [3] A. Ramani, B. Grammaticos, T. Tamizhmani and K.M. Tamizhmani, *Special function solutions of the discrete Painlevé equations*, Comp. Math. Appl. **42**(2001) 603–614.
- [4] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada,  ${}_{10}E_9$  solution to the elliptic Painlevé equation, J. Phys. A: Math. Gen. **36** (2003) L263–L272.
- [5] H. Sakai, *Casorati determinant solutions for the  $q$ -difference sixth Painlevé equation*, Nonlinearity **11** (1998) 823–833.
- [6] M. Murata, H. Sakai and J. Yoneda, *Riccati solutions of discrete Painlevé equations with Weyl group symmetry of type  $E_8^{(1)}$* , J. Math. Phys. **44** (2003) 1396–1414.
- [7] K. Kajiwara, M. Noumi and Y. Yamada, *A study on the fourth  $q$ -Painlevé equation*, J. Phys. A: Math. Gen. **34** (2001) 8563–8581.
- [8] K. Kajiwara and K. Kimura, *On a  $q$ -difference Painlevé III equation. I: Derivation, symmetry and Riccati type solutions*, J. Nonlin. Math. Phys. **10** (2003) 86–102.
- [9] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, Encyclopedia of Mathematics and its Applications, **35**(1990) (Cambridge: Cambridge University Press).
- [10] D.P. Gupta, M.E.H. Ismail and D.R. Masson, *Contiguous relations, basic hypergeometric functions, and orthogonal polynomials. II. associated big  $q$ -Jacobi polynomials*, J. Math. Anal. Appl. **171**(1992) 477–497.
- [11] M. Ismail and M. Rahman, *The Associated Askey-Wilson Polynomials*, Trans. Amer. Math. Soc. **328**(1991) 201-237.
- [12] Y. Ohta, A. Ramani and B. Grammaticos, *An affine Weyl group approach to the eight-parameter discrete Painlevé equation*, J. Phys. A: Math. Gen. **34** (2001) 10523–10532.
- [13] D.P. Gupta and D.R. Masson, *Contiguous relations, continued fractions and orthogonality*, Trans. Amer. Math. Soc. **350** (1998) 769–808.



- [14] B. Grammaticos, A. Ramani and Y. Ohta, *A unified description of the asymmetric  $q$ - $P_V$  and  $d$ - $P_{IV}$  equations and their Schlesinger transformations*, J. Nonlin. Math. Phys. **2**(2003) 215-228.
- [15] K.M. Tamizhmani, A. Ramanim B. Grammaticos and Y. Ohta, *A study of the discrete  $P_V$  equation: Miura transformation and particular solutions*, Lett. Math. Phys. **38**(1996) 289–296.
- [16] R. Koekoek and R. F. Swarttouw, *The Askey-scheme of hypergeometric orthogonal polynomials and its  $q$ -analogue*, Delft University of Technology, Faculty of Information Technology and Systems, Department of Technical Mathematics and Informatics, Report no. 98-17 (1998).
- [17] K. Kajiwara, *On a  $q$ -difference Painlevé III equation. II: Rational solutions*, J. Nonlin. Math. Phys. **10** (2003) 282–303.
- [18] A. Ramani and B. Grammaticos, *Discrete Painlevé equations: coalescences, limits and degeneracies*, Physica A **228** (1996) 160–171.