

ROUTE TO CHAOS IN THREE-DIMENSIONAL MAPS OF LOGISTIC TYPE.

Danièle Fournier-Prunaret(*), Ricardo Lopez-Ruiz(**), Abdel-Kaddous Taha(*)¹

Abstract : A route to chaos is studied in 3-dimensional maps of logistic type. Mechanisms of period doubling for invariant closed curves (ICC) are found for specific 3-dimensional maps. These bifurcations cannot be observed for ICC in the 2-dimensional case. When the parameter of the system is modified, localized oscillations occur on the ICC that give rise to weakly chaotic rings, then to chaotic attractors, which finally disappear by contact bifurcations. These maps can be considered as models for the symbiotic interaction of three species.

I INTRODUCTION

Many papers have been devoted to the study of two-dimensional coupled logistic maps [1] - [4], [6]. Some of them can be considered as biological models, corresponding to interactions between species [1] - [4]. In this paper, we consider two models of the same kind in the 3-dimensional case. The first one, given by the map T_1 , corresponds to a symbiotic interaction between correlative pairs of species :

$$T_1 \begin{cases} x_{n+1} = \lambda(3y_n + 1)x_n(1 - x_n), \\ y_{n+1} = \lambda(3z_n + 1)y_n(1 - y_n), \\ z_{n+1} = \lambda(3x_n + 1)z_n(1 - z_n), \end{cases} \quad (1)$$

where λ is a real positive parameter and (x,y,z) represent the species populations.

The second one, given by the map T_2 :

$$T_2 \begin{cases} x_{n+1} = \lambda(x_n + y_n + z_n + 1)x_n(1 - x_n), \\ y_{n+1} = \lambda(x_n + y_n + z_n + 1)y_n(1 - y_n), \\ z_{n+1} = \lambda(x_n + y_n + z_n + 1)z_n(1 - z_n), \end{cases} \quad (2)$$

corresponds to a global symbiotic interaction among the 3 species (see [1] for more informations on the model), λ is also a real positive parameter and (x,y,z) represent the species populations

Our aim is to study the routes to the chaos in such 3-dimensional models when the parameter λ is modified. Then, we consider the existence of invariant closed curves and their evolutions towards chaotic attractors.

II PERIOD DOUBLING OF INVARIANT CLOSED CURVES

Let consider first the map T_1 . There exists a fixed point P, whose coordinates are given by :

$$\frac{1}{3} \left(1 + \sqrt{\frac{4\lambda - 3}{\lambda}}\right) (1,1,1). \quad (3)$$

When λ belongs to $[0.75, 0.895]$, this point is stable ; when $\lambda=0.895$ it undergoes a Neïmark-Hopf bifurcation and gives rise to a stable invariant closed curve (ICC) C_1 (Figure 1a). C_1 undergoes a kind of period doubling (the doubling of a toric surface in 4 dimensions) when $\lambda=0.9705$ (Figure 1b-c), the ICC becoming doubled with a curle. Such phenomenon has been observed in [7][8], also for 3-dimensional maps. This kind of phenomena does not exist in 2-dimensional case. Then a second period doubling occurs when $\lambda=0.9985$, the ICC C_1 becomes formed with 4 curles (Figures 1d-1e). Two other ICC C_2 (Figures 3a-b) and

¹ (*)Danièle Fournier-Prunaret, Abdel-Kaddous Taha, SYD - LESIA - INSA - Complexe scientifique de Rangueil, 31077 Toulouse, France.

E-mail : daniele.fournier@insa-toulouse.fr

(**)Ricardo Lopez-Ruiz, Department of Computer Science and BIFI, Facultad de Ciencias-Edificio B, Universidad de Zaragoza, 50009 - Zaragoza (Spain).

C_3 (Figures 5a-b) exist for other parameter values and also undergo such period doubling (Figures 3c-d-e for C_2 and Figure 5c for C_3).

Concerning the map T_2 , the same phenomena can be observed for an order 2 cyclic ICC (Figures 7a-b).




Figure 1a : First invariant closed curve (ICC) C_1 of map T_1




Figure 1b : Period doubling of ICC C_1 .




Figure 1c : Projection of ICC C_1 on (x,y) plane.




Figure 1d : Projection of ICC C_1 on (x,y) plane after second period doubling.




Figure 1e : ICC C_1 after second period doubling.




Figure 2a : Frequency locking on ICC C_1 .




Figure 2b : Enlargement of Figure 2a.




Figure 2c : ICC C_1 becomes a weakly chaotic ring (WCR).




Figure 2d : Enlargement of Figure 2c.




Figure 3a : Second ICC C_2 of map T_1 .




Figure 3b : Projection of ICC C_2 on (x,y) plane.




Figure 3c : Projection of ICC C_2 on (x,y) plane after period doubling.




Figure 3d : Projection of ICC C_2 on (x,y) plane after second period doubling.




Figure 3e : ICC C_2 after second period doubling.




Figure 4a : Projection of ICC C_2 on (x,y) plane after it becomes a WCR.




Figure 4b : Enlargement of Figure 4a.




Figure 5a : Third ICC C_3 of map T_1 .




Figure 5b : Projection of ICC C_3 on (x,y) plane.




Figure 5c : Enlargement of Figure 5b, one can see a period doubling which cannot be easily seen on Figure 5b.




Figure 6a : Projection of ICC C_3 on (x,y) plane after it becomes a WCR.




Figure 6b : Enlargement of Figure 6a.




Figure 6c : The WCR of Figure 6a becomes a chaotic attractor.





Figure 7b : Map T_2 , period doubling of order 2 cyclic ICC.




Figure 8a : The two cyclic ICC of Figures 7 become cyclic chaotic attractors.




Figure 8b : After contact bifurcation, the two cyclic chaotic attractors of Figure 8a become a single one attractor.

III FROM WEAKLY CHAOTIC RINGS TO CHAOS

After the phenomena of period doubling for the different ICC of the two maps T_1 and T_2 , bifurcations analogous to those occurring in the 2-dimensional case, can be observed. Frequency lockings (Figures 2a-b for C_1) and appearance of weakly chaotic rings (WCR) [5] (Figures 2c-d for ICC C_1 , Figures 4a-b for ICC C_2 and Figures 6a-b for ICC C_3 of map T_1). Some “chaotic pockets” appear on parts of curves (Figures 2d & 4b) as it has also been observed for the 2-dimensional case [4].

Then WCR change to chaotic attractors (Figures 6c for ICC C_3 of T_1 and 8a for the order 2 cyclic ICC of T_2).

IV MULTISTABILITY

In both cases (T_1 and T_2), phenomena of multistability can be obtained.

For T_1 , the ICC C_1 coexists with an order 3 cyclic ICC (Figure 9a), which becomes an order 3 WCR (Figures 9b-c) before disappearing, as also observed in 2-dimensional case [4]. The ICC C_1 coexists with the ICC C_2 (Figure 10a) and with the chaotic attractor C_2 becomes (Figure 10b). For other parameter value, the ICC C_1 coexists with the ICC C_3 (Figure 11). ICC C_2 and C_3 never coexist.

For map T_2 , the two order 2 cyclic ICC coexist before becoming two order 2 cyclic chaotic attractors, as observed in previous paragraph (Figures 7a-b & 8a). Then, by contact bifurcations of heteroclinic type, the two chaotic attractors become a single one (Figure 8b).




Figure 9a : Map T_1 , coexistence of ICC C_1 and of an order 3 cyclic ICC.




Figure 9b : Map T_1 , the order 3 cyclic ICC becomes a WCR.




Figure 9c : Projection of ICC of Figure 9b on (x,y) plane.




Figure 10a : Map T_1 , coexistence of ICC C_1 and C_2 .




Figure 10b : Map T_1 , coexistence of ICC C_1 and chaotic attractor C_2 .




Figure 11 : Map T_1 , coexistence of chaotic attractor C_1 and WCR C_2 .

V DISAPPEARANCE OF ATTRACTORS

As it is usual in 2-dimensional case, chaotic attractors disappear for 3-dimensional maps T_1 and T_2 . One can observe two kinds of such bifurcations. The first one corresponds to the disappearance of one of both attractors by a contact bifurcation between the attractor and its basin boundary, the second attractor remains.




Figure 12 : Map T_1 , the attractor C_3 disappears after contact bifurcation. One can see a chaotic transient close to the bifurcation before joining the attractor C_1 .




Figure 13 : Map T_1 , the attractor C_2 disappears after contact bifurcation. One can see a chaotic transient close to the bifurcation before joining the attractor C_1 .




Figure 14 : Map T_1 , the attractor C_1 is going to disappear after contact bifurcation.




Figure 15 : Map T_2 , the chaotic attractor is going to disappear after contact bifurcation

This is the case for C_3 and C_2 ; the attractor C_1 only remains in both cases and the disappearance of C_2 and C_3 gives rise to a chaotic transient before joining the remaining attractor (Figures 12 - 13).

The second bifurcation gives rise to global instability of the modelled system, it also corresponds to a contact bifurcation between the attractor and its basin boundary; for T_1 , the chaotic attractor disappears when $\lambda=1.0201$ and for T_2 , when $\lambda=1.1805$ (Figures 14-15) [5].

VI CONCLUSION

Two 3-dimensional maps of logistic type involving a mutualistic coupling among the variables have been studied. Routes to chaos with specificities to 3-dimensional case regarding the 2-dimensional case have been observed. The most interesting phenomenon, which seems very usual in 3-dimensional case, is period doubling of ICC. Then WCR giving rise to chaotic attractors have also been observed, as contact bifurcations leading to attractor disappearance.

To complete such work, it would be of most interest to introduce studies with critical manifolds, as it usually done in the 2-dimensional case. Phenomena of multistability have been observed, so the study of basins has to be introduced, but basins are much more complicated to obtain and to analyze in the 3-dimensional case. As maps T_1 and T_2 can represent the evolution of species population, it would be also interesting to explain all this behaviour in such a context.

REFERENCES

- [1] Lopez-Ruiz R., Fournier-Prunaret D., "Complex behaviour in a discrete coupled logistic model for the symbiotic interaction of two species", *Mathematical Bioscience and Engineering*, vol. 1, n° 2, (2004), 307 –324.
- [2] Lopez-Ruiz R., Fournier-Prunaret D., "Indirect allee effect, bistability and chaotic oscillations in a predator-prey discrete model of logistic type", *Chaos, Solitons and Fractals*, 24 (2005), 85-101.
- [3] Lopez-Ruiz R., Fournier-Prunaret D. , "Complex patterns on the plane : different types of basin fractalization in a two-dimensional mapping", *International Journal of Bifurcations and Chaos*, vol. 13, n° 5, (2003), 287-311.
- [4] Fournier-Prunaret D., Lopez-Ruiz R., "Basin Bifurcation in a 2-dimensional logistic map", *European Conference on Iteration Theory (ECIT)* , Evora, Portugal, Grazer Mathematischen Berichte publisher, Austria (2002).
- [5] C. Mira, L. Gardini, A. Barugola, J.C. Cathala, «Chaotic dynamics in two-dimensional noninvertible maps», World Scientific, Series A, Vol. 20, 636 pages, June 1996.
- [6] L.Gardini - R. Abraham - D. Fournier-Prunaret - R. Record – « A double logistic map ». *International Journal of Bifurcation and Chaos in Applied Sciences and Engineering*, vol.4, n°1, (1994), 145-176.
- [7] L. Gardini, R. Lupini, C.Mamana, M.G.Messia, « Bifurcations and transitions to chaos in the three-dimensional Lotka-Volterra map ». *SIAM J. Appl. Math.*, vol. 47, n°3, (1987), 455-482.
- [8] F.Argoul, A. Arneodo, "From quasiperiodicity to chaos : an unstable scenario via period-doubling bifurcations of tori". *J. Mécanique Théor. Appl.*, n° spécial, (1984), 241-288.