# PHASE SPACE MODELS FOR STOCHASTIC NONLINEAR PARABOLIC WAVES: WAVE SPREAD AND SINGULARITY

ALBERT C. FANNJIANG

ABSTRACT. We derive several kinetic equations to model the large scale, low Fresnel number behavior of the nonlinear Schrödinger (NLS) equation with a rapidly fluctuating random potential. There are three types of kinetic equations the longitudinal, the transverse and the longitudinal with friction. For these nonlinear kinetic equations we address two problems: the rate of dispersion and the singularity formation.

For the problem of dispersion, we show that the kinetic equations of the longitudinal type produce the cubic-in-time law, that the transverse type produce the quadratic-in-time law and that the one with friction produces the linear-in-time law for the variance prior to any singularity.

For the problem of singularity, we show that the singularity and blow-up conditions in the transverse case remain the same as those for the homogeneous NLS equation with critical or supercritical self-focusing nonlinearity, but they have changed in the longitudinal case and in the frictional case due to the evolution of the Hamiltonian.

# 1. INTRODUCTION

In this paper we consider nonlinear Schrödinger (NLS) equation with a random potential

(1) 
$$i\frac{\partial}{\partial z}\Psi(z,\mathbf{x}) + \frac{\gamma}{2}\Delta_{\mathbf{x}}\Psi(z,\mathbf{x}) + \gamma^{-1}g|\Psi|^{2\sigma}\Psi(z,\mathbf{x}) + \mu V(zL_z,\mathbf{x}L_x)\Psi(z,\mathbf{x}) = 0,$$
$$\mathbf{x} \in \mathbb{R}^d, \quad \sigma > 0$$

where  $\gamma$  is the Fresnel number (defined below),  $\mu$  is the linear coupling coefficient for the random potential V, which is rescaled by two large parameters  $L_z$  and  $L_x$ , and g is nonlinear coupling coefficient with g > 0 representing the self-focusing case and g < 0 the self-defocusing case. Here  $\sigma$  is a positive constant and  $\sigma = 1$  corresponds to the *cubic* NLS equation. We are particularly interested in the regime of low Fresnel number  $\gamma \ll 1$  with a rapidly oscillating potential  $L_x, L_z, \mu \gg 1$ .

Finite-time blow-up or wave collapse is a well-known effect for the self-focusing, (super)-critical  $(d\sigma \geq 2)$  NLS equation without a random potential [23]. In this case the nonlinear focusing effect dominates over the linear diffraction effect of  $i\Delta$ . The question remains whether the added random potential and its scattering effect would prevent the formation of singularity. Given the wide range of scales present in this problem the numerical simulation as well as theoretical analysis is undoubtedly extremely challenging. Although current numerical results, e.g. [5], have already indicated that a white-noise-in-z potential without the feature of rapid fluctuation in **x** does not prevent blow-up. Indeed, without the self-averaging effect of a **x**-rapidly fluctuating potential the large z-fluctuations in the white-noise potential may drive the system to a state with low, negative Hamiltonian, thus developing singularities in very short time. These singularities in our view are the small scale, randomly fluctuating singularities which may be delayed or removed in the presence of a **x**-rapidly fluctuating potential.

To capture the robust, large scale blow-ups we derive several phase-space model equations corresponding to different scalings (in  $\gamma, \mu, L_x, L_z$ ). We use these kinetic equations to elucidate the

Department of Mathematics, University of California at Davis, Davis, CA 95616, E-mail: cafannjiang@ucdavis.edu. Research partially supported by the Centennial Fellowship from the American Mathematical Society, the UC Davis Chancellor's Fellowship and U.S. National Science Foundation grant DMS 0306659.

problems of singularity formation and rate of dispersion. The main ingredient of our analysis is the variance identity for these phase space transport equations. In general, we show that random scattering changes the blow-up conditions for the homogeneous NLS eq. but does not prevent singularity formation on the large scales (see Conclusion for the discussion and summary of our results). We also give various upper bounds for the blow-up time.

As a simple by-product of the variance identity, we also derive various bounds on the rate of dispersion for random Schrödinger waves, including exact expressions in the critical case,  $d\sigma = 2$ . We show that depending on the random scattering mechanism, one can have the cubic  $(z^3)$ , the quadratic  $(z^2)$  and the linear (z) rate of dispersion prior to singularity, if there is any. To our knowledge, these are significant improvements over current results (e.g. [3], [24], [7]) which are mostly for the linear problem.

The cubic NLS equation is a model equation in nonlinear optics, describing the spatial distribution of the stationary electromagnetic field in a nonlinear medium with the Kerr effect [20], [15]. Here let us give a brief derivation of the NLS equation with a random potential.

The electric field E with a fixed polarization in a lossless medium satisfies the Helmholtz equation

$$\nabla_{\vec{\mathbf{x}}}^2 E(\vec{\mathbf{x}}) + k^2 n^2 E(\vec{\mathbf{x}}) = 0, \quad \vec{\mathbf{x}} \in \mathbb{R}^3$$

where k the wavenumber and  $n(\vec{\mathbf{x}}, |E|^2) > 0$  is the refractive index of the medium. For a weakly nonlinear and weakly fluctuating inhomogeneous medium we write

$$n(\vec{\mathbf{x}}, |E|^2) = \bar{n}(1 + \tilde{n}(\vec{\mathbf{x}})) + n_2|E|^2$$

with  $\tilde{n} \ll 1$  and the Kerr coefficient  $n_2$  being small. Consider the approximate equation

(2) 
$$\nabla_{\vec{\mathbf{x}}}^2 E(\vec{\mathbf{x}}) + k^2 \bar{n}^2 E(\vec{\mathbf{x}}) = -2\bar{n}^2 \tilde{n}(\vec{\mathbf{x}}) E(\vec{\mathbf{x}}) - 2\bar{n}n_2 |E|^2 E(\vec{\mathbf{x}}).$$

The electric field is taken as

$$E(z, \mathbf{x}) = \Psi(z, \mathbf{x}) \exp(i\bar{n}kz), \quad \vec{\mathbf{x}} = (z, \mathbf{x}) \in \mathbb{R}^3$$

where z is the coordinate in the direction of propagation and  $\mathbf{x}$  is the transverse coordinates. The modulation  $\Psi$  is assumed to be slowly varying on the scale  $k^{-1}$ .

Substituting the ansatz into eq. (2) and making the parabolic approximation we obtain for the modulation  $\Psi$  the equation

(3) 
$$ik\bar{n}\frac{\partial}{\partial z}\Psi(z,\mathbf{x}) + \frac{1}{2}\nabla_{\mathbf{x}}^{2}\Psi(z,\mathbf{x}) + k^{2}\bar{n}n_{2}|\Psi|^{2}\Psi(z,\mathbf{x}) + k^{2}\bar{n}^{2}\tilde{n}(z,\mathbf{x})\Psi(z,\mathbf{x}) = 0.$$

Let us write

$$\tilde{n}(z, \mathbf{x}) = \nu V(z, \mathbf{x})$$

where  $\nu \ll 1$  is the standard deviation of  $\tilde{n}$  and  $V(z, \mathbf{x})$  is a zero-mean, z-stationary, **x**-homogeneous random field. Let  $L_x, L_z \gg 1$  be the two length scales of the wave beam in the transverse and longitudinal directions, respectively and

$$\tilde{\mathbf{x}} = \mathbf{x}/L_x, \quad \tilde{z} = z/L_z,$$

be the rescaled variables. Let us introduce the dimensionless parameters

$$\mu = k\bar{n}L_z\nu, \quad \gamma = \frac{L_z}{k\bar{n}L_x^2}, \quad g = L_z n_2 k\gamma.$$

Writing the wave field  $\Psi$  in the new variables  $\tilde{\mathbf{x}}, \tilde{z}$ , dropping the tilde and rescaling we obtain (1) with  $\sigma = 1$ .

The critical case of the cubic NLS with d = 2 and g > 0 is our primary example; another is the sub-critical case of a planar waveguide (thin film) with d = 1. The opposite case of a self-defocusing Kerr nonlinearity occurs in semiconductor waveguide [19], [18]. Similar equations also arise in many other contexts such as the Langmuir wave in plasma and Bose-Einstein condensation, see [23] and

references therein. Therefore, we will formulate the results for the general power-law NLS eq. (1) with transverse dimension  $d \ge 1$ . Because the variable z is time-like we will refer to it as "time" occasionally, especially in the discussion of finite-time singularity.

# 2. WIGNER DISTRIBUTION AND WIGNER-MOYAL EQUATION

We consider several families of scaling limits, as  $\gamma \to 0, L_z, L_x \to \infty$ , which are first distinguished by whether

(4) 
$$\theta \equiv \gamma L_x \to 0 \quad \text{or} \quad \theta = 1.$$

This, of course, is not sufficient to ensure the existence of scaling limit until we specify the strength of  $\mu$ .

Our phase-space model equations for the low Fresnel number regime is based on the Wigner equations. We consider the Wigner distribution of the form

(5) 
$$W(\mathbf{x}, \mathbf{p}) = \frac{1}{(2\pi)^d} \int e^{-i\mathbf{p}\cdot\mathbf{y}} \Psi\left(z, \mathbf{x} + \frac{\gamma\mathbf{y}}{2}\right) \Psi^*\left(z, \mathbf{x} - \frac{\gamma\mathbf{y}}{2}\right) d\mathbf{y}$$

The Wigner distribution has many important properties. For instance, it is real and its **p**-integral is the modulus square of the wave function,

(6) 
$$\int_{\mathbb{R}^d} W^{\varepsilon}(\mathbf{x}, \mathbf{p}) d\mathbf{p} = |\Psi(\mathbf{x})|^2 \equiv \rho(\mathbf{x}),$$

so we may think of  $W(\mathbf{x}, \mathbf{p})$  as wave number-resolved mass density. Consequently, the mean  $M_x$ and variance  $V_x$  of **x** are given by, respectively,

$$\bar{\mathbf{x}} = \int \mathbf{x} W(\mathbf{x}, \mathbf{p}) d\mathbf{x} d\mathbf{p},$$

$$V_x = \int |\mathbf{x} - \bar{\mathbf{x}}|^2 W(\mathbf{x}, \mathbf{p}) d\mathbf{x} d\mathbf{p}$$

$$= S_x - \bar{\mathbf{x}}^2$$

where  $S_x$  is the second moment of **x** 

$$S_x = \int |\mathbf{x}|^2 W(\mathbf{x}, \mathbf{p}) d\mathbf{x} d\mathbf{p}.$$

Also, we have that

$$\int_{\mathbb{R}^d} W^{\varepsilon}(\mathbf{x},\mathbf{p}) d\mathbf{x} \ = \ (\frac{2\pi}{\gamma})^d |\widehat{\Psi}^{\varepsilon}|^2(\mathbf{p}/\gamma).$$

and the energy flux

(7) 
$$\frac{1}{2i}(\Psi\nabla\Psi^* - \Psi^*\nabla\Psi) = \int_{\mathbb{R}^d} \mathbf{p} W^{\varepsilon}(\mathbf{x}, \mathbf{p}) d\mathbf{p}$$

so that the mean and variance of  $\mathbf{p}$ ,  $V_p$ , are given by, respectively

$$\bar{\mathbf{p}} = \int \mathbf{p} W(\mathbf{x}, \mathbf{p}) d\mathbf{x} d\mathbf{p},$$

$$V_p = \int |\mathbf{p} - \bar{\mathbf{p}}|^2 W(\mathbf{x}, \mathbf{p}) d\mathbf{x} d\mathbf{p} = S_p - \bar{\mathbf{p}}^2$$

where  $S_p$  is the second moment of **p** 

$$S_p = \int |\mathbf{p}|^2 W(\mathbf{x}, \mathbf{p}) d\mathbf{x} d\mathbf{p}.$$

In view of these properties it is tempting to think of the Wigner distribution as a phase-space probability density, which is unfortunately not the case, since it is not everywhere non-negative (it is always real-valued though).

It is straightforward to derive the closed-form equation for the Wigner distribution [9]

(8) 
$$\frac{\partial W}{\partial z} + \mathbf{p} \cdot \nabla_{\mathbf{x}} W + \mathcal{U}_{\gamma} W + \mu \mathcal{V} W = 0,$$

~ \* \* \*

with the Moyal operators

(9) 
$$\mathcal{U}_{\gamma}W(\mathbf{x},\mathbf{p}) = i\int e^{i\mathbf{q}\cdot\mathbf{x}}\gamma^{-1} \left[W(\mathbf{x},\mathbf{p}+\gamma\mathbf{q}/2) - W(\mathbf{x},\mathbf{p}-\gamma\mathbf{q}/2)\right] \widehat{U}(z,\mathbf{q})d\mathbf{q}, \quad U = g|\Psi|^{2\sigma}$$
  
(10) $\mathcal{V}W(z,\mathbf{x},\mathbf{p}) = i\mu\int e^{i\mathbf{q}\cdot\mathbf{x}L_x} \left[W_z(\mathbf{x},\mathbf{p}+\theta\mathbf{q}/2) - W_z(\mathbf{x},\mathbf{p}-\theta\mathbf{q}/2)\right] \widehat{V}(zL_z,d\mathbf{q})$ 

where  $\hat{U}, \hat{V}$  denote the Fourier transforms of U, V, respectively. Formally we see that as  $\gamma \to 0$ 

(11) 
$$\mathcal{U}_{\gamma}W(\mathbf{x},\mathbf{p}) \to \mathcal{U}_{0}W(\mathbf{x},\mathbf{p}) \equiv \nabla_{\mathbf{x}}U(z,\mathbf{x}) \cdot \nabla_{\mathbf{p}}W(\mathbf{x},\mathbf{p}).$$

We use the following definition of the Fourier transform and inversion:

$$\mathcal{F}f(\mathbf{p}) = \frac{1}{(2\pi)^d} \int e^{-i\mathbf{x}\cdot\mathbf{p}} f(\mathbf{x}) d\mathbf{x}$$
$$\mathcal{F}^{-1}g(\mathbf{x}) = \int e^{i\mathbf{p}\cdot\mathbf{x}} g(\mathbf{p}) d\mathbf{p}.$$

We shall refer to eq. (8) as the nonlinear Wigner-Moyal (NWM) equation. We only need to consider the weak formulation

(12) 
$$\frac{\partial}{\partial z} \int \Theta W d\mathbf{x} d\mathbf{p} - \int \mathbf{p} \cdot \nabla_{\mathbf{x}} \Theta W d\mathbf{x} d\mathbf{p} - \int \mathcal{U}_{\gamma} \Theta W d\mathbf{x} d\mathbf{p} - \mu \int \mathcal{V} \Theta W d\mathbf{x} d\mathbf{p} = 0$$

for smooth, rapidly decaying test functions  $\Theta$ .

One advantage of working with eq. (8) is that one can use it to evolve the mixed-state initial condition, instead of the pure-state one given in (8). This is important in the context of modeling quantum open systems. A mixed state Wigner distribution is a convex combination of the pure-state Wigner distributions (5) described as follows.

Let  $\{\Psi_{\alpha}\}$  be a family of  $L^2$  functions parametrized by  $\alpha$  which is weighted by a probability measure  $P(d\alpha)$ . Denote the pure-state Wigner distribution (8) by  $W[\Psi]$ . A mixed-state Wigner distribution is given by

(13) 
$$\int W[\Psi_{\alpha}]P(d\alpha)$$

The limits as  $\gamma \to 0$  of the mixed state Wigner distributions constitute the so called Wigner measures which are always positive [12], [13], [17]. Evolution by eq. (12) preserves the form (13). In particular, for such initial data we have

(14) 
$$\int W(z, \mathbf{x}, \mathbf{p}) d\mathbf{p} \geq 0, \quad \forall \mathbf{x} \in \mathbb{R}^d$$

(15) 
$$\int W(z, \mathbf{x}, \mathbf{p}) d\mathbf{x} \geq 0, \quad \forall \mathbf{p} \in \mathbb{R}^d$$

Multiplying (12) by W and integrating by parts we also see that the evolution preserves the  $L^2(\mathbb{R}^{2d})$ -norm of W, i.e.

$$\frac{\partial}{\partial z} \int |W|^2 d\mathbf{x} d\mathbf{p} = 0$$

Let us be more explicit about the random potential  $V(z, \mathbf{x})$ . We assume that z-stationary, x-homogeneous random field V is square-integrable and admits the spectral representation

$$V(z, \mathbf{x}) = \int \exp{(i\mathbf{p} \cdot \mathbf{x})} \hat{V}(z, d\mathbf{p})$$

with the z-stationary spectral measure  $\hat{V}(z, \cdot)$  satisfying

$$\mathbb{E}[\hat{V}(z, d\mathbf{p})\hat{V}(z, d\mathbf{q})] = \delta(\mathbf{p} + \mathbf{q})\Phi_0(\mathbf{p})d\mathbf{p}d\mathbf{q}.$$

Here  $\mathbb{E}$  denotes the ensemble expectation and the function  $\Phi_0(\mathbf{p})$  is the transverse power spectrum density which is assumed to be rapidly decaying as  $|\mathbf{p}| \to 0$  or  $|\mathbf{p}| \to \infty$ . The transverse power spectrum density is related to the full power spectrum density  $\Phi(w, \mathbf{p})$  in the following way

$$\Phi_0(\mathbf{p}) = \int \Phi(w, \mathbf{p}) dw$$

with  $\Phi(\mathbf{k})$  satisfying  $\Phi(\mathbf{k}) = \Phi(-\mathbf{k}), \forall \mathbf{k} = (w, \mathbf{p}) \in \mathbb{R}^{d+1}$ . Without loss of generality we will also assume

(16) 
$$\Phi(w,\mathbf{p}) = \Phi(-w,\mathbf{p}) = \Phi(w,-\mathbf{p}) = \Phi(-w,-\mathbf{p}).$$

#### 3. Phase space transport equations

3.1. Linear kinetic equations. First we summarize what has been established in the linear case when  $U(z, \mathbf{x})$  is a given function. In this case we refer to eq. (8) with a given U as the linear Wigner-Moyal (LWM) equation. The principal feature of the scaling is the separation of scales in the given potential U and the random potential V as they appear in the LWM equation, c.f. (9) and (10). Under pretty general conditions (the most important of which being the integrability of the maximum correlation coefficient of  $V(z, \cdot)$  as **x**-homogeneous-field-valued z-stationary process) one can prove that, as  $\gamma \to 0$  and  $\mu$ ,  $L_x$ ,  $L_z \to \infty$ , the weak solution of the LWN equation converges in law to the weak solution of the linear Boltzmann (LB) equation or the linear Fokker-Planck (LFP) equation, described below, depending on whether  $\theta$  also tends to zero or not [9],[10] (see also [2]).

## Linear Boltzmann equation $(\theta = 1)$ :

(17) 
$$\frac{\partial}{\partial z} W(z, \mathbf{x}, \mathbf{p}) + \mathbf{p} \cdot \nabla_{\mathbf{x}} W(z, \mathbf{x}, \mathbf{p}) + \nabla_{\mathbf{x}} U(z, \mathbf{x}) \cdot \nabla_{\mathbf{p}} W(z, \mathbf{x}, \mathbf{p})$$
$$= 2\pi \int K(\mathbf{p}, \mathbf{q}) [W(z, \mathbf{x}, \mathbf{q}) - W(z, \mathbf{x}, \mathbf{p})] d\mathbf{q}$$

with a nonnegative kernel  $K(\mathbf{p}, \mathbf{q})$  given respectively as follows.

(a): If  $\mu \sim \sqrt{L_z}, L_x \ll L_z$  then

(18) 
$$K(\mathbf{p},\mathbf{q}) = \Phi(0,\mathbf{q}-\mathbf{p})$$

(b): If  $\mu \sim \sqrt{L_x}, L_z \ll L_x \ll L_z^{4/3}, d \ge 3$  then

(19) 
$$K(\mathbf{p},\mathbf{q}) = \delta(\frac{|\mathbf{q}|^2 - |\mathbf{p}|^2}{2}) \left[ \int \Phi(w,\mathbf{q}-\mathbf{p}) dw \right].$$

(c): If  $\mu \sim \sqrt{L_z}, L_x \sim L_z$  then

(20) 
$$K(\mathbf{p},\mathbf{q}) = \Phi(\frac{|\mathbf{q}|^2 - |\mathbf{p}|^2}{2}, \mathbf{q} - \mathbf{p}).$$

Linear Fokker-Planck equation  $(\theta \rightarrow 0)$ :

(21) 
$$\frac{\partial}{\partial z} W(z, \mathbf{x}, \mathbf{p}) + \mathbf{p} \cdot \nabla_{\mathbf{x}} W(z, \mathbf{x}, \mathbf{p}) + \nabla_{\mathbf{x}} U(z, \mathbf{x}) \cdot \nabla_{\mathbf{p}} W(z, \mathbf{x}, \mathbf{p}) \\ = \nabla_{\mathbf{p}} \cdot \mathbf{D} \nabla_{\mathbf{p}} W(z, \mathbf{x}, \mathbf{p})$$

with a symmetric, nonnegative-definite matrix **D** given as follows.

(a): If  $\mu \sim \theta^{-1} \sqrt{L_z}$ ,  $L_x \ll L_z$  then

(22) 
$$\mathbf{D} = \pi \int \Phi(0, \mathbf{q}) \mathbf{q} \otimes \mathbf{q} d\mathbf{q}.$$

(b): If 
$$\mu \sim \theta^{-1} \sqrt{L_x}, L_z \ll L_x \ll L_z^{4/3}, d \ge 3$$
 then

(23) 
$$\mathbf{D}(\mathbf{p}) = \pi |\mathbf{p}|^{-1} \int_{\mathbf{p} \cdot \mathbf{p}_{\perp} = 0} \int \Phi(w, \mathbf{p}_{\perp}) dw \ \mathbf{p}_{\perp} \otimes \mathbf{p}_{\perp} d\mathbf{p}_{\perp}.$$

(c): If  $\mu \sim \theta^{-1} \sqrt{L_z}, L_x \sim L_z$  then

(24) 
$$\mathbf{D}(\mathbf{p}) = \pi \int \Phi(\mathbf{p} \cdot \mathbf{q}, \mathbf{q}) \mathbf{q} \otimes \mathbf{q} d\mathbf{q}$$

We shall use  $\mathcal{L}$  to denote either the scattering operator on the right side of eq. (17) or the diffusion operator on the right side of eq. (21). The self-adjoint operator  $\mathcal{L}$  is non-positive definite and represents various decoherence effects due to random fluctuations of the medium, see (27). The evolution by Eq. (17) or (21) preserves the mixed-state structure (13) and hence the positivity (14), (15).

We shall refer to regime (a), for either (17) or (21), as the *longitudinal* case and regime (b) as the *transverse* case because the longitudinal and the transverse scales are dominant, respectively. Regime (c) is the borderline case. We will consider hereafter the longitudinal and transverse cases only.

We note that the additional restrictions of  $L_x \ll 4/3L_z$  and  $d \ge 3$  in regime (b) are due to technical reasons. We believe that the results should hold for any  $L_z \ll L_x$ ,  $d \ge 2$  along with the other assumptions used in proving the above scaling limits (see [9], [10] for details). The mean field result for a Gaussian potential with  $L_z = 0$  and  $d \ge 3$  has been established previously [22], [8] (see also [21]).

3.2. Nonlinear kinetic equation. When  $U = g|\Psi|^{2\sigma}$ , the convergence of the above scaling limits is not known. But because the phase of  $\Psi$  is canceled in the expression of U one can reasonably expect that, at least before any singularity formation, U has significantly less oscillation than  $\Psi$ and  $V(zL_z, \mathbf{x}L_x)$  as appearing in the NLS equation (1). The property of separation of scales in U and  $\Psi$  holds in the linear case and will be assumed to hold prior to singularity formation.

Hence we will assume below the validity of the above scaling limits for the nonlinear case and use the kinetic equation (17) or (21) with  $U = q |\Psi|^{2\sigma}$  to investigate the issue of random diffraction and singularity formation in the case of self-focusing (q > 0), (super)-critical  $(d\sigma \ge 2)$  nonlinearity as well as the question of rate of spread in all situations. Such a model with the kernel (19) was considered in [11]. The Fokker-Planck equation with the diffusion coefficient (22) is an example of those which describe open quantum systems in contact with a heat bath of linear oscillators in thermal equilibrium (see, e.g., [4], [6], [1], [25]). In this context, z would be the physical time,  $\mathbf{x}$ the physical coordinates and **p** the momentum of the quantum particle. We will discuss the master equation for the open system including dissipation mechanism in the final section.

Let us state the nonlinear kinetic equation which we will analyze subsequently:

(25) 
$$\frac{\partial}{\partial z}W(z,\mathbf{x},\mathbf{p}) + \mathbf{p} \cdot \nabla_{\mathbf{x}}W(z,\mathbf{x},\mathbf{p}) + \mathcal{U}_0W(z,\mathbf{x},\mathbf{p}) = \mathcal{L}W(z,\mathbf{x},\mathbf{p})$$

where  $\mathcal{U}_0$  is given by (11) and  $\mathcal{L}$  is either the linear scattering operator

$$\mathcal{L}W = 2\pi \int K(\mathbf{p}, \mathbf{q}) [W(z, \mathbf{x}, \mathbf{q}) - W(z, \mathbf{x}, \mathbf{p})] d\mathbf{q}$$

or the linear diffusion operator

$$\mathcal{L}W = \nabla_{\mathbf{p}} \cdot \mathbf{D} \nabla_{\mathbf{p}} W(z, \mathbf{x}, \mathbf{p}).$$

The weak formulation of (25) is given by

(26) 
$$\frac{\partial}{\partial z} \int \Theta W d\mathbf{x} d\mathbf{p} - \int \mathbf{p} \cdot \nabla_{\mathbf{x}} \Theta W d\mathbf{x} d\mathbf{p} - \int \nabla_{\mathbf{x}} U \cdot \nabla_{\mathbf{p}} \Theta W d\mathbf{x} d\mathbf{p} - \int \mathcal{L} \Theta W d\mathbf{x} d\mathbf{p} = 0$$

for smooth, rapidly decaying test functions  $\Theta$ . The evolution preserves the positivity of the initial condition which is consistent with the fact that the Wigner measures are positive and will be assumed subsequently (see the comment following (13)).

The nonlinear kinetic eq. (26) preserves the total mass, i.e.

$$\frac{\partial}{\partial z}N = 0, \quad N = \int W(z, \mathbf{x}, \mathbf{p}) d\mathbf{x} d\mathbf{p}$$

but in general decreases the  $L^2$ -norm

(27) 
$$\frac{\partial}{\partial z} \int |W|^2(z, \mathbf{x}, \mathbf{p}) d\mathbf{x} d\mathbf{p} = \int W \mathcal{L} W d\mathbf{x} d\mathbf{p} \le 0$$

because the operator  $\mathcal{L}$  is non-positive definite. The inequality (27) expresses certain irreversibility as a result of the weak convergence of solutions [9]. One can absorb the effect of the total mass Ninto g by the obvious rescaling of W in the eq. (26). Henceforth we assume that

$$N = 1$$

which is the case when in (13)

$$\int |\Psi_{\alpha}(\mathbf{x})|^2 d\mathbf{x} = 1$$

for *P*-almost all  $\alpha$ .

3.3. Local existence. A natural space of initial data and solutions is the space S of the non-negative measures with square integrable density W

$$\int |W|^2 d\mathbf{x} d\mathbf{y} < \infty,$$

finite Dirichlet form

$$-\int W\mathcal{L}Wd\mathbf{x}d\mathbf{p}<\infty$$

and finite, positive variances

$$V_x = \int |\mathbf{x} - \bar{\mathbf{x}}|^2 W d\mathbf{x} d\mathbf{p} = S_x - |\bar{\mathbf{x}}|^2 \in (0, \infty), \quad V_p = \int |\mathbf{p} - \bar{\mathbf{p}}|^2 W d\mathbf{x} d\mathbf{p} = S_p - |\bar{\mathbf{p}}|^2 \in (0, \infty)$$

where  $S_x, S_p$  are the second moments

(28) 
$$S_x = \int |\mathbf{x}|^2 W d\mathbf{x} d\mathbf{p} \in (0,\infty), \quad S_p = \int |\mathbf{p}|^2 W d\mathbf{x} d\mathbf{p} \in (0,\infty)$$

In addition, we shall also assume that the initial data have a finite Hamiltonian H

$$H = \frac{1}{2}S_p - \frac{g}{\sigma+1}\int \rho^{\sigma+1}d\mathbf{x} \in (-\infty,\infty), \quad \rho(\mathbf{x}) = \int W d\mathbf{p}$$

the first term of which is the kinetic energy and the last two terms are the potential energy. A finite Hamiltonian and a finite variance  $V_p$  then imply

(29) 
$$\int \rho^{\sigma+1} d\mathbf{x} < \infty,$$

namely a finite potential energy.

We recall that for the NLS eq. without a random potential and the scaling (4) the local existence in the analogous space of functions (i.e.  $\Psi, \mathbf{x}\Psi, \nabla\Psi, |\Psi|^{\sigma+1} \in L^2(\mathbb{R}^d)$ ) holds under the additional assumption

$$0 \le \sigma < \frac{2}{d-2}, \quad \text{for } d \ge 3$$

and  $\sigma \in [0, \infty)$ , for  $d \leq 2$ , see [23] and the references therein.

The local existence for the nonlinear kinetic equation (26) in the space S is assumed throughout the paper and will be addressed elsewhere.

3.4. Energy law. Next, we discuss the evolution of the Hamiltonian H and the variance  $V_x$ . Let us first note the result of  $\mathcal{U}_{\gamma}$  applied to the quadratic polynomials.

(30) 
$$\mathcal{U}_{\gamma}\mathbf{x} = 0$$

(31) 
$$\mathcal{U}_{\gamma}\mathbf{p} = i\int e^{i\mathbf{q}\cdot\mathbf{x}}\mathbf{q}\hat{U}(d\mathbf{q}) = \nabla_{\mathbf{x}}U$$

(32) 
$$\mathcal{U}_{\gamma}|\mathbf{x}|^2 = 0$$

(33) 
$$\mathcal{U}_{\gamma}\mathbf{x}\cdot\mathbf{p} = i\int e^{i\mathbf{q}\cdot\mathbf{x}}\mathbf{x}\cdot\mathbf{q}\hat{U}(d\mathbf{q}) = \mathbf{x}\cdot\nabla_{\mathbf{x}}U$$

(34) 
$$\mathcal{U}_{\gamma}|\mathbf{p}|^2 = i \int e^{i\mathbf{q}\cdot\mathbf{x}} 2\mathbf{p} \cdot \mathbf{q}\hat{U}(d\mathbf{q}) = 2\mathbf{p} \cdot \nabla_{\mathbf{x}} U.$$

It is noteworthy that the results of the calculation are independent of  $\gamma \ge 0$  and identical to those for  $\gamma = 0$  (see more on this in Conclusion).

Consider the mean dynamics for

(35) 
$$\bar{\mathbf{x}} = \int \mathbf{x} W d\mathbf{x} d\mathbf{p}, \quad \bar{\mathbf{p}} = \int \mathbf{p} W d\mathbf{x} d\mathbf{p}$$

with the mean Hamiltonian defined as

(36) 
$$\bar{H} = \frac{1}{2} |\bar{\mathbf{p}}|^2.$$

Using the above and integrating by parts we obtain the following

$$\begin{aligned} \frac{\partial}{\partial z} \bar{\mathbf{p}} &= \int \mathcal{L}^* \mathbf{p} W d\mathbf{x} d\mathbf{p} \\ \frac{\partial}{\partial z} V_p &= \int \nabla_{\mathbf{x}} U \cdot \mathbf{p} W d\mathbf{x} d\mathbf{p} + \int \mathcal{L}^* |\mathbf{p}|^2 W d\mathbf{p} d\mathbf{x} \\ \frac{\partial}{\partial z} \frac{g}{\sigma + 1} \int \rho^{\sigma + 1} d\mathbf{x} &= \int \nabla_{\mathbf{x}} U \cdot \mathbf{p} W d\mathbf{p} d\mathbf{x} \end{aligned}$$

and hence

(37) 
$$\frac{\partial}{\partial z}\bar{H} = \bar{\mathbf{p}} \cdot \int \mathcal{L}^* \mathbf{p} W d\mathbf{x} d\mathbf{p}$$

(38) 
$$\frac{\partial}{\partial z}H = \int \mathcal{L}^* |\mathbf{p}|^2 W d\mathbf{p} d\mathbf{x}$$

which will take a more explicit form once we calculate  $\mathcal{L}^*\mathbf{p}, \mathcal{L}^*|\mathbf{p}|^2$  with  $\mathcal{L}^*$  of each case.

As is the case without a random potential and the scaling limit (4) [23], we assume subsequently that the laws (37), (38) for the Hamiltonian holds for the local solutions of eq. (26) in the space S.

In the following sections, we first derive the variance identity for eq. (26) in the longitudinal case (regime (a)), then the transverse case (regime (b)) and finally the case with friction. As is the case without a random potential and the scaling limit (4) [23], we assume throughout the paper that the variance identity derived below holds true for the maximally extended local solutions of eq. (26) in the space S. Let  $[0, z_*)$  be the interval on which the maximally extended local solution is defined. When  $z_* = \infty$  then the local solutions become global solutions; when  $z_* < \infty$  then the solutions are said to develop finite time singularity. Our main goal then is to analyze what these formally derived relations tell us about singularity formation and rate of spread of low Fresnel number nonlinear parabolic waves in random media.

Before concluding this section let us state some elementary inequalities which will be useful later. An application of the Cauchy-Schwartz inequality and the marginal positivity (14), (15) to the first moments  $\bar{\mathbf{x}}, \bar{\mathbf{p}}$  then leads to

(39) 
$$|\bar{\mathbf{x}}|^2 \leq S_x, \quad |\bar{\mathbf{p}}|^2 \leq S_p.$$

Furthermore, by using the mixed-state structure (13) in estimating the cross moment

(40) 
$$S_{xp} = \int \mathbf{x} \cdot \mathbf{p} W d\mathbf{x} d\mathbf{p}$$

one deduces that

$$(41) S_{xp}^2 \le S_x S_p < \infty.$$

Likewise the covariance

$$V_{xp} = \int (\mathbf{x} - \bar{\mathbf{x}}) \cdot (\mathbf{p} - \bar{\mathbf{p}}) W d\mathbf{x} d\mathbf{p} = S_{xp} - \bar{\mathbf{x}} \cdot \bar{\mathbf{p}}$$

can be bounded by  $V_x$  and  $V_p$  as

(42)  $V_{xp}^2 \le V_x V_p.$ 

## 4. The longitudinal case

4.1. Variance identity. The variance identity has been long used to derive the wave collapse condition for the NLS without the random potential [23]. Below we reformulate it and extend it to the phase-space models (26).

We have the following simple calculations for  $\mathcal{L}^* = \mathcal{L}$ :

$$\mathcal{L}\mathbf{x} = 0$$
  

$$\mathcal{L}\mathbf{p} = 2\pi \int \Phi(0, \mathbf{p} - \mathbf{q})(\mathbf{p} - \mathbf{q})d\mathbf{q} = 0 \quad (by \ (16))$$
  

$$\mathcal{L}|\mathbf{x}|^2 = 0$$
  

$$\mathcal{L}\mathbf{x} \cdot \mathbf{p} = \mathbf{x} \cdot 2\pi \int \Phi(0, \mathbf{p} - \mathbf{q})(\mathbf{p} - \mathbf{q})d\mathbf{q} = 0 \quad (by \ (16))$$
  

$$\mathcal{L}|\mathbf{p}|^2 = -2\pi \int \Phi(0, \mathbf{p} - \mathbf{q}) \left[|\mathbf{p}|^2 - |\mathbf{q}|^2\right] d\mathbf{q}$$
  

$$= -2\pi \int \Phi(0, \mathbf{q})(2\mathbf{p} - \mathbf{q}) \cdot \mathbf{q} d\mathbf{q}$$
  

$$= 2\pi \int \Phi(0, \mathbf{q})|\mathbf{q}|^2 d\mathbf{q} \equiv R.$$

We have used the linear Boltzmann operator  $\mathcal{L}$  in the above calculation; the same result holds for the linear diffusion operator  $\mathcal{L}$  for which case

$$R = 2 \times \text{trace}(\mathbf{D})$$

where the diffusion matrix  $\mathbf{D}$  is given by (22). We shall use the above identities to perform integrating by parts in the derivation of the variance identity.

The evolution of the mean position  $\bar{\mathbf{x}}$  and momentum  $\bar{\mathbf{p}}$  is then given by

$$\frac{\partial}{\partial z} \bar{\mathbf{x}} = \bar{\mathbf{p}} 
\frac{\partial}{\partial z} \bar{\mathbf{p}} = \frac{g\sigma}{\sigma+1} \int \nabla_{\mathbf{x}} \rho^{\sigma+1} d\mathbf{x} 
= 0,$$

as a consequence the mean Hamiltonian  $\overline{H}$  is invariant. Moreover, the evolution of  $(\overline{\mathbf{x}}, \overline{\mathbf{p}})$  for the autonomous Hamiltonian system of the harmonic oscillator.

The evolution of the variance  $V_x$  is given by

$$\frac{\partial}{\partial z}V_x = 2V_{xp}$$

Differentiating  $S_{xp}$  we obtain

$$\frac{\partial}{\partial z} S_{xp} = S_p - \frac{gd\sigma}{\sigma+1} \int \rho^{\sigma+1} d\mathbf{x} d\mathbf{p}$$

Hence the second derivative of  $V_x$  becomes

(43) 
$$\frac{\partial^2}{\partial z^2} V_x = 4(H - \bar{H}) + \frac{2(2 - d\sigma)g}{\sigma + 1} \int \rho^{\sigma + 1} d\mathbf{x}$$

An alternative expression for the variance identity is

(44) 
$$\frac{\partial^2}{\partial z^2} V_x = 2d\sigma (H - \bar{H}) + (2 - d\sigma) V_p.$$

Both forms (43) and (44) will be used to obtain dispersion estimates below.

4.2. **Dispersion rate.** Although the medium is lossless, reflected in the fact that the total mass N = 1 is conserved, but the Hamiltonian is not conserved by the evolution since the scattering with the random potential is not elastic. Indeed, its rate of change is

(45) 
$$\frac{\partial}{\partial z}H = R, \quad H(z) = H(0) + Rz$$

due to the diffusion-like spread in the momentum **p**.

In the critical case  $d\sigma = 2$ , we have the exact result

$$\frac{\partial^2}{\partial z^2} V_x = 4H - 2|\bar{\mathbf{p}}|^2 = 4(H(0) - \bar{H}) + 4Rz$$

before any singularity formation and hence the following.

**Proposition 1.** If  $d\sigma = 2$  or g = 0, then

(46) 
$$V_x(z) = V_x(0) + 2V_{xp}(0)z + 2(H(0) - \bar{H})z^2 + \frac{2R}{3}z^3, \quad z \in [0, z_*).$$

The analogous result  $(V_x \sim z^3)$  for the *linear* Schrödinger equation (d = 1, g = 0) with a random potential has been proved previously [3], [7].

We have from (43) that

(47) 
$$\frac{\partial^2}{\partial z^2} V_x = 4H + \frac{(4 - 2d\sigma)g}{\sigma + 1} \int \rho^{\sigma + 1} d\mathbf{x} - 2|\bar{\mathbf{p}}|^2$$

and hence

$$\frac{\partial^2}{\partial z^2} V_x \leq 4H - 4\bar{H} = 4(H(0) + Rz) - 4\bar{H}, \text{ for } g(2 - d\sigma) < 0$$
  
$$\frac{\partial^2}{\partial z^2} V_x \geq 4H - 4\bar{H} = 4(H(0) + Rz) - 4\bar{H}, \text{ for } g(2 - d\sigma) \ge 0$$

On the other hand, from (44) we obtain for any g

$$\frac{\partial^2}{\partial z^2} V_x \leq 2d\sigma (H - \bar{H}), \quad \text{for } 2 - d\sigma \leq 0$$
$$\frac{\partial^2}{\partial z^2} V_x \geq 2d\sigma (H - \bar{H}), \quad \text{for } 2 - d\sigma \geq 0.$$

Integrating the above inequalities twice, we obtain the following.

**Proposition 2.** The estimates hold

$$V_x(z) \leq V_x(0) + 2V_{xp}(0)z + 2(H(0) - \bar{H})z^2 + \frac{2}{3}Rz^3, \quad \text{for } g(2 - d\sigma) \leq 0$$
  
$$V_x(z) \geq V_x(0) + 2V_{xp}(0)z + 2(H(0) - \bar{H})z^2 + \frac{2}{3}Rz^3, \quad \text{for } g(2 - d\sigma) \geq 0$$

and

$$V_{x}(z) \leq V_{x}(0) + 2V_{xp}(0)z + d\sigma(H(0) - \bar{H})z^{2} + \frac{d\sigma}{3}Rz^{3}, \quad \text{for } 2 \leq d\sigma$$
$$V_{x}(z) \geq V_{x}(0) + 2V_{xp}(0)z + d\sigma(H(0) - \bar{H})z^{2} + \frac{d\sigma}{3}Rz^{3}, \quad \text{for } 2 \geq d\sigma$$

for all  $z \in [0, z_*)$ .

Therefore

**Corollary 1.** Assume g < 0 (hence  $H \ge 0$ ). Then

$$V_x(z) \le V_x(0) + 2V_{xp}(0)z + (d\sigma \lor 2)[H(0) - \bar{H}]z^2 + \frac{d\sigma \lor 2}{3}Rz^3$$

and

(48) 
$$V_x(z) \ge V_x(0) + 2V_{xp}(0)z + (d\sigma \wedge 2)[H(0) - \bar{H}]z^2 + \frac{d\sigma \wedge 2}{3}Rz^3$$

for  $z \in [0, z_*)$ 

4.3. Singularity formation. Finite-time singularity for the critical or supercritical NLS  $(d\sigma \ge 2)$  without a random potential is a well known effect [23]. In this case the singularity is the blow-up type  $V_p$ ,  $\|\rho\|_{\sigma+1} \to \infty$ . Here we take (26) as a model equation to gain some insight into singularity formation in the presence of a random potential.

First we consider the self-focusing case  $g \ge 0$ . For  $d\sigma \ge 2$  one can bound  $V_x$  as

(49) 
$$V_x(z) \le V_x(0) + 2V_{xp}(0)z + d\sigma(H(0) - \bar{H})z^2 + \frac{d\sigma R}{3}z^3 \equiv F(z)$$

and looks for the situation when F(z) vanishes.

The sufficient conditions for F(z) to vanish at a finite positive z are that F(z) takes a nonpositive value  $F(z_0) \leq 0$  at its local minimum point  $z_0 > 0$ . The local minimum point  $z_0$  is given by

(50) 
$$z_0 = \frac{\bar{H} - H(0) + \sqrt{(H(0) - \bar{H})^2 - 2RV_{xp}(0)/(d\sigma)}}{R}$$

Therefore we are led to the singularity conditions for  $g \ge 0$ .

**Proposition 3.** For  $d\sigma \geq 2, g > 0$ , the local solution of the nonlinear kinetic eq. (26) with the longitudinal randomness develops singularity at a finite time  $z_* \leq z_0$  given by (50) under the condition  $F(z_0) \leq 0$  and either one of the following conditions

$$(51) V_{xp}(0) < 0$$

(52) 
$$V_{xp}(0) > 0, \quad H(0) < \bar{H} - \sqrt{\frac{2RV_{xp}(0)}{d\sigma}}.$$

**Remark 1.** Clearly, the condition  $F(z_0) \leq 0$  requires H(0) to be sufficiently below  $\overline{H}$  by allowing the potential energy

$$-\frac{g}{\sigma+1}\int \rho^{\sigma+1}d\mathbf{x}d\mathbf{p}$$

at z = 0 to be sufficiently negative. We will give an explicit blow-up condition for the supercritical nonlinearity in the next section.

In the linear or self-defocusing case  $g \leq 0$  the right side of (46) or (48) is always positive in view of the inequality

$$2|V_{xp}| \le V_x + V_p,$$

cf. (42).

Therefore, the above result suggests that the scattering term does not prevent large scale singularity, even though it appears (from the upper bound  $z_0$ ) that it takes longer time than the time to singularity in the homogeneous NLS [23].

With the assumption that the variance identity and the energy laws (37), (38) hold for the maximally extended local solutions of eq. (26) then it is clear that the singularity is the blow-up type

$$\lim_{z \to z_*} \int \rho^{\sigma+1} d\mathbf{x} = \infty$$

or equivalently

$$\lim_{z \to z_*} S_p = \infty$$

by the finitude of the Hamiltonian.

Next we will follow the argument of [14] to show more explicitly the blow-up mechanism in the case with the supercritical, self-focusing nonlinearity and give a sharper bound on  $z_*$  under certain circumstances.

**Proposition 4.** Suppose  $d\sigma > 2, g > 0$ . Then under the conditions (51),

(53) 
$$H(z_0) = H(0) + Rz_0 \le 0$$

with  $z_0$  given by (50),  $V'_x$  as well as  $V_p$  blow up at a finite time  $z_*$  where

$$z_* \le z_0 \wedge \frac{2V_x(0)}{V_{xp}(0)(2-d\sigma)}$$

Let us give the argument below. Since blow-up is a local phenomenon,  $V_x$  is a poor indicator of its occurrence. To this end a more useful object to consider is  $V_p$ .

From (44) it follows that

(54) 
$$\frac{\partial^2}{\partial z^2} V_x \leq (2 - d\sigma) V_p < 0, \quad z < z_*.$$

Hence  $V'_x(z)$  is a negative, decreasing function for  $z < z_*$ . Also by the Cauchy-Schwartz inequality, we get

(55) 
$$0 \le V_{xp}^2 \le V_x V_p \le V_x(0) V_p, \quad 0 < z < z_*$$

and hence

(56) 
$$V_p \ge \frac{V_{xp}^2}{V_x(0)}$$

Let  $A(z) = -V'_x(z) > 0, z < z_*$ . We have from (44) and (55) the differential inequality

(57) 
$$\frac{\partial}{\partial z}A \ge CA^2, \quad C = \frac{d\sigma - 2}{4V_x(0)} > 0$$

which yields the estimate

$$A(z) \ge \frac{A(0)}{1 - zCA(0)}, \quad z < \frac{1}{CA(0)}$$

and thus the blow-up of  $-V'_x(z)$  at a finite time. This along with (56) then (53) implies the divergence of  $V_p$  at a finite time

$$\lim_{z \to z_*} V_p(z) = \infty$$

with

(58) 
$$z_* \le \frac{1}{CA(0)} = \frac{2V_x(0)}{V_{xp}(0)(2 - d\sigma)}$$

For high power  $d\sigma \gg 2$ , (58) is a better upper bound for  $z_*$  than  $z_0$  given by (50).

The preceding argument demonstrates clearly the blow-up mechanism, namely the quadratic growth property (57) as well as gives a sharper bound on the blow-up time for large  $d\sigma$ .

# 5. The transverse case

The scattering operator  $\mathcal{L}$  in this case corresponds to elastic scattering, instead of the inelastic scattering of the longitudinal case. This affects the evaluation of  $\mathcal{L}^* = \mathcal{L}$  on quadratic polynomials:

$$\begin{aligned} \mathcal{L}\mathbf{x} &= 0\\ \mathcal{L}\mathbf{p} &= 2\pi \int \delta(\frac{|\mathbf{p}|^2 - |\mathbf{q}|^2}{2}) \int \Phi(w, \mathbf{p} - \mathbf{q}) dw (\mathbf{p} - \mathbf{q}) d\mathbf{q} = 0\\ \mathcal{L}|\mathbf{x}|^2 &= 0\\ \mathcal{L}\mathbf{x} \cdot \mathbf{p} &= 0\\ \mathcal{L}|\mathbf{p}|^2 &= -2\pi \int \delta(\frac{|\mathbf{p}|^2 - |\mathbf{q}|^2}{2}) \int \Phi(w, \mathbf{p} - \mathbf{q}) dw \left[|\mathbf{p}|^2 - |\mathbf{q}|^2\right] d\mathbf{q} = 0. \end{aligned}$$

The same results hold for the diffusion operator  $\mathcal{L}$  with the diffusion matrix (23).

5.1. Rate of dispersion. The variance identity still holds Hence the second derivative of  $V_x$  becomes

(59) 
$$\frac{\partial^2}{\partial z^2} V_x = 4(H - \bar{H}) + \frac{2(2 - d\sigma)g}{\sigma + 1} \int \rho^{\sigma + 1} d\mathbf{x}$$

(60) 
$$= 2d\sigma(H - \bar{H}) + (2 - d\sigma)V_p.$$

The main difference from the longitudinal case is that the Hamiltonian is invariant

$$\frac{\partial}{\partial z}H = 0.$$

By the same argument as in the longitudinal case we have the following analogous estimates.

**Proposition 5.** If  $d\sigma = 2$  or g = 0, then

(61) 
$$V_x(z) = V_x(0) + 2V_{xp}(0)z + 2(H(0) - \bar{H})z^2, \quad z \in [0, z_*).$$

**Proposition 6.** The following estimates hold for  $z \in [0, z_*)$ :

$$V_x(z) \ge V_x(0) + 2V_{xp}(0)z + 2(H - \bar{H})z^2, \quad \text{for } g(2 - d\sigma) \ge 0$$
  
$$V_x(z) \le V_x(0) + 2V_{xp}(0)z + 2(H - \bar{H})z^2, \quad \text{for } g(2 - d\sigma) \le 0$$

and

$$V_x(z) \ge V_x(0) + 2V_{xp}(0)z + d\sigma(H - H)z^2, \text{ for } 2 \ge d\sigma$$
  
$$V_x(z) \le V_x(0) + 2V_{xp}(0)z + d\sigma(H - \bar{H})z^2, \text{ for } 2 \le d\sigma.$$

Therefore,

Corollary 2. Assume g < 0 (hence  $H \ge 0$ ). Then for  $z \in [0, z_*)$  $(62V_x(0) + 2V_{xp}(0)z + (d\sigma \land 2)[H - \bar{H}]z^2 \le V_x(z) \le V_x(0) + 2V_{xp}(0)z + (d\sigma \lor 2)[H - \bar{H}]z^2$ .

From these estimates we see that a ballistic kind of motion takes place in the transverse case.

5.2. Singularity. By finding the zeros of the upper bounds in Proposition 6 we can derive the conditions for singularity formation. As in the longitudinal case, for  $g \leq 0$ , (61) and the left side of (62) are always positive.

**Proposition 7.** For  $g > 0, d\sigma \ge 2$ , the local solution of the nonlinear kinetic eq. (26) with the transverse randomness develops singularity in a finite time  $z_* < \infty$  under either of the following conditions

or

(64) 
$$H \ge \bar{H}, \quad V_{xp}(0) < 0, \quad H \le \bar{H} + |V_{xp}(0)|^2 / (d\sigma V_x(0)).$$

**Remark 2.** It is easy to see that the singularity time  $z_*$  is bounded from above by

(65) 
$$z_0 = \frac{-V_{xp}(0) + \sqrt{|V_{xp}(0)|^2 - d\sigma V_x(0)(H - \bar{H})}}{d\sigma (H - \bar{H})}$$

under condition (63) and by

$$z_0 = \frac{-V_{xp}(0) - \sqrt{|V_{xp}(0)|^2 - d\sigma V_x(0)(H - \bar{H})}}{d\sigma (H - \bar{H})}$$

under condition (64).

The preceding singularity conditions are identical to those for the homogeneous (super)-critical NLS equation [23]. This is already suggested by the previous result [11] where (63) is shown to be the instability condition for the diffusion approximation of the kinetic equation (26) with the kernel (19).

With more stringent conditions one can demonstrate a stronger sense of blow-up. Let us state the result.

**Proposition 8.** Suppose  $d\sigma > 2, g > 0$ . Then under the conditions

$$H < H$$
,

 $V'_x$  as well as  $V_p$  blow up at a finite time  $z_* < z_0$ .

Let us sketch the argument below. From (60) it follows that

(66) 
$$\frac{\partial^2}{\partial z^2} V_x \leq (2 - d\sigma) V_p < 0, \quad z < z_*$$

Since  $V'_x(z)$  is a decreasing function and becomes negative after a finite time  $\bar{z}$  when  $V_x(z)$  achieves its maximum  $\bar{V}_x$ .

Again for  $A(z) = -V'_x(z) > 0, z > \overline{z}$  we have from (66) and (55) the differential inequality

$$\frac{\partial}{\partial z}A \ge CA^2, \quad C = \frac{d\sigma - 2}{4V_x(\bar{z})} > 0, \quad z > \bar{z}.$$

This is a plausible mechanism for the finite-time blow-up of the kinetic energy and the potential energy.

## 6. Fokker-Planck equation with dissipation

In this section we will apply the same analysis to the Fokker-Planck equation (26) with, instead, the Caldeira-Leggett operator [4]

(67) 
$$\mathcal{L}W = D\Delta_{\mathbf{p}}W + \lambda \nabla_{\mathbf{p}}[\cdot \mathbf{p}W], \quad D, \lambda > 0$$

which is, perhaps, the simplest example of the phase-space models for open, dissipative systems [6], [1]. As remarked before in this context z is the physical time,  $\mathbf{x}$  the physical coordinates and  $\mathbf{p}$  the momentum in  $\mathbb{R}^3$ . If the open quantum system is the electron motion in a semi-conductor then the nonlinearity arises from the electrostatic Hartree potential self-induced by the density  $\rho$  with self-defocusing nonlinearity. Since operator (67) is not self-adjoint (w.r.t. the Lebesgue measure) it is outside the scope of the NLS eq. with a real-valued random potential (1). The method used here can be applied to such a problem. However, in order to compare how different decoherence mechanisms affect wave spreading and collapse we will still consider the same power-law nonlinearity as in the previous cases.

The dissipative effect is of frictional nature and the Fokker-Planck equation with (67) still preserves the total mass

$$\frac{\partial}{\partial z}N = 0$$

As before we set N = 1.

We re-calculate the action of  $\mathcal{L}^*$ , the adjoint operator, on the quadratic polynomials:

$$\mathcal{L}^* \mathbf{x} = 0$$
  

$$\mathcal{L}^* \mathbf{p} = -\lambda \mathbf{p}$$
  

$$\mathcal{L}^* |\mathbf{x}|^2 = 0$$
  

$$\mathcal{L}^* \mathbf{x} \cdot \mathbf{p} = -\lambda \mathbf{x} \cdot \mathbf{p}$$
  

$$\mathcal{L}^* |\mathbf{p}|^2 = 2dD - 2\lambda |\mathbf{p}|^2.$$

6.1. Rate of Spread. Using these results and integrating by parts we get

$$\frac{\partial}{\partial z} \bar{\mathbf{x}} = \bar{\mathbf{p}}$$
$$\frac{\partial}{\partial z} \bar{\mathbf{p}} = -\lambda \bar{\mathbf{p}}$$

which again constitute an exactly solvable system:

(68) 
$$\bar{\mathbf{p}}(z) = e^{-\lambda z} \bar{\mathbf{p}}(0)$$

(69) 
$$\bar{\mathbf{x}}(z) = \bar{\mathbf{x}}(0) + \frac{\bar{\mathbf{p}}(0)}{\lambda}(1 - e^{-\lambda z}).$$

The Hamiltonian  $\overline{H}$ , defined by (36), of the mean motion is then given by

$$\bar{H}(z) = \bar{H}(0)e^{-2\lambda z}.$$

Also,

$$\frac{\partial}{\partial z} V_x = 2V_{xp}$$

$$\frac{\partial}{\partial z} V_{xp} = d\sigma (H - \bar{H}) + (1 - \frac{d\sigma}{2})V_p - \lambda V_{xp}$$

$$= 2(H - \bar{H}) + \frac{(2 - d\sigma)g}{\sigma + 1} \int \rho^{\sigma + 1} d\mathbf{x} - \lambda V_{xp}$$

From these we obtain the upper bound

(70) 
$$V_{xp}(z) \leq V_{xp}(0) + d\sigma e^{-\lambda z} \int_{0}^{z} e^{s\lambda} (H(s) - \bar{H}(s)) ds, \text{ for } d\sigma \geq 2$$

(71) 
$$V_{xp}(z) \geq V_{xp}(0) + d\sigma e^{-\lambda z} \int_0^z e^{s\lambda} (H(s) - \bar{H}(s)) ds, \text{ for } d\sigma \le 2$$

and the bounds

(72) 
$$V_{xp}(z) \leq V_{xp}(0) + 2e^{-\lambda z} \int_0^z e^{s\lambda} (H(s) - \bar{H}(s)) ds$$
, for  $g(2 - d\sigma) \leq 0$ 

(73) 
$$V_{xp}(z) \ge V_{xp}(0) + 2e^{-\lambda z} \int_0^z e^{s\lambda} (H(s) - \bar{H}(s)) ds$$
, for  $g(2 - d\sigma) \ge 0$ .

On the other hand, the rate of change of the Hamiltonian is given by

$$\frac{\partial}{\partial z} [H(z) - \bar{H}(z)] = dD - \lambda V_p$$
  
=  $dD - 2\lambda (H - \bar{H}) - \frac{2\lambda g}{\sigma + 1} \int \rho^{\sigma + 1} d\mathbf{x}.$ 

For self-focusing nonlinearity  $g \geq 0$  the upper bound on H follows

(74) 
$$H(z) - \bar{H}(z) \leq H(0) - \bar{H}(0) + (1 - e^{-2\lambda z}) \frac{dD}{2\lambda}$$

whereas for the self-defocusing nonlinearity  $g \leq 0$  we have both the lower as well the upper bounds

$$H(0) - \bar{H}(0) + (1 - e^{-2\lambda z})\frac{dD}{2\lambda} \le H(z) - \bar{H}(z) \le H(0) - \bar{H}(0) + dDz.$$

Therefore for  $g \geq 0$  we have the upper bounds

$$V_{xp}(z) \le V_{xp}(0) + d\sigma \left[ \frac{H(0) - \bar{H}(0)}{\lambda} (1 - e^{-\lambda z}) + (1 - e^{-\lambda z})^2 \frac{dD}{2\lambda^2} \right]_{16}$$

whereas for  $g \leq 0$ , we have the estimates

$$V_{xp}(z) \geq V_{xp}(0) + (d\sigma \wedge 2) \left[ \frac{H(0) - \bar{H}(0)}{\lambda} (1 - e^{-\lambda z}) + (1 - e^{-\lambda z})^2 \frac{dD}{2\lambda^2} \right]$$
  
$$V_{xp}(z) \leq V_{xp}(0) + (d\sigma \vee 2) \left[ \frac{H(0) - \bar{H}(0)}{\lambda} - \frac{dD}{\lambda^2} \right] (1 - e^{-\lambda z}) + \frac{d^2 \sigma D}{\lambda} z.$$

Therefore,

**Proposition 9.** For  $g \ge 0, z \in [0, z_*)$ 

$$V_x(z) \leq V_x(0) + 2V_{xp}(0)z + \frac{d\sigma}{\lambda}(2H(0) - 2\bar{H}(0) + \frac{dD}{\lambda})z \\ -\frac{2d\sigma}{\lambda^2} \left[H(0) - \bar{H}(0) + \frac{dD}{\lambda}\right](1 - e^{-\lambda z}) + \frac{d^2\sigma D}{2\lambda^3}(1 - e^{-2\lambda z});$$

whereas for  $g \leq 0, z \in [0, z_*)$ 

$$V_x(z) \geq V_x(0) + 2V_{xp}(0)z + \frac{d\sigma \wedge 2}{\lambda}(2H(0) - 2\bar{H}(0) + \frac{dD}{\lambda})z \\ -\frac{2(d\sigma \wedge 2)}{\lambda^2} \left[H(0) - \bar{H}(0) + \frac{dD}{\lambda}\right](1 - e^{-\lambda z}) + \frac{(d\sigma \wedge 2)dD}{2\lambda^3}(1 - e^{-2\lambda z}) \\ V_x(z) \leq V_x(0) + 2V_{xp}(0)z + 2(d\sigma \vee 2) \left[\frac{H(0)}{\lambda} - \frac{dD}{\lambda^2}\right] \left[z + \frac{1}{\lambda}(e^{-\lambda z} - 1)\right] + \frac{d^2\sigma D}{\lambda}z^2.$$

6.2. Singularity. The singularity conditions for the self-focusing case are determined from that the right side of (75) has a positive root.

**Proposition 10.** For  $d\sigma \ge 2, g > 0$ , the local solution of the Fokker-Planck eq. (26) with (67) develops finite-time singularity if

(76) 
$$V_{xp}(0) + \frac{d\sigma}{\lambda} (H(0) - \bar{H}(0) + \frac{dD}{2\lambda}) < 0.$$

The condition (76) is better suited for large  $\lambda$  and can be much improved for small  $\lambda$ . As  $\lambda \to 0$  this case should approach to that of the longitudinal case with **D** given by (22) being a scalar.

Let us sketch an explicit blow-up mechanism in the super-critical case under different conditions.

**Proposition 11.** Suppose  $d\sigma > 2, g > 0$ . Then under the conditions

(77) 
$$V_{xp}(0) \le 0, \quad H(0) - \bar{H}(0) + \frac{dD}{2\lambda} < 0$$

and

(78) 
$$V_x(0) < \frac{d\sigma - 2}{2\lambda}$$

 $V_x'$  as well as  $V_p$  blow up at a finite time  $z_*$  where

$$z_* < \frac{1}{\lambda} \ln \frac{d\sigma - 2}{d\sigma - 2 - 2\lambda V_x(0)}.$$

We present the argument here. Rewriting some of the preceding calculations, we have

$$\frac{\partial}{\partial z}V_x = 2V_{xp}$$

and

(79) 
$$\frac{\partial}{\partial z} V_{xp} = d\sigma (H - \bar{H}) + (1 - \frac{d\sigma}{2}) V_p - \lambda V_{xp}$$

Condition (77) and (74) imply that

$$H(z) - \bar{H}(z) \le -e^{-2\lambda z} \frac{dD}{2\lambda}.$$

Using this in (79) we obtain

$$\frac{\partial}{\partial z}(e^{\lambda z}V_{xp}(z)) \le -e^{-\lambda z}\frac{d^2\sigma D}{2\lambda} + e^{\lambda z}(1-\frac{d\sigma}{2})V_p$$

which together with (77) implies  $V_{xp}(z) < 0, z \le z_*$  and  $V_x(z)$  is decreasing. Using the inequality

$$V_p \ge \frac{V_{xp}^2}{V_x(0)}$$

in (79) we obtain again the quadratic growth estimate

$$\frac{\partial}{\partial z}A \ge \frac{d\sigma - 2}{2V_x(0)}A^2 - \lambda A, \quad z < z_*$$

for  $A = -V_{xp} > 0$ , or equivalently

$$\frac{\partial}{\partial z}B \geq \frac{d\sigma - 2}{2V_x(0)}e^{-\lambda z}B^2, \quad z < z_*$$

for

$$B = e^{\lambda z} A.$$

From this it follows that

$$B(z) \ge \frac{B(0)}{1 - \frac{d\sigma - 2}{2\lambda V_x(0)}(1 - e^{-\lambda z})}$$

the right-hand side of which blows up at a finite time if and only if (78) holds.

The finite-time blow-up of B would also lead to the finite-time blow-up of A and hence of  $V_p$ .

## 7. Conclusion

We have derived several kinetic equations meant to model the large scale, low Fresnel number behavior of the NLS equation with a rapidly fluctuating random potential based on the rigorous theory [9], [10] for the linear Schrödinger waves in the same situation. The low Fresnel number waves interacting with a rapidly oscillating potential give rise to, in the scaling (4), a self-averaging limit of a deterministic kinetic equation with a scattering operator.

We analyze these kinetic equations in order to shed light on two problems: the rate of dispersion and the singularity formation. Our main assumptions are the existence of local solutions of the kinetic models in the space S, that the local solutions satisfy the Hamiltonian laws (37), (38) and the variance identity.

The scattering operator in the kinetic equations may be a Boltzmann-type operator or a Fokker-Planck operator, depending on whether the random potential is fully resolved by the wave or not. This, however, does not affect our results for either problem.

What is more important for our investigation is the *structure* of the scattering kernel in the Boltzmann operator or the diffusion matrix of the Fokker-Planck operator. There are two types of structures: the longitudinal regime when the longitudinal scale of the random potential is dominant and the transverse regime when the transverse scale of the random potential is dominant. The third kind of kinetic equation that we have analyzed is based on the Fokker-Planck equation with the longitudinal type of diffusion matrix plus a frictional term. This model is motivated by the Caldeira-Leggett operator for open quantum dynamics.

For the problem of dispersion, we have shown that the kinetic equations of the longitudinal type produce the cubic  $(z^3)$  law, that the kinetic equations of the transverse type produce the quadratic

 $(z^2)$  law and that the Fokker-Planck equation with friction produces the linear (z) law for the variance  $V_x(z)$  for  $z \in [0, z_*)$ .

For the problem of singularity, we have shown by analyzing the variance identity that the singularity and blow-up conditions in the transverse case remain the same as those for the homogeneous NLS equation with critical or supercritical self-focusing nonlinearity, that the singularity and blowup conditions have changed in the longitudinal case and in the frictional case.

Finally let us make a slight extension of our results by considering the nonlinear kinetic equation of the more general form

(80) 
$$\frac{\partial}{\partial z}W(z,\mathbf{x},\mathbf{p}) + \mathbf{p} \cdot \nabla_{\mathbf{x}}W(z,\mathbf{x},\mathbf{p}) + \mathcal{U}_{\gamma}W(z,\mathbf{x},\mathbf{p}) = \mathcal{L}W(z,\mathbf{x},\mathbf{p})$$

with  $\gamma > 0$  small, but positive and the mixed-state Wigner distributions (13) as initial data. Eq. (26) is formally the geometrical optics limit of eq. (80). Eq. (80) retains the diffraction effect of the nonlinear potential term that is absent in eq. (26) with  $\gamma = 0$ . Even though eq. (80) does not preserve positivity, it can be shown to preserve the mixed state structure (13) and therefore the marginal positivity (14), (15) as well as the inequalities (41), (42) in both the longitudinal and transverse cases. Since the calculations (30)-(34) are independent of  $\gamma \geq 0$ , all the results in Section 4 and 5 hold for eq. (80) with  $\gamma > 0$  as well. However, because the Caldeira-Leggett operator (67) is in the Lindblad form only in the classical limit [6], therefore in order to maintain the solution in the the mixed-state form (13) we need to set  $\gamma = 0$  in this particular case.

## References

- A. Arnold, J.L. López, P.A. Markowich and J. Soler, An analysis of quantum Fokker-Planck Models: A Wigner function approach, preprint 2003.
- [2] G. Bal, G. Papanicolaou and L. Ryzhik, Radiative transport limit for the random Schrödinger equation, Nonlinearity, Vol. 15, 513-529, (2002).
- [3] L. Bunimovich, H.R. Jauslin, J.L. Lebowitz, A. Pellegrinotti and P. Nielaba, Diffusive energy growth in classical and quantum driven oscillators. J. Stat. Phys.62(1991), 793-817.
- [4] A.O. Caldeira and A.J. Leggett, Path integral approach to quantum Brownian motion. Physica A.121(1983), 587-616.
- [5] A. Debussche and L. Di Menza, Numerical simulation of focusing stochastic nonlinear Schrödinger equations. Phys. D 162 (2002), 131-154.
- [6] L. Diósi, Caldeira-Leggett master equation and medium temperatures. Physica A.199(1993), 517-526.
- [7] E.B. Erdogan, R. Killip and W. Schlag, Energy growth in Schrödinger's equation with Markovian driving. Commun. Math. Phys.240(2003), 1-29.
- [8] L. Erdös and H.T. Yau, Linear Boltzmann equation as the weak coupling limit of a random Schrödinger equation, Comm. Pure Appl. Math. 53, 667-735 (2000).
- [9] A. Fannjiang, Self-averaging in scaling limits for random parabolic waves. Archives for Rational Mechanics and Analysis, in press.
- [10] A. Fannjiang, Self-averaging limits of a quantum system in random environments, preprint.
- [11] A. Fannjiang, S. Jin and G. Papanicolaou, High frequency behavior of the focusing nonlinear Schroedinger equation with random inhomogeneities. SIAM J. Appl. Math. 63(2003), 1328-1358.
- [12] P. Gérard and E. Leichtnam, Ergodic properties of eigenfunctions for the Dirichlet problem, Duke Math. J.71 (1993), 559-607.
- [13] P. Gérard, P.A. Markowich, N.J. Mauser and F. Poupaud, Homogenization limits and Wigner transforms, Commun. Pure Appl. Math. I (1997), 323-379.
- [14] R.T. Glassey, On the blow-up of solutions to the Cauchy problem for nonlinear Schrödinger equations, J. Math. Phys. 18(1977), 1794-1797.
- [15] A. Hasegawa and Y. Kodama, Solitons in Optical Communications, Oxford University Press, Oxford, 1995.
- [16] T. Kato, On nonlinear Schrödinger equations, Ann. INst. H. Poincaré, Phys. Théor. 46 (1987), 113-129.
- [17] P.L. Lions and T. Paul, Sur les mesures de Wigner. Rev. Mat. Iberoamericana—sf 9, 553-618 (1993).
- [18] B.A. Malomed, Variational methods in nonlinear fiber optics and related fields. Prog. Optics43(2002), 71-193.
- [19] D. Michaelis, U. Peschel and F. Lederer, Phys. Rev. A 56 (1997), 3366.

- [20] A. C. Newell and J. V. Moloney, Nonlinear Optics, Addison-Wesley Publishing Company, Redwood City, CA., 1992.
- [21] F. Poupaud and A. Vasseur, Classical and quantum transport in random media. Math. Pure et Appli. 82, 711-748 (2003).
- [22] H. Spohn, Derivation of the transport equation for electrons moving through random impurities, J. Stat. Phys., 17, 385-412 (1977).
- [23] C. Sulem and P.-L. Sulem, The nonlinear Schrdinger equation: self-focusing and wave collapse. Springer, New York, 1999.
- [24] S. Tcheremchantsev, Transport properties of Markovian Anderson model, Commun. Math. Phys. 196:1 (1998), 105-131.
- [25] N.G. van Kampen, A new approach to noise in quantum mechanics. J. Stat. Phys.115(2004), 1057-1071.