

# A note on the forced Burgers equation

S. Eule<sup>1</sup>, R. Friedrich  
Institute of Theoretical Physics  
Westfälische Wilhelms Universität  
Münster  
Wilhelm-Klemm-Str. 9  
G-48149 Münster

## Abstract

We obtain the exact solution for the Burgers equation with a time dependent forcing, which depends linearly on the spatial coordinate. For the case of a stochastic time dependence an exact expression for the joint probability distribution for the velocity fields at multiple spatial points is obtained. A connection with stretched vortices in hydrodynamic flows is discussed.

PACS number(s): 02.50.-r, 05.40.-a, 05.90.-a, 47.32.Cc

Burgers equation [1], [2], [3] has extensively been investigated as a model equation for the study of the problem of turbulence. The homogeneous equation can be exactly solved by a Hopf-Cole transformation [4], [5] such that many properties of the spatio-temporal behaviour can be obtained by analytical means. In recent years, the stochastically forced Burgers equation has attracted much interest due to the work of Polyakov [6]. He considered a white noise forcing  $f(x, t)$  with spatial correlations  $\langle f(x, t)f(x', t) \rangle = \kappa(x - x')\delta(t - t')$ . This equation has been found to exhibit scaling behaviour. Polyakov suggested to use an operator product expansion in order to assess the scaling behaviour of the velocity increments.

The purpose of the present note is to present the general solution of the forced Burgers equation

$$\left[\frac{\partial}{\partial t} + v(r, t)\frac{\partial}{\partial r}\right]v(r, t) = \nu\frac{\partial^2}{\partial r^2}v(r, t) + G(t)r \quad , \quad (1)$$

where  $G(t)$  is an arbitrary function of time  $t$ . As we shall discuss below,

---

<sup>1</sup>yeahhart@uni-muenster.de, Tel.:+49-251-8334924, Fax: +49-251-8336328

this problem is related to the Polyakov problem. The investigation of the special type of forcing,  $f(r, t) = rG(t)$ , has the advantage that one can find an analytic solution for an arbitrary function  $G(t)$ .

This solution can be found by direct calculation using the ansatz

$$v(r, t) = a(t)r + \Lambda(t)w(\Lambda(t)r, \tau(t)) \quad . \quad (2)$$

Thereby, the functions  $a(t)$ ,  $\Lambda(t)$ ,  $\tau(t)$  are determined by the following solvability conditions:

$$\begin{aligned} \dot{a}(t) + a(t)^2 &= G(t) \\ \dot{\Lambda} &= -a(t)\Lambda \quad , \\ \dot{\tau} &= \Lambda^2 \quad . \end{aligned} \quad (3)$$

The quantities  $\Lambda(t)$  and  $\tau(t)$  can be determined as a function of  $a(t)$ ,

$$\Lambda(t) = e^{-\int_0^t dt' a(t')} \quad , \quad \tau = \int_0^t dt' e^{-2\int_0^{t'} dt'' a(t'')} \quad (4)$$

In the following we shall see that the quantity

$$\frac{\Lambda^2(t)}{\tau(t)} = 1/\sigma^2(t) = \frac{d}{dt} \ln \tau(t) \quad (5)$$

will play a major role.

In order that the ansatz eq. (2) obeys the forced Burgers equation the function  $w(\xi, \tau)$  has to obey the homogeneous equation. For instance,  $w(\xi, \tau)$  could be the single shock solution

$$w(\xi, \tau) = -A \tanh \frac{A}{2\nu} \xi \quad (6)$$

Of interest are also scaling solutions of the form [3]

$$w(\xi, \tau) = \frac{1}{\sqrt{\tau}} W\left(\frac{\xi}{\sqrt{\tau}}\right) \quad , \quad (7)$$

which e.g. evolve from the initial condition  $w(\xi, 0) = \delta(\xi)$ .

Since the solution of the homogeneous Burgers equation is known in terms of the initial conditions the function  $\Lambda(t)w(\xi, \tau)$  can be represented according to

$$\Lambda(t)w(\Lambda r, \tau) = -2\nu\Lambda(t) \frac{\partial}{\partial r} \ln \int_{-\infty}^{\infty} dr' e^{-(r-r')^2/(4\nu\sigma^2(t))} e^{-\int_{-\infty}^{r'} dr'' \Lambda(t)w(\Lambda r'', 0)/2\nu} \quad (8)$$

We mention that the structure of the solution, eq. (2), is similar to the solution describing stretched vortices in the Navier-Stokes equation (see e.g. [7]). The analogy becomes even more clear since, as we shall briefly mention below, the treatment can be extended to the case of the multidimensional Burgers equation. Stretched vortex solutions of the Navier-Stokes equation are supposed to play a fundamental role in the organization of turbulence. Stretching has the effect to render a decaying vortex like the Lamb-Oseen vortex into a stationary vortex (e.g. the Burgers vortex). Similar phenomena are expected in the present case of the forced Burgers equation.

As can be seen from the solvability conditions, the crucial point is to obtain a solution of the nonlinear differential equation for  $a(t)$ . The behaviour of  $a(t)$  can be visualized by introducing the potential  $V(a) = \frac{1}{3}a^3$ : The variable  $a(t)$  behaves as an overdamped particle moving in the potential  $V(a)$  under the influence of the time dependent force  $G(t)$ . By introducing the variable  $b(t)$  according to

$$a(t) = \frac{\dot{b}(t)}{b(t)} \quad (9)$$

the equation of motion for  $a(t)$  can be transformed into a linear equation of second order (Hill's equation)

$$\ddot{b}(t) = G(t)b(t) \quad . \quad (10)$$

In terms of  $b(t)$ , the variables  $\Lambda(t)$  and  $\tau(t)$  are determined by

$$\begin{aligned} \Lambda(t) &= \frac{b(0)}{b(t)} \\ \tau(t) &= \int_0^t dt' \left( \frac{b(0)}{b(t')} \right)^2 \end{aligned} \quad (11)$$

In the following, we shall briefly consider some cases of functions  $G(t)$ .

The first case is a stationary forcing,  $G(t) = g$ . If we take  $g$  to be a positive number, one obtains two solutions for a

$$a = \pm\sqrt{g} \quad \Lambda = e^{\mp\sqrt{g}t} \quad \tau = \frac{e^{\mp 2\sqrt{g}t} - 1}{\mp 2\sqrt{g}} \quad . \quad (12)$$

Let us first consider the case of a positive value of  $a = \sqrt{g}$ . This solution is linearly stable. The contribution of a shock

$$- A\Lambda(t)\tanh\frac{A\Lambda(t)}{2\nu}r \quad (13)$$

decays to zero. A scale invariant solution also decays, since

$$\frac{\Lambda(t)}{\sqrt{\tau(t)}} W\left(\frac{\Lambda(t)}{\sqrt{\tau(t)}} r\right) \quad (14)$$

and

$$\lim_{t \rightarrow \infty} \frac{\Lambda(t)}{\sqrt{\tau(t)}} = 2\sqrt{g} \frac{e^{-2\sqrt{g}t}}{1 - e^{-2\sqrt{g}t}} = 0 \quad (15)$$

The case of a negative value of  $a = -\sqrt{g}$  is linearly unstable. Furthermore, shocklike contributions increase in time connected with a steepening of the shock. Scale invariant contributions in the long time limit become stationary since

$$\lim_{t \rightarrow \infty} \frac{\Lambda(t)}{\sqrt{\tau(t)}} = 2\sqrt{g} \frac{e^{2\sqrt{g}t}}{e^{2\sqrt{g}t} - 1} = \sqrt{2\sqrt{g}} \quad (16)$$

such that the solution takes the form

$$v(r, t) = -\sqrt{gr} + \sqrt{2\sqrt{g}} W(\sqrt{2\sqrt{g}} r) \quad . \quad (17)$$

If we take  $g = -\omega^2$  to be negative, we obtain an oscillating solution for  $b(t)$  and a finite time singularity for  $a(t)$ :

$$b(t) = A \cos \omega t \quad a(t) = -\omega \tan \omega t \quad (18)$$

Furthermore, the quantity  $\sigma(t)$  reads:

$$\frac{1}{\sigma(t)} = \frac{\Lambda(t)}{\sqrt{\tau(t)}} = \sqrt{\frac{2\omega}{\sin 2\omega t}} \quad (19)$$

We shall now consider the case where  $G(t)$  is a stochastic white noise force. The Fokker-Planck equation for the probability distribution  $f(a, t) = \langle \delta(a - a(t)) \rangle$  reads

$$\frac{\partial}{\partial t} f(a, t) = \left[ \frac{\partial}{\partial a} \frac{\partial V(a)}{\partial a} + Q \frac{\partial^2}{\partial a^2} \right] f(a, t) \quad (20)$$

A stationary solution to this equation can be obtained:

$$f_0(a) = N e^{-V(a)/Q} \quad (21)$$

However, since the potential  $V(a)$  is not bounded from below, this stationary solution does not define a probability distribution, i.e. a stationary state does

not exist. Due to the quadratic nonlinearity in the Langevin equation a kind of stochastic finite time singularity exists, where  $a(t)$  tends to  $a(t) \rightarrow -\infty$ . This can be seen from the fact that the variable  $a$  moves like an overdamped particle in the unbounded potential under the influence of a stochastic force. As a consequence the variable  $\Lambda(t)$  increases, too.

Although there is a finite time singularity it is quite interesting to investigate the structure of the probability distribution of the velocity field  $v(r, t)$ . To this end we introduce the distribution  $P(a, \Lambda, \tau; t)$ , which obeys the Fokker-Planck equation

$$\frac{\partial}{\partial t}P = \left\{ \frac{\partial}{\partial \Lambda}a\Lambda - \frac{\partial}{\partial \tau}\Lambda^2 + \frac{\partial}{\partial a}a^2 + \frac{\partial^2}{\partial a^2} \right\}P \quad (22)$$

Using  $P(a, \Lambda, \tau; t)$  the probability distribution of the velocity field  $v(r, t)$ ,  $f(v, r, t) = \langle \delta(v - v(r, t)) \rangle$  takes the following form

$$\begin{aligned} f(v, r, t) &= \int d\Lambda d\tau da P(a, \Lambda, \tau, t) \delta(v - ar - \Lambda w(\Lambda r, \tau)) \\ &= \frac{1}{r} \int d\tau d\Lambda P\left(\frac{v - \Lambda w(\Lambda r, \tau)}{r}, \Lambda, \tau, t\right) \end{aligned} \quad (23)$$

The joint pdf for the velocity field at multiple scales  $r_i$  takes the form

$$f(\{v_i, r_i\}, t) = \int d\Lambda d\tau da P(a, \Lambda, \tau, t) \prod_{i=1}^N \delta(v_i - ar_i - \Lambda w(\Lambda r_i, \tau)) \quad . \quad (24)$$

This formula demonstrates how the spatial coherence of the stochastic force governs the statistics of the multiple scales pdf of the velocity field.

Using standard methods one may easily derive an evolution equation for the pdf  $f(v, r, t)$ :

$$\begin{aligned} \frac{\partial}{\partial t}f(v, r, t) &+ v \frac{\partial}{\partial r}f(v, r, t) + \frac{\partial}{\partial r} \int_{-\infty}^v dv' f(v', r, t) \\ &= -\nu \int dv' \int dr' \delta(r - r') v' \frac{\partial^2}{\partial r'^2} f(v', r'; v, r; t) \\ &+ Q(r, r) \frac{\partial^2}{\partial v^2} f(v, r, t) \quad . \end{aligned} \quad (25)$$

This equation is the first equation in a whole chain of equations for the joint pdf of the velocity field at multiple scales,

$$\begin{aligned} \frac{\partial}{\partial t}f(v_i, r_i, t) &+ \sum_i v_i \frac{\partial}{\partial r_i} f(v_i, r_i, t) + \sum_i \frac{\partial}{\partial r} \int_{-\infty}^{v_i} dv'_i f(v'_i, r_i, t) \\ &= -\sum_i \nu \int dv' \int dr' \delta(r - r') v' \frac{\partial^2}{\partial r'^2} f(v', r'; v_i, r_i; t) \\ &+ \sum_{i,j} Q(r_i, r_j) \frac{\partial^2}{\partial v_i \partial v_j} f(v_i, r_i, t) \end{aligned} \quad (26)$$

Here, the matrix  $Q(r_i, r_j)$  is given by

$$Q(r_i, r_j) = Qr_i r_j \quad (27)$$

We present this hierarchy, since Polyakov [6] considered the same hierarchy for the case of a translational invariant white noise forcing. Here, we could obtain a solution to this chain of equations by a direct calculation in terms of equation (24).

Let us now discuss the relationship of the above problem of forced Burgers equation to the problem considered by Polyakov [6] of translational invariant white noise forcing:

$$\begin{aligned} \left[ \frac{\partial}{\partial t} + u(x, t) \frac{\partial}{\partial x} \right] u(x, t) &= \nu \frac{\partial^2}{\partial x^2} u(x, t) + F(x, t) \\ \langle F(x, t) F(x', t) \rangle &= q(x - x') \delta(t - t') \quad . \end{aligned} \quad (28)$$

Of interest are the statistical properties of the velocity increments

$$v_x(r, t) = u(x + r, t) - u(x, t) \quad . \quad (29)$$

The increment  $v_x(r, t)$  obeys the evolution equation

$$\begin{aligned} \left[ \frac{\partial}{\partial t} + u(x, t) \frac{\partial}{\partial x} \right] v_x(r, t) + v_x(r, t) \frac{\partial}{\partial r} v_x(r, t) \\ = \nu \left[ \frac{\partial^2}{\partial r^2} v_x(r, t) - \frac{\partial^2}{\partial r^2} v_x(r, t) \Big|_{r=0} \right] + f_x(r, t) \\ f_x(r, t) = F(x + r, t) - F(x, t) \\ \langle f_x(r, t) f_x(r', t) \rangle = q(r - r') - q(r) - q(r') + q(0) \end{aligned} \quad (30)$$

The additional dissipative term arises since one has the condition

$$v_x(r = 0, t) = 0 \quad . \quad (31)$$

In a comoving coordinate system, i.e. considering the increments  $v_{x(t)}(r, t)$ , where the path  $x(t)$  is defined by

$$\dot{x}(t) = u(x(t), t) \quad (32)$$

we obtain the forced Burgers equation for the increment

$$\begin{aligned} \left[ \frac{\partial}{\partial t} + v(r, t) \frac{\partial}{\partial r} \right] v(x, t) \\ = \nu \left[ \frac{\partial^2}{\partial r^2} v(r, t) - \frac{\partial^2}{\partial r^2} v(r, t) \Big|_{r=0} \right] + G(r, t) \quad , \end{aligned} \quad (33)$$

Here,  $G(r, t)$  is given by

$$G(r, t) = F(x(t) + r, t) - F(x(t), t) \quad (34)$$

Polyakov investigated the statistics of increments in the limit of small values of  $r$  and approximated the function  $q(\xi)$  by

$$q(\xi) = q(0) - \frac{Q}{2}\xi^2 + .. \quad (35)$$

This relationship can only hold for  $|\xi| \ll \sqrt{q(0)/Q}$ , since for large values of  $\xi$   $q(\xi) \rightarrow 0$ . This approximation, however, is equivalent to retain only the term linear in  $r$  in the expression of the fluctuating force:

$$G(r, t) = rG(t) + h.o. \quad (36)$$

The function  $G(t)$  is related to the stochastic force  $F(x, t)$  via the relationship

$$G(t) = \frac{\partial}{\partial x} F(x, t)|_{x=x(t)} \quad (37)$$

In this approximation we recover our starting equation, eq. (33), but now formulated for increments. That means, that the solutions are given by a similar expression compared to eq. (2)

$$v(r, t) = a(t)r + \Lambda(t)\tilde{w}(\Lambda(t)r, \tau(t)) \quad (38)$$

provided we take  $\tilde{w}(\xi, \tau)$  as increment solutions, i.e.  $\tilde{w}(\xi, \tau) = w(x + \xi, \tau) - w(x, \tau)$ , where  $w(\xi, \tau)$  is a solution of the homogeneous Burgers equation.

Let us now briefly consider the treatment of the velocity increment statistics in the translationally invariant white noise forcing case, which has been performed by Polyakov on the basis of a hierarchy similar to eq. (26). The only difference in this hierarchy is the form of the matrix  $Q(r_i, r_j)$ . It is defined according to

$$Q(r_i, r_j) = q(r_i) + q(r_j) - q(0) - q(r_i - r_j) \quad (39)$$

Taking the approximation  $q(r) = q(0) - Q/2r^2$  the same hierarchy arises since in this approximation  $Q(r_i, r_j) = Qr_i r_j$ . Under this approximation the hierarchy of Polyakov is solved by the probability distributions given in eq. (24). However, as we have indicated above, the limit  $t \rightarrow \infty$  of the solution does not exist. Therefore, the evaluation of the stationary pdf's for the Polyakov problem has to go beyond the approximation of the force eq. (36). This indicates that all calculations which have been performed so far

for the translationally invariant white noise forced Burgers equation exclusively based on the approximation (35) should lead to the above probability distributions, which, as we have indicated above, have no stationary time limit.

Let us now finally consider the multidimensional forced Burgers equation:

$$\frac{\partial}{\partial t}\mathbf{v}(\mathbf{r}, t) + \mathbf{v}(\mathbf{r}, t) \cdot \nabla_r \mathbf{v}(\mathbf{r}, t) = \nu \Delta \mathbf{v}(\mathbf{r}, t) + G(t)\mathbf{r} \quad (40)$$

Here,  $G(t)$  denotes a time dependent matrix. We look for solutions of the form

$$\mathbf{v} = A(t)\mathbf{r} + \mathbf{U}(\mathbf{r}, t) \quad (41)$$

The matrix  $A(t)$  is determined in such a way as to cancel the force field. This leads to the matrix equation

$$\dot{A} + AA = G(t) \quad , \quad (42)$$

whereas the additional field  $\mathbf{U}(\mathbf{r}, t)$  has to fulfill the equation

$$\frac{\partial}{\partial t}\mathbf{U}(\mathbf{r}, t) + A \cdot \mathbf{U}(\mathbf{r}, t) + (A \cdot \mathbf{r}) \cdot \nabla \mathbf{U}(\mathbf{r}, t) + \mathbf{U}(\mathbf{r}, t) \cdot \nabla_r \mathbf{U}(\mathbf{r}, t) = \nu \Delta \mathbf{U}(\mathbf{r}, t) \quad (43)$$

We consider the ansatz

$$\mathbf{U}(\mathbf{r}, t) = \mathbf{W}(\Lambda(t) \cdot \mathbf{r}, t) \quad (44)$$

and assume the matrix  $\Lambda(t)$  to obey the equation

$$\dot{\Lambda}^T + A^T \Lambda^T = 0 \quad (45)$$

This leads to the equation

$$\frac{\partial}{\partial t}\mathbf{W}(\xi, t) + A \cdot \mathbf{W}(\xi, t) + \mathbf{W}(\xi, t) \cdot \Lambda^T \cdot \nabla_\xi \mathbf{W}(\xi, t) = \nu \nabla_\xi \Lambda \Lambda^T \cdot \nabla_\xi \mathbf{W}(\xi, t) \quad (46)$$

The ansatz

$$\mathbf{W}(\xi, t) = \Phi(t)\mathbf{w}(\xi, t) \quad (47)$$

where  $\Phi(t)$  has to obey the equation

$$\dot{\Phi}(t) + A(t)\Phi(t) = 0 \quad (48)$$

leads to

$$\frac{\partial}{\partial t}\mathbf{w}(\xi, t) + \mathbf{w}(\xi, t) \cdot \Phi^T \Lambda^T \cdot \nabla_\xi \mathbf{w}(\xi, t) = \nu \nabla_\xi \Lambda \Lambda^T \cdot \nabla_\xi \mathbf{w}(\xi, t) \quad (49)$$



The matrix  $\Phi(t)$ ,  $\Lambda(t)$  obeys the equation

$$\begin{aligned}\dot{\Phi} + A\Phi &= 0 \\ \dot{\Lambda} + \Lambda A &= 0\end{aligned}\tag{50}$$

We do not pursue the question further, what conditions have to be imposed on the matrix  $G(t)$  in order to allow for a complete reduction of equation (49) to the Burgers equation (possibly in a space with lower dimensions). The simplest case is clearly the case of a diagonal matrix

$$G(t) = g(t)E\tag{51}$$

Then, all matrices  $\Phi$ ,  $\Phi^T$ ,  $\Lambda$ ,  $\Lambda^T$  are diagonal matrices and, therefore, coincide. The matrices

$$\Lambda\Lambda^T = \Phi^T\Lambda^T = \lambda^2(t)E\tag{52}$$

are proportional to the unit matrix such that a transformation

$$\frac{d\tau(t)}{dt} = \lambda^2\tag{53}$$

actually defines  $\mathbf{w}(\xi, \tau)$  as the solution of the multidimensional Burgers equation. We mention that a similar treatment has been performed for the case of stretched vortices in hydrodynamic flows [7]

Summarizing we have considered a forced Burgers equation with a forcing linear in the spatial coordinate. We have shown that in the one dimensional case this problem can be solved completely. We have discussed the relationship to the velocity increment statistics of the Burgers equation with translationally invariant forcing. In the multidimensional case we have indicated the possibility of a similar treatment, which shows an analogy to the treatment of stretched vortices in hydrodynamic flows.

## References

- [1] J.M. Burgers, *The Nonlinear Diffusion Equation*, D. Reidel, Dordrecht 1974.
- [2] U. Frisch and J. Bec, *Les Houches 2000: New Trends in Turbulence*, M Lesieur, ed., Springer, Berlin (2001)
- [3] W.A. Woyczyński, *Burgers-KPZ Turbulence*, Springer, Berlin (1998)

- [4] E. Hopf, *The partial differential equation  $u_t + uu_x = u_{xx}$* , Comm. Pure Appl. Math. **3** (1950), 201–230.
- [5] J.D. Cole, *On a quasi-linear parabolic equation occurring in aerodynamics*, Quart. Appl. Math. **9** (1951), 225–236.
- [6] A.M. Polyakov, *Turbulence without pressure*, Phys. Rev. E **52** (1995), 6183–6188.
- [7] D.I. Pullin, P.G. Saffman, *Vortex dynamics in turbulence*, Anu. Rev. Fluid Mech. 30, 31 (1998)