

An exact solution to a k th order, n th degree nonlinear differential equation

C. Radhakrishnan Nair
Centre for Mathematical Sciences,
Thiruvananthapuram,
PIN - 695 581, INDIA

Abstract

An explicit analytic solution to the nonlinear differential equation $\left(\frac{d^k y}{dx^k}\right)^n = y^l$ is obtained for arbitrary integer values of k, l and n .

Introduction

Quantum mechanics, relativity and nonlinear physics are often pointed out [1] as the landmarks of the 20th century physics. The development of nonlinear physics occurred mostly in the last quarter of the 20th century. Chaos, Solitons and Fractals are the new disciplines that emerged out of the nonlinear physics. Industrial and technological exploits of the nonlinear phenomena are now common. Nonlinear differential equations is one of the tools for the study of nonlinear phenomena [2, 3]. Therefore, naturally more and more nonlinear differential equations are set up and methods of solving nonlinear differential equations are being vigorously pursued.

It is often very difficult to find exact solutions of nonlinear differential equations. The higher the nonlinearity, the greater the difficulty in getting an exact solution. Again, the higher the order of the nonlinear differential equation, the process of solving the nonlinear differential equation gets still more complicated. Owing to the lack of knowledge of solving the nonlinear differential equations exactly, more and more people try numerical methods to solve nonlinear differential equations. But no numerical solution, however great the accuracy, is a substitute for the analytical solutions. Therefore, even a small contribution to the analytical solution of differential equation is a welcome step.

The Nonlinear Differential Equation and its Solution

We consider the following nonlinear differential equation of the k th order and n th degree.

$$\left(\frac{d^k y}{dx^k}\right)^n = y^l \quad (1)$$

where k, l, n are integers and $l \neq n$.

Let us assume that a solution of the form

$$y = ax^m \quad (2)$$

exists, then

$$\frac{d^k y}{dx^k} = [am(m-1)(m-2)\dots\{m-(k-1)\}]x^{m-k}$$

and

$$y^l = a^l x^{ml}$$

substituting the values of $\frac{d^k y}{dx^k}$ and y^l into equation (1), we get

$$[\{am(m-1)(m-2)\dots(m-(k-1))\}x^{m-k}]^n = a^l x^{ml} \quad (3)$$

Equating the powers of x , on the left hand side and right hand side of equation (3), we get

$$\begin{aligned} n(m-k) &= ml \\ \therefore m &= \frac{nk}{n-l} \end{aligned} \quad (4)$$

Equating the coefficients of x ,

$$\begin{aligned} a^n m^n (m-1)^n (m-2)^n \dots \{m-(k-1)\}^n &= a^l \\ a &= [m(m-1)(m-2)\dots(m-(k-1))]^{\frac{n}{l-n}} \end{aligned} \quad (5)$$

Since a and m are known the solution

$$y = ax^m \text{ is known}$$

Illustration

By putting $k = 2$, $l = 4$ and $n = 6$ in (1), we get the equation

$$\left(\frac{d^2 y}{dx^2}\right)^6 = y^4 \quad (6)$$

From equation (5) $a = [6^6 \cdot 5^6]^{-1/2}$.

Substituting the values of m and a in the solution (2), we get

$$y = \frac{1}{\sqrt{6^6 5^6}} x^6 \quad (7)$$

It is verified that equation (7) is a solution of equation (6).

Discussion

The solution we found for equation (1) is simple. But the importance of this solution is that we can find the solution for any order and any degree of the differential equation. This implies that a very large class of physical phenomena can be studied using equation (1). If we know one of the solutions of a nonlinear differential equation, this knowledge may prove useful to generate some other solutions.

Acknowledgement

The author wishes to express his gratitude towards E. Krishnan and V. Vipindas for fruitful discussions and to L. A. Ajith for beautifully putting the material in \LaTeX . General support from Focal Image (India) Pvt Ltd was always a source of strength.

References

- [1] Grapnov-Grekhov A V and Rabinovich M I (1992) Nonlinearities in Action. Springer-Verlag, Berlin, pp. 1–9.
- [2] John D W and Smith P (1986) Nonlinear Ordinary Differential Equations—II edition, Clarendon Press, Oxford.
- [3] Dodd R K, Eilbeck J C, Gibben J D (1984) Solitons and Nonlinear wave equation, Academic, New York.