

# Multi Solitons of a Bose-Einstein Condensate in a Three-Dimensional Ring

L. Salasnich

*CNR-INFM, Research Unit of Milano University,  
Dipartimento di Fisica, Università di Milano,  
via Celoria 16, 20133 Milano, Italy  
E-mail: salasnich@mi.infm.it*

(Received 30 October 2005)

A Bose-Einstein Condensate (BEC) made of alkali-metal atoms at ultra-low temperatures is well described by the three-dimensional cubic nonlinear Schrödinger equation, the so-called Gross-Pitaevskii equation (GPE). Here we consider an attractive BEC in a ring and by solving the GPE we predict the existence of bright solitons with single and multi-peaks, showing that they have a limited domain of dynamical stability. We also discuss finite-temperature effects on the transition from the uniform phase to the localized phase.

**Key words:** Matter waves; Solitons in Bose-Einstein condensates

**PACS numbers:** 03.75.-b; 03.75.Lm

Single and multiple bright soliton configurations in a Bose Einstein condensates (BEC) have been observed in two different experiments [1, 2]. These solitons, which can travel for long distances without dispersion, have been theoretically investigated by various authors due to their relevance in nonlinear atom optics [3, 4]. Very interesting appears the not yet experimentally studied case of an attractive BEC in a ring. For this system it has been predicted, on the basis of the one-dimensional (1D) theory, a quantum phase transition from a uniform condensate to a bright soliton [5].

Here we consider an attractive BEC in a 3D ring, taking into account transverse variations of the BEC width. We show that the phase diagram of the system reveals novel and peculiar phenomena: the localized soliton has a limited existence and stability domain and the system supports also multi-peak solitons, which can be dynamically stable.

The time-dependent Gross-Pitaevskii equation (GPE) for a BEC in a trapping potential  $U(\mathbf{r})$  is given by

$$\left[ i\hbar \frac{\partial}{\partial t} + \frac{\hbar^2}{2m} \nabla^2 - U(\mathbf{r}) - \frac{4\pi\hbar^2 a_s (N-1)}{m} |\psi|^2 \right] \psi = 0$$

where  $N$  is the number of condensed atoms and  $a_s$  the s-wave scattering length of inter-atomic interaction. The GPE describes  $N$  trapped bosons in the same quantum state  $\psi(\mathbf{r}, t)$ : a pure BEC of  $N$  atoms. The GPE is a mean-field Hartree equation and it is accurate for a dilute gas of condensed bosons at ultra-low temperature. The cubic term is obtained by using a contact pseudo-potential  $V(\mathbf{r}, \mathbf{r}') = 4\pi\hbar^2 a_s / m \delta^3(\mathbf{r} - \mathbf{r}')$  to model the two-body interaction. If  $a_s > 0$  there is repulsion between particles, if  $a_s < 0$  there is attraction between particles [6].

We describe an attractive BEC in a cylinder, i.e. under transverse radial harmonic confinement and without axial longitudinal confinement, by using the self-focusing GPE ( $a_s < 0$ ) with  $\omega_\perp$  the transverse frequency of harmonic confinement. This self-focusing 3D GPE is the Euler-Lagrange equation of the following Lagrangian density

$$\mathcal{L} = \psi^* \left[ i \frac{\partial}{\partial t} + \frac{1}{2} \nabla^2 - \frac{1}{2} (x^2 + x^2) + \frac{1}{2} \Gamma |\psi|^2 \right] \psi$$

where  $\Gamma = 4\pi(N-1)|a_s|/a_\perp$  is the scaled inter-atomic strength. Here we use scaled units: energy

in units  $\hbar\omega_{\perp}$ , time in units of  $\omega_{\perp}^{-1}$ , length in units of  $a_{\perp} = \sqrt{\hbar/(m\omega_{\perp})}$ . To simplify the 3D problem, let us choose the trial wave function  $\psi(\mathbf{r}, t) = f(z, t)\exp[-(x^2 + y^2)/2\sigma(z, t)^2]/\pi\sigma(z, t)^2$ . Inserting it into the Lagrangian, integrating over  $x$  and  $y$  and neglecting the derivatives of  $\sigma(z, t)$  one finds an effective Lagrangian density  $\tilde{\mathcal{L}}$  which depends on the axial wave function  $f(z, t)$  and the transverse width  $\sigma(z, t)$ . The two Euler-Lagrange equations of the effective Lagrangian  $\tilde{\mathcal{L}}$  with respect to  $f$  and  $\sigma$  are

$$i\frac{\partial}{\partial t}f = \left[ -\frac{1}{2}\frac{\partial^2}{\partial z^2} + \frac{1}{2}\left(\frac{1}{\sigma^2} + \sigma^2\right) - \frac{g}{\sigma^2}|f|^2 \right] f$$

$$\sigma^2 = \sqrt{1 - g|f|^2}$$

where  $g = \Gamma/(2\pi) = 2(N-1)|a_s|/a_{\perp}$ . For  $\sigma = 1$  the first equation becomes the familiar self-focusing 1D GPE, also called cubic nonlinear Schrödinger equation. The equation for  $\sigma$  can be inserted into the equation for  $f$ . In this way one gets the time-dependent non-polynomial Schrödinger equation (NPSE) [7].

In [7] we have verified that the NPSE gives results always very close to that of the 3D GPE under transverse harmonic confinement for both positive and negative scattering length  $a_s$ . The self-focusing NPSE ( $a_s < 0$ ) admits a shape-invariant solution, called bright soliton, due to the interplay between the dispersive kinetic term and the attractive interaction term. Note that bright solitons of the self-focusing 3D GPE do not exist without transverse confinement. We have found [3, 4] that the bright soliton reduces, in the limit of very strong transverse confinement, namely for  $g \ll 1$  (1D regime), to the text-book bright soliton of the self-focusing 1D GPE, whose axial wave function is  $f(z) = \sqrt{g/4} \operatorname{sech}(gz/2)$ . Contrary to the self-focusing 1D GPE, the bright soliton of the self-focusing NPSE has a limited domain of existence:  $0 < g < \frac{4}{3}$ . If the strength  $g$  exceeds  $4/3$  there is the so-called collapse of the BEC. Thus, the transverse dynamics of the BEC induces the instability of the solitonic solution.

As previously stressed, bright solitons have been observed in two experiments: a single soliton in repulsive trap at the Ecole Normale Supérieure (ENS)

of Paris [2], and a soliton train in a harmonic trap at Rice University [1]. We believe that in the next future there will be experiments on attractive BEC in a toroidal trap. In fact, toroidal traps have been recently produced: at Georgia Tech cold atoms in a magnetic ring (diameter of 2 cm) [8] and at Berkeley a BEC in a magnetic ring (diameter of 3 mm) [9].

To model an attractive BEC in a toroidal trap we consider now the self-focusing NPSE where the solution  $f(z, t)$  has periodic boundary conditions:

$$f(z + L, t) = f(z, t)$$

where  $L = 2\pi R$  is the azimuthal longitudinal length of the BEC and  $R$  is the azimuthal radius of the trap. Due to the periodic boundary conditions the stationary self-focusing NPSE admits a uniform solution, given by  $f(z) = 1/\sqrt{L}$ . One sees immediately that the uniform solution exists only for  $g < 2\pi R$ . The stationary NPSE admits also nodeless and nodal localized solutions (solitonic solutions). In Fig. 1 we plot the existence diagram of these solitonic solutions in the  $(R, g)$  plane. The diagram is obtained by numerically solving the NPSE. The one-peak localized nodeless solution exists between the two solid lines of Fig. 1. Nodeless two-peak localized solution exists between the two dashed lines while the nodal two-peak localized solution exists between the two dot-dashed lines.

We analyze the dynamical stability of the solutions by numerically solving the time-dependent NPSE taking as initial condition the stationary localized solution  $f(z)$  with a very weak perturbation. In the left panels of Fig. 2 we plot the density profile  $\rho(z) = |f(z)|^2$  of the solitonic solutions. In the right panels we show the time-dependence of the mean squared widths  $\langle z^2 \rangle$  for the weakly perturbed solitonic solutions. A periodic oscillation means that the solitonic solution is dynamically stable. In the calculations we choose a ring axial length  $L = 15$  and an interaction strength  $g = 1$ . We have studied the dynamical stability for various initial conditions. The main results are the following: i) the uniform solution is stable only below the lower curve of existence of the one-peak soliton; ii) the one-peak soliton is stable where it exists; iii) the nodal two-




FIG. 1. Existence diagram in the  $(R, g)$  plane, where  $g = 2N|a_s|/a_\perp$  is the interaction coupling and  $R = L/(2\pi)$  is the azimuthal radius of the ring (in units  $a_\perp$ ). Uniform solution: exists for  $g < 2\pi R$ . One-peak localized nodeless solution: exists between the two solid lines. Nodeless two-peak localized solution exists between the two dashed lines. The nodal two-peak localized solution exists between the two dot-dashed lines.

peak soliton is stable in the plane  $(R, g)$  only below the upper curve of existence of the one-peak soliton; iv) the nodeless two-peak soliton is unstable. Similar results are found for solitonic solutions with a larger number of peaks.

To conclude, we observe that the critical strength  $g^* = \pi a_\perp/2R$  of the transition from the uniform state to the localized state at zero temperature has been determined in [5] by investigating the Bogoliubov energy spectrum of the self-focusing 1D GPE. The inclusion of finite-temperature effects is quite simple if one consider a non-interacting thermal cloud. In this way we find that the critical temperature  $T^*$  of transition from the uniform to the localized state is given by

$$\frac{k_B T^*}{\hbar\omega_\perp} = \frac{a_\perp^2}{4R^2} \left( N - \frac{\pi a_\perp^2}{4R|a_s|} \right)$$

where  $k_B$  is the Boltzmann constant, and using not-scaled quantities. Obviously if  $N < \pi a_\perp^2/(4R|a_s|)$  the system remains uniform at all temperatures.




FIG. 2. Left panels: density profile  $\rho(z)$  of the solitonic solutions. Right panels: time-dependence of the mean squared widths  $\langle z^2 \rangle$  for the weakly perturbed solitonic solutions. Ring axial length:  $L = 15$ . Interaction strength:  $g = 1$ .

## References

- [1] K.E. Strecker *et al.*, Nature **417**, 150 (2002).
- [2] L. Khaykovich *et al.*, Science **296**, 1290 (2002).
- [3] U. Al Khawaja *et al.*, Phys. Rev. Lett. **89**, 200404 (2002); L. Salasnich, A. Parola, and L. Reatto, Phys. Rev. A **66**, 043603 (2002); L. Salasnich, A. Parola, and L. Reatto, Phys. Rev. Lett. **91**, 080405 (2003); L.D. Carr and J. Brand, Phys. Rev. Lett. **92**, 040401 (2004).
- [4] L.D. Carr and Y. Castin, Phys. Rev. A **66**, 063602 (2002); L. Salasnich, Phys. Rev. A **70**, 053617 (2004).
- [5] G.M. Kavoulakis, Phys. Rev. A **67**, 011601(R) (2003); R. Kanamoto, H. Saito, and M. Ueda, Phys. Rev. A **67**, 013608 (2003).
- [6] L.P. Pitaevskii and S. Stringari, *Bose-Einstein Condensation* (Oxford University Press, Oxford, 2003).
- [7] L. Salasnich, Laser Phys. **12**, 198 (2002); L. Salasnich, A. Parola, and L. Reatto, Phys. Rev. A **65**, 043614 (2002).
- [8] J.A. Sauer, M.D. Barrett, and M.S. Chapman, Phys. Rev. Lett. **87**, 270401 (2001);
- [9] S. Gupta *et al.*, Phys. Rev. Lett. **95**, 143201 (2005).