The Sasa–Satsuma (complex mKdV II) and the complex sine-Gordon II equation revisited: recursion operators, nonlocal symmetries, and more

Artur Sergyeyev[†] and Dmitry Demskoi[‡]

[†] Mathematical Institute, Silesian University in Opava,

Na Rybníčku 1, 74601 Opava, Czech Republic

‡ School of Mathematics and Statistics,

The University of New South Wales,

Sydney, NSW 2052, Australia

E-mail: Artur.Sergyeyev@math.slu.cz and demskoi@maths.unsw.edu.au

July 1, 2021

We present a new symplectic structure and a hereditary recursion operator for the Sasa– Satsuma equation which is widely used in nonlinear optics. Using an integro-differential substitution relating this equation to a third-order symmetry flow of the complex sine-Gordon II equation enabled us to find a hereditary recursion operator and higher Hamiltonian structures for the latter equation.

We also show that both the Sasa–Satsuma equation and the third-order symmetry flow for the complex sine-Gordon II equation are bi-Hamiltonian systems, and we construct several hierarchies of local and nonlocal symmetries for these systems.

I Introduction

Finding a recursion operator for a system of PDEs is of paramount importance, as the whole integrable hierarchy for the system in question is then readily generated by the repeated application of the recursion operator to a suitably chosen seed symmetry, see e.g. [7, 32, 11] and references therein. Moreover, using formal adjoint of the recursion operator enables us to produce infinitely many conserved quantities for our hierarchy (see e.g. [7, 11] for further details). The recursion operators of multicomponent systems often have a richer structure of nonlocalities than in the one-component case, which makes such operators more difficult to find. For instance, there are two-component integrable generalizations, Eqs.(2) and (3), of the sine-Gordon equation and of the mKdV equation respectively. No recursion operator was found for Eqs.(2) and (3) so far even though the corresponding Lax pairs [36, 17, 38] and bilinear representations [17, 20, 18] are well known. The goal of the present paper is to fill this gap. Namely, we find and study the recursion operators for the so-called complex sine-Gordon II equation (2) and for the Sasa–Satsuma equation (3).

The Sasa–Satsuma equation [36] (see also [46, 44]) for a complex function U has the form

$$U_t = U_{xxx} + 6U\bar{U}U_x + 3U(U\bar{U})_x.$$
 (1)

Here and below the bar refers to the complex conjugate.

It is natural to refer to this equation as to the *complex mKdV II* as (1) is one of the two integrable complexifications of the famous modified KdV equation, the other complexification (complex mKdV I) being simply $U_t = U_{xxx} + 6U\bar{U}U_x$.

In turn, the complex sine-Gordon II equation [17, 38] is a hyperbolic PDE for a complex function ψ of the form

$$\psi_{xy} = \frac{\psi\psi_x\psi_y}{\psi\bar{\psi}+c} + (2\psi\bar{\psi}+c)(\psi\bar{\psi}+c)k\psi,$$

where c and k are arbitrary constants. The usual sine-Gordon equation $\phi_{xy} = c^2 k \sin(\phi)$ is recovered upon setting $\psi = \bar{\psi} = \sqrt{-c} \sin(\phi/4)$.

The Sasa–Satsuma and the complex sine-Gordon II equation are of considerable interest for applications. The Sasa–Satsuma equation is widely used in nonlinear optics, see e.g. [24, 34] and references therein, because the integrable cases of the so-called higher order nonlinear Schrödinger equation [25] describing the propagation of short pulses in optical fibers are related through a gauge transformation either to the Sasa–Satsuma equation [36] or to the so-called Hirota equation [19].

The complex sine-Gordon II equation, along with the Pohlmeyer–Lund–Regge model [33, 26, 16, 30] also known as the complex sine-Gordon I, define integrable perturbations of conformal field theories [12, 2, 8, 3]; see e.g. [35, 37] for other applications. Moreover, the complex sine-Gordon I and II are the only equations for one complex field in the plane for which the (multi)vortex solutions are found in closed form [31, 5, 6].

Following [21] we set $u = \psi$ and $v = \overline{\psi}$ and write the complex sine-Gordon II equation along with its complex conjugate as a system for u and v:

$$u_{xy} = \frac{vu_x u_y}{uv + c} + (2uv + c)(uv + c)ku, \qquad v_{xy} = \frac{uv_x v_y}{uv + c} + (2uv + c)(uv + c)kv.$$
(2)

Likewise, upon setting p = U and $q = \overline{U}$ in the Sasa–Satsuma equation, proceeding in the same fashion as above and writing out $(pq)_x$ as $pq_x + p_xq$, we obtain

$$p_t = p_{xxx} + 9pqp_x + 3p^2q_x, \quad q_t = q_{xxx} + 9pqq_x + 3q^2p_x.$$
(3)

From now on we shall treat u, v, p and q as independent variables that can be real or complex and consider systems (2) and (3) that are more general than the original complex sine-Gordon II and Sasa–Satsuma (complex mKdV II) equations which can be recovered under the reductions $v = \bar{u}$ and $q = \bar{p}$ respectively. In what follows we shall refer to (2) as to the *complex sine-Gordon II system* and to (3) as to the *Sasa–Satsuma system*.

Let us briefly address the relationship of the Sasa–Satsuma system (3) with other integrable systems. First of all, Eq.(3) can be obtained [46] as a reduction of the four-component Yajima– Oikawa system (8) from [46].

On the other hand, consider the vector modified KdV equation

$$V_t = V_{xxx} + \langle V, V \rangle V_x + \langle V, V_x \rangle V \tag{4}$$

studied by Svinolupov and Sokolov [42] (see also [43, 45]). Here $V = (V^1, \ldots, V^n)^T$ is an *n*-component vector, $\langle \cdot, \cdot \rangle$ stands for the usual Euclidean scalar product of two vectors, and the superscript *T* here and below indicates the transposed matrix. A simple linear change of variables $V^1 = \sqrt{3}(p+q)/\sqrt{2}$, $V^2 = i\sqrt{3}(p-q)/\sqrt{2}$ takes (3) into (4) with n = 2.

Setting $r = \sqrt{3}p$ and $s = \sqrt{3}q$ turns (3) into the system

$$r_t = r_{xxx} + r(3sr_x + rs_x),$$

$$s_t = s_{xxx} + s(3rs_x + sr_x),$$
(5)

studied by Foursov [15], who found a Hamiltonian structure for (5) of the form

$$\tilde{P} = \begin{pmatrix} -\frac{1}{3}rD_x^{-1} \circ r & D_x + \frac{1}{3}rD_x^{-1} \circ s \\ D_x + \frac{1}{3}sD_x^{-1} \circ r & -\frac{1}{3}sD_x^{-1} \circ s \end{pmatrix}$$
(6)

where D_x is the operator of total x-derivative, see e.g. [7, 11, 32] for details and for the background on the recursion operators, Hamiltonian and symplectic structures. The corresponding Hamiltonian density is [15] $(2/3)r^2s^2 - r_xs_x$. This immediately yields a Hamiltonian structure (8) for (3). Note that in [14, 15] there seems to be a misprint in \tilde{P} , and in (6) we corrected this misprint.

In [14, p.89] Foursov claims to have found some skew-symmetric operators that are likely to provide higher Hamiltonian structures for (5) but he failed to verify that these operators are indeed Hamiltonian. The explicit form of these operators was not presented in [15] or [14], so apparently it was never proved in the literature that (5) (and hence (3)) are bi-Hamiltonian systems. We establish the bi-Hamiltonian nature of (3) in Theorem 1 below.

Now turn to the complex sine-Gordon II system (2). Recall that Eq.(2) can be obtained (see e.g. [17, 21, 28]) as the Euler–Lagrange equation for the functional $S = \int L \, dx \, dy$, where

$$L = \frac{1}{2} \frac{u_x v_y + u_y v_x}{uv + c} + k(uv + c)uv.$$

A few conservation laws and generalized symmetries for (2) can be readily found e.g. using computer algebra [21, 28]. In particular [21], Eq.(2) is compatible with

$$u_{t} = u_{xxx} - \frac{3uv_{x}u_{xx}}{uv + c} - \frac{9u_{x}^{2}v_{x}}{uv + c} + \frac{3u^{2}v_{x}^{2}u_{x}}{(uv + c)^{2}},$$

$$v_{t} = v_{xxx} - \frac{3vu_{x}v_{xx}}{uv + c} - \frac{9v_{x}^{2}u_{x}}{uv + c} + \frac{3v^{2}u_{x}^{2}v_{x}}{(uv + c)^{2}}.$$
(7)

The compatibility here means that the flow (7) commutes with the nonlocal flow (14) associated with (2) or, equivalently, the right-hand sides of (7) constitute the characteristic of a third-order generalized symmetry for (2), see e.g. [32] for general background on symmetries.

The rest of paper is organized as follows. In section II we present symplectic structure and recursion operator for the Sasa–Satsuma system (3) and show that (3) is a bi-Hamiltonian system. In section III we employ a nonlocal change of variables relating systems (7) and (3) in order to construct recursion operator, Hamiltonian and symplectic structures for (2) and (7) from those of (3), and we show that (7) is a bi-Hamiltonian system. Finally, in section IV we discuss the hierarchies of local and nonlocal symmetries for (2), (3) and (7).

II Recursion operator and symplectic structure

for the Sasa–Satsuma system

A straightforward but tedious computation proves the following assertion.

Theorem 1 The Sasa–Satsuma system (3) possesses a Hamiltonian structure

$$\mathcal{P} = \begin{pmatrix} -pD_x^{-1} \circ p & D_x + pD_x^{-1} \circ q \\ D_x + qD_x^{-1} \circ p & -qD_x^{-1} \circ q \end{pmatrix},$$
(8)

a symplectic structure

$$\mathcal{J} = \begin{pmatrix} 3pD_x^{-1} \circ p & D_x + 5pD_x^{-1} \circ q \\ D_x + 5qD_x^{-1} \circ p & 3qD_x^{-1} \circ q \end{pmatrix},$$
(9)

and a hereditary recursion operator $\mathcal{R} = \mathcal{P} \circ \mathcal{J}$ that can be written as

$$\mathcal{R} = \begin{pmatrix} D_x^2 + 6pq + q_x D^{-1} \circ p + bD_x^{-1} \circ p + 3p_x D_x^{-1} \circ b & 2p^2 - 2z_1 p D_x^{-1} \circ q - 3p_x D_x^{-1} \circ a \\ 2q^2 - 2z_2 q D_x^{-1} \circ p + 3q_x D_x^{-1} \circ b & D_x^2 + 6pq + p_x D^{-1} \circ q + a D_x^{-1} \circ q - 3q_x D_x^{-1} \circ a \end{pmatrix}$$
(10)

where $a = p_x + 2z_1q$, $b = q_x + 2z_2p$; $z_1 = D_x^{-1}(p^2)$ and $z_2 = D_x^{-1}(q^2)$ are potentials for the following conservation laws of (3):

$$D_t(p^2) = D_x(2pp_{xx} - p_x^2 + 6p^3q), \quad D_t(q^2) = D_x(2qq_{xx} - q_x^2 + 6pq^2).$$

Hence (3) has an infinite hierarchy of compatible Hamiltonian structures $\mathcal{P}_k = \mathcal{R}^k \circ \mathcal{P}$, $k = 0, 1, 2, ..., \mathcal{P}_0 \equiv \mathcal{P}$, an infinite hierarchy of symplectic structures $\mathcal{J}_k = \mathcal{J} \circ \mathcal{R}^k$, k = 0, 1, 2, ...,and an infinite hierarchy of commuting symmetries of the form $\mathcal{K}_i = \mathcal{R}^i(\mathcal{K}_0)$, i = 0, 1, 2, ..., where $\mathcal{K}_0 = (p_x, q_x)^T$.

The Sasa-Satsuma system (3) is bi-Hamiltonian with respect to \mathcal{P}_0 and \mathcal{P}_1 :

$$\begin{pmatrix} p_t \\ q_t \end{pmatrix} = \mathcal{P}_0 \begin{pmatrix} \delta \mathcal{H}_1 / \delta p \\ \delta \mathcal{H}_1 / \delta q \end{pmatrix} = \mathcal{P}_1 \begin{pmatrix} \delta \mathcal{H}_0 / \delta p \\ \delta \mathcal{H}_0 / \delta q \end{pmatrix},$$

where $\mathcal{H}_i = \int H_i dx$, $i = 0, 1, H_0 = pq, H_1 = 2p^2q^2 - p_xq_x$.

Here $\delta/\delta p$ and $\delta/\delta q$ denote variational derivatives with respect to p and q.

Note that the Hamiltonian structure \mathcal{P} above is nothing but the second Hamiltonian structure of the AKNS system, see e.g. [27]. This Hamiltonian structure (more precisely, its counterpart (6) for Eq.(5)) has already appeared in [14, 15].

The symmetries \mathcal{K}_i commute because \mathcal{R} is hereditary. Applying \mathcal{R} to an obvious symmetry $\mathcal{K}_0 = (p_x, q_x)^T$ of (3) yields the symmetry $\mathcal{K}_1 = \mathcal{R}(\mathcal{K}_0) = (p_{xxx} + 9pqp_x + 3p^2q_x, q_{xxx} + 9pqq_x + 3q^2p_x)^T$, i.e., the right-hand side of (3). In turn, $\mathcal{R}(\mathcal{K}_1)$ is a *local* fifth-order symmetry for (3). We guess that \mathcal{K}_i are local for all natural *i* but so far we were unable to provide a rigorous proof of this.

Unlike the overwhelming majority of the hitherto known recursion operators, see e.g. discussion in [39] and references therein, the nonlocal variables appear explicitly in the coefficients of \mathcal{R} . Perhaps this is the very reason why \mathcal{R} was not found earlier. The nonlocal variables in the coefficients of \mathcal{R} are *abelian* pseudopotentials as in [23] and unlike e.g. the nonlocalities in the recursion operator discovered by Karasu et al. [22] and later rewritten in [40]: the nonlocalities in the operator from [22, 40] are *nonabelian* pseudopotentials.

III Recursion operator

for the complex sine-Gordon II system

There is a well-known (see e.g. [1] for discussion and references) transformation $z = \sqrt{2/3}g_x$ relating the symmetry flow $g_t = g_{xxx} + g_x^3/2$ of the sine-Gordon equation $g_{xy} = \sin(g)$ and the mKdV equation $z_t = z_{xxx} + z^2 z_x$. Moreover, there exists [1] a nonlocal generalization of this transformation that sends the third-order symmetry flow

$$u_{t} = u_{xxx} - 3\frac{uv_{x}u_{xx}}{uv + c} + 3\frac{(-uvu_{x} - cu_{x} + u^{2}v_{x})u_{x}v_{x}}{(uv + c)^{2}},$$
$$v_{t} = v_{xxx} - 3\frac{vu_{x}v_{xx}}{uv + c} - 3\frac{(-v^{2}u_{x} + uvv_{x} + cv_{x})u_{x}v_{x}}{(uv + c)^{2}}$$

of the complex sine-Gordon I equation (see e.g. [16, 21, 28])

$$u_{xy} = \frac{vu_x u_y}{uv + c} + (uv + c)ku, \qquad v_{xy} = \frac{uv_x v_y}{uv + c} + (uv + c)kv, \tag{11}$$

into the two-component generalization of the mKdV equation

$$p_t = p_{xxx} + 6pqp_x, \qquad q_t = q_{xxx} + 6pqq_x$$

that belongs to the hierarchy of the well-known AKNS system

$$p_t = p_{xx} + p^2 q, \qquad q_t = -q_{xx} - q^2 p.$$
 (12)

Note that this nonlocal transformation also sends [1] the second-order symmetry flow

$$u_t = u_{xx} - 2\frac{uu_x v_x}{uv + c}, \qquad v_t = -v_{xx} + 2\frac{vu_x v_x}{uv + c}$$

of (11) into the AKNS system (12).

It turns out that upon a suitable redefinition of nonlocal variables the nonlocal transformation in question in combination with a suitable rescaling of dependent variables p and q also sends the third-order symmetry flow (7) of the complex sine-Gordon II equation (2) into the Sasa–Satsuma system (3). Namely, we have the following result.

Theorem 2 The substitution

$$p = \frac{\mathrm{i}\sqrt{2}u_x \exp(-\frac{1}{2}w_1)}{\sqrt{uv+c}}, \quad q = \frac{\mathrm{i}\sqrt{2}v_x \exp(\frac{1}{2}w_1)}{\sqrt{uv+c}},$$

where $i = \sqrt{-1}$, $w_1 = D_x^{-1}(\rho_1)$ is a potential for the conservation law $D_t(\rho_1) = D_x(\sigma_1)$ of (7),

$$\rho_{1} = \frac{uv_{x} - vu_{x}}{uv + c}, \quad \sigma_{1} = \frac{uv_{xxx} - vu_{xxx}}{uv + c} + \frac{(3uv + 2c)(v_{x}u_{xx} - u_{x}v_{xx})}{(uv + c)^{2}} + \frac{(12uv + 11c)(vu_{x} - uv_{x})u_{x}v_{x}}{(uv + c)^{3}},$$

takes Eq.(7) into the Sasa-Satsuma system (3).

Here and below D_x and D_t stand for the total x- and t-derivatives as defined e.g. in [32]. Remark 1 Let

$$\tilde{p} = \frac{\mathrm{i}\sqrt{2}u_x \exp(-\frac{1}{2}w_1)}{\sqrt{uv+c}}, \quad \tilde{q} = \frac{\mathrm{i}\sqrt{2}v_x \exp(\frac{1}{2}w_1)}{\sqrt{uv+c}},$$

where $w_1 = D_x^{-1}(\rho_1)$ is now a potential for the conservation law $D_y(\rho_1) = D_x(\theta_1)$ of (2), ρ_1 is given above and

$$\theta_1 = \frac{vu_y - uv_y}{uv + c}$$

Then for k = 0 we have

$$D_y(\tilde{p}) = 0, \qquad D_y(\tilde{q}) = 0,$$

where D_y stands for the total y-derivative. In other words, if k = 0 then \tilde{p} and \tilde{q} provide nonlocal y-integrals for (2), and $D_x \ln(\tilde{p})$ and $D_x \ln(\tilde{q})$ are local y-integrals of (2). Therefore, the complex sine-Gordon II system (2) for k = 0 is Liouvillean and C-integrable, see [9, 10] for the construction of symmetries of such systems using the integrals thereof. Using the above local integrals enables us to find the general solution for (2) with k = 0 along the lines of [47].

Passing from p and q to u and v yields from \mathcal{R} a recursion operator \mathfrak{R}_0 for (7). The operator \mathfrak{R}_0 is hereditary because so is \mathcal{R} , see [13]. It is easily seen that \mathfrak{R}_0 is a recursion operator for (2) as well. Upon removing an inessential overall constant factor in \mathfrak{R}_0 we obtain the following result:

Theorem 3 The complex sine-Gordon II system (2) has a hereditary recursion operator

$$\begin{split} \Re &= \left(\begin{array}{cc} D_x^2 - \frac{2uv_x}{uv+c} + \frac{uv_{xx} - 12u_xv_x}{uv+c} + \frac{cu_xv_x}{(uv+c)^2} & \frac{-2uu_{xx} + 6u_x^2}{uv+c} + \frac{4u^2u_xv_x}{(uv+c)^2} \\ & \frac{-2vv_{xx} + 6v_x^2}{uv+c} + \frac{4v^2u_xv_x}{(uv+c)^2} & D_x^2 - \frac{2vu_x}{uv+c} + \frac{vu_{xx} - 12u_xv_x}{uv+c} + \frac{cu_xv_x}{(uv+c)^2} \end{array} \right) \\ & + \left(\begin{array}{c} \sum\limits_{\alpha=1}^5 Q_\alpha^1 D_x^{-1} \circ \gamma_{\alpha,1} & \sum\limits_{\alpha=1}^5 Q_\alpha^1 D_x^{-1} \circ \gamma_{\alpha,2} \\ \\ \sum\limits_{\alpha=1}^5 Q_\alpha^2 D_x^{-1} \circ \gamma_{\alpha,1} & \sum\limits_{\alpha=1}^5 Q_\alpha^2 D_x^{-1} \circ \gamma_{\alpha,2} \end{array} \right), \end{split}$$

where w_1 is as in Remark 1; $y_1 = D_x^{-1}\left((u_x^2 \exp(-w_1))/(uv+c)\right)$ and $y_2 = D_x^{-1}\left((v_x^2 \exp(w_1))/(uv+c)\right)$ are potentials for the nonlocal conservation laws $D_y(\zeta_i) = D_x(\xi_i)$, i = 1, 2, of(2),

$$\zeta_1 = \frac{u_x^2 \exp(-w_1)}{uv + c}, \quad \xi_1 = ku^2(uv + c) \exp(-w_1),$$

$$\zeta_2 = \frac{v_x^2 \exp(w_1)}{uv + c}, \quad \xi_2 = kv^2(uv + c) \exp(w_1);$$

$$\begin{aligned} \boldsymbol{Q}_{1} &= \begin{pmatrix} y_{1}u \\ u_{x}\exp(-w_{1}) - vy_{1} \end{pmatrix}, \quad \boldsymbol{Q}_{2} &= \begin{pmatrix} v_{x}\exp(w_{1}) - uy_{2} \\ y_{2}v \end{pmatrix}, \\ \boldsymbol{Q}_{3} &= \begin{pmatrix} u \\ -v \end{pmatrix}, \quad \boldsymbol{Q}_{4} &= c \begin{pmatrix} -u_{xx} + \frac{2uu_{x}v_{x}}{uv + c} + 4\exp(w_{1})y_{1}v_{x} - 4uy_{2}y_{1} \\ v_{xx} - \frac{2vu_{x}v_{x}}{uv + c} - 4\exp(w_{1})y_{2}u_{x} + 4vy_{2}y_{1} \end{pmatrix}, \quad \boldsymbol{Q}_{5} &= \begin{pmatrix} u_{x} \\ v_{x} \end{pmatrix} \end{aligned}$$

are symmetries for (2), and

$$\begin{split} \gamma_1 &= \left(\frac{\exp(w_1)vv_x^2 + cy_2v_x}{(uv+c)^2}, \frac{\exp(w_1)v_{xx}}{uv+c} - \frac{2\exp(w_1)vu_xv_x + cy_2u_x}{(uv+c)^2}\right), \\ \gamma_2 &= \left(\frac{\exp(-w_1)u_{xx}}{uv+c} - \frac{2\exp(-w_1)uu_xv_x + cy_1v_x}{(uv+c)^2}, \frac{\exp(-w_1)uu_x^2 + cy_1u_x}{(uv+c)^2}\right) \\ \gamma_3 &= \left(-\frac{v_{xxx}}{uv+c} + \frac{2(vv_x+2\exp(-w_1)uvy_2 + 2\exp(-w_1)cy_2)u_{xx} + (2vu_x + uv_x)v_{xx}}{(uv+c)^2} + \frac{2v_xu_x(-v^2u_x + 2cv_x)}{(uv+c)^3} - \frac{2v_x(-u_xv_x + 4\exp(-w_1)uy_2u_x + 2\exp(w_1)vy_1v_x + 2cy_2y_1)}{(uv+c)^2}, \\ \frac{u_{xxx}}{uv+c} - \frac{2(uu_x+2\exp(w_1)uvy_1 + 2\exp(w_1)cy_1)v_{xx} - (2uv_x + vu_x)u_{xx}}{(uv+c)^2} - \frac{2v_xu_x(-u^2v_x + 2cu_x)}{(uv+c)^3} + \frac{2u_x(-u_xv_x + 4\exp(w_1)vy_1v_x + 2\exp(-w_1)uy_2u_x + 2cy_2y_1)}{(uv+c)^2} \right) \\ \gamma_4 &= \left(\frac{v_x}{(uv+c)^2}, -\frac{u_x}{(uv+c)^2}\right), \quad \gamma_5 = \left(\frac{v_{xx}}{(uv+c)} - \frac{uv_x}{(uv+c)^2}, \frac{u_{xx}}{(uv+c)} - \frac{vu_x}{(uv+c)^2}\right) \end{split}$$

are cosymmetries for (2).

Here Q_{α}^{i} and $\gamma_{\alpha,j}$ denote i^{th} component of \boldsymbol{Q}_{α} and j^{th} component of $\boldsymbol{\gamma}_{\alpha}$ respectively, i.e.,

$$oldsymbol{Q}_{lpha} \equiv \left(egin{array}{c} Q_{lpha}^1 \ Q_{lpha}^2 \end{array}
ight), \quad oldsymbol{\gamma}_{lpha} \equiv (\gamma_{lpha,1},\gamma_{lpha,2}),$$

Note that if we use the tensorial notation (see e.g. [28]), we can rewrite the operator

$$\left(\begin{array}{c}\sum_{\alpha=1}^{5}Q_{\alpha}^{1}D_{x}^{-1}\circ\gamma_{\alpha,1} & \sum_{\alpha=1}^{5}Q_{\alpha}^{1}D_{x}^{-1}\circ\gamma_{\alpha,2}\\ & \sum_{\alpha=1}^{5}Q_{\alpha}^{2}D_{x}^{-1}\circ\gamma_{\alpha,1} & \sum_{\alpha=1}^{5}Q_{\alpha}^{2}D_{x}^{-1}\circ\gamma_{\alpha,2}\end{array}\right)$$

in a more concise form, namely $\sum_{\alpha=1}^{5} \boldsymbol{Q}_{\alpha} \otimes D_{x}^{-1} \circ \boldsymbol{\gamma}_{\alpha}$.

Remark 2 Eq.(7) has a recursion operator of precisely the same form as the \Re given above, but in this case the nonlocal variables should be defined in a slightly different way: w_1 should be as in Theorem 2, and $y_1 = D_x^{-1}((u_x^2 \exp(-w_1))/(uv+c)), y_2 = D_x^{-1}((v_x^2 \exp(w_1))/(uv+c))$ should now be potentials for the nonlocal conservation laws $D_t(\zeta_i) = D_x(\chi_i), i = 1, 2, of$ (7), where ζ_i are as in Theorem 1 and

$$\chi_{1} = \exp(-w_{1}) \left(\frac{2u_{x}u_{xxx}}{uv+c} - \frac{u_{xx}^{2}}{uv+c} - \frac{2uu_{x}v_{x}u_{xx}}{(uv+c)^{2}} - \frac{2uu_{x}^{2}v_{xx}}{(uv+c)^{2}} - \frac{2(6uv+7c)u_{x}^{3}v_{x}}{(uv+c)^{3}} + \frac{3u^{2}u_{x}^{2}v_{x}^{2}}{(uv+c)^{3}} \right),$$

$$\chi_{2} = \exp(w_{1}) \left(\frac{2v_{x}v_{xxx}}{uv+c} - \frac{v_{xx}^{2}}{uv+c} - \frac{2vv_{x}^{2}u_{xx}}{(uv+c)^{2}} - \frac{2vu_{x}v_{x}v_{xx}}{(uv+c)^{2}} + \frac{3v^{2}u_{x}^{2}v_{x}^{2}}{(uv+c)^{3}} - \frac{2(6uv+7c)u_{x}v_{x}^{3}}{(uv+c)^{3}} \right).$$

Recall that Eq.(2) possesses [28, 29] a local symplectic structure

$$\mathfrak{J} = \begin{pmatrix} 0 & \frac{1}{uv+c}D_x - \frac{uv_x}{(uv+c)^2} \\ \frac{1}{uv+c}D_x - \frac{vu_x}{(uv+c)^2} & 0 \end{pmatrix},$$

i.e., Eq.(2) can be written in the form $\mathfrak{J}\boldsymbol{u}_y = k(2uv+c)(v,u)^T = \delta \mathcal{H}/\delta \boldsymbol{u}$, where $\boldsymbol{u} = (u,v)^T$ and $\mathcal{H} = \int k(uv+c)uvdx$. The symplectic structure \mathfrak{J} is readily seen to be shared by Eq.(7).

Inverting \mathfrak{J} yields a nonlocal Hamiltonian structure for (7) of the form

$$\mathfrak{P} = \begin{pmatrix} 0 & \exp(-w_1/2)\sqrt{uv+c} \circ D_x^{-1} \circ \exp(w_1/2)\sqrt{uv+c} \\ \exp(w_1/2)\sqrt{uv+c} \circ D_x^{-1} \circ \exp(-w_1/2)\sqrt{uv+c} & 0 \end{pmatrix},$$

where w_1 should be interpreted as in Theorem 2. It can be shown that if we go from the (u, v)- to the (p, q)-variables using the transformation from Theorem 2 then we obtain from \mathfrak{P} the Hamiltonian structure (8) for the Sasa–Satsuma system up to an inessential overall constant factor.

The higher Hamiltonian (resp. symplectic) structures for (7) are given by the formulas $\mathfrak{P}_n = \mathfrak{R}^n \circ \mathfrak{P}$ (resp. $\mathfrak{J}_n = \mathfrak{J} \circ \mathfrak{R}^n$), where $n = 1, 2, 3, \ldots$ The operator \mathfrak{R} is hereditary, and hence all of these structures are compatible. However, the structures \mathfrak{P}_1 and \mathfrak{J}_1 are already very cumbersome, so we do not display them here. Nevertheless, it is readily checked that the following assertion holds.

Theorem 4 Eq. (7) is a bi-Hamiltonian system:

$$\begin{pmatrix} u_t \\ v_t \end{pmatrix} = \mathfrak{P}_0 \begin{pmatrix} \delta \mathfrak{H}_1 / \delta u \\ \delta \mathfrak{H}_1 / \delta v \end{pmatrix} = \mathfrak{P}_1 \begin{pmatrix} \delta \mathfrak{H}_0 / \delta u \\ \delta \mathfrak{H}_0 / \delta v \end{pmatrix},$$

where $\mathfrak{H}_i = \int \tilde{H}_i dx$, i = 0, 1,

$$\tilde{H}_0 = -\frac{u_x v_x}{uv+c}, \quad \tilde{H}_1 = \frac{u_{xx} v_{xx}}{uv+c} - \frac{v u_x v_x u_{xx} + u u_x v_x v_{xx} - 4u_x^2 v_x^2}{(uv+c)^2} + \frac{u v u_x^2 v_x^2}{(uv+c)^3}.$$

It is straightforward to verify that the action of \mathfrak{R} on the obvious symmetry u_x of (7) yields the right-hand side of (7), and $\mathfrak{R}^2(u_x)$ is a fifth-order *local* generalized symmetry for (2). We guess that the repeated application of \mathfrak{R} to u_x yields a hierarchy of *local* generalized symmetries for (2) but we were not able to prove this in full generality so far, as presence of the nonlocalities y_i in the coefficients of \mathfrak{R} appears to render useless the hitherto known ways of proving locality for hierarchies of symmetries generated by the recursion operator, cf. e.g. [41] and references therein. Nevertheless, as \mathfrak{R} is a recursion operator for (7), Proposition 2 from [39] tells us that the only nonlocalities that could possibly appear in the hierarchy of the symmetries $\mathfrak{R}^k(u_x)$, $k = 1, 2, \ldots$, are potentials of (possibly nonlocal) conservation laws for (2).

IV Nonlocal symmetries

First of all, we can consider Eq.(2) as a nonlocal symmetry flow of Eq.(7). We have already noticed above that Eq.(2) can be written as

$$\mathfrak{J}\boldsymbol{u}_y = \delta\mathfrak{H}/\delta\boldsymbol{u},\tag{13}$$

where $\mathfrak{H} = \int k(uv+c)uvdx$. Acting by $\mathfrak{P} = \mathfrak{J}^{-1}$ on both sides of (13) we can formally rewrite the complex sine-Gordon II system (2) in the evolutionary form $u_y = \mathfrak{P}(\delta \mathfrak{H} / \delta u)$, that is,

$$u_y = k \exp(-w_1/2)\sqrt{uv + c} \ \omega_2, \quad v_y = k \exp(w_1/2)\sqrt{uv + c} \ \omega_1,$$
 (14)

where $\omega_1 = D_x^{-1} \left(v(2uv+c) \exp(-w_1/2)\sqrt{uv+c} \right)$ and $\omega_2 = D_x^{-1} \left(u(2uv+c) \exp(w_1/2)\sqrt{uv+c} \right)$ are potentials for the following nonlocal conservation laws of (7):

$$D_t \left(v(2uv+c) \exp(-w_1/2)\sqrt{uv+c} \right) = D_x \left(\exp(-w_1/2)\sqrt{uv+c} \left(\frac{(4uv+3c)v^2 u_{xx}}{uv+c} + (4uv+c)v_{xx} - \frac{2v^3 u_x^2}{uv+c} - \frac{v(20u^2v^2 + 26cuv + 7c^2)u_x v_x}{(uv+c)^2} - 2uv_x^2 \right) \right),$$

$$D_t \left(u(2uv+c) \exp(w_1/2)\sqrt{uv+c} \right) = D_x \left(\exp(w_1/2)\sqrt{uv+c} \left((4uv+c)u_{xx} + \frac{u^2(4uv+3c)v_{xx}}{uv+c} - \frac{2v^3 v_x^2}{uv+c} - \frac{2v^3 v_x^2}{uv+c} \right) \right).$$

Eq.(14) is compatible with Eq.(7), i.e., the right-hand side of (14) divided by k,

$$\boldsymbol{Q}_{-1} = \left(\begin{array}{c} \exp(-w_1/2)\sqrt{uv+c} \ \omega_2 \\\\ \exp(w_1/2)\sqrt{uv+c} \ \omega_1 \end{array} \right),$$

is a nonlocal symmetry for (7). Moreover, we have $\Re(\mathbf{Q}_{-1}) = c^2 \mathbf{u}_x$, i.e., the action of \Re on \mathbf{Q}_{-1} gives the 'zeroth' (obvious) symmetry \mathbf{u}_x up to a constant factor. Thus, Eq.(14) (and hence the complex sine-Gordon II system (2)) can be considered as a first negative flow in the hierarchy of (7). Note that if we consider \Re as a recursion operator for Eq.(2), we have $\Re(\mathbf{u}_y) = kc^2 \mathbf{u}_x$, in perfect agreement with the above result.

Eqs.(3) and (2) possess nonlocal symmetries of the form

$$\mathcal{G}_1 = \begin{pmatrix} q_x + z_2 p \\ -z_2 q \end{pmatrix}, \quad \mathcal{G}_2 = \begin{pmatrix} -z_1 p \\ p_x + z_1 q \end{pmatrix},$$

and

$$\boldsymbol{Q}_1 = \begin{pmatrix} y_1 u \\ u_x \exp(-w_1) - v y_1 \end{pmatrix}, \quad \boldsymbol{Q}_2 = \begin{pmatrix} v_x \exp(w_1) - u y_2 \\ y_2 v \end{pmatrix},$$

respectively, where z_1 and z_2 were defined in Theorem 1. Therefore we have four hierarchies of nonlocal symmetries: $\mathcal{R}^i(\mathcal{G}_j)$ for (3) and $\mathfrak{R}^i(\mathbf{Q}_j)$ for (2), where in both cases j = 1, 2 and i = 0, 1, 2, ...

Two more hierarchies of nonlocal symmetries have the form $\mathcal{R}^i(\mathcal{G}_0)$ for (3) and $\mathfrak{R}^i(\mathcal{Q}_0)$ for (2), where $i = 1, 2, \ldots$,

$$\mathcal{G}_0 = \begin{pmatrix} p \\ -q \end{pmatrix}, \quad \mathcal{Q}_0 = \begin{pmatrix} u \\ -v \end{pmatrix}.$$

Two somewhat more 'usual' nonlocal hierarchies of master symmetries (the latter are, roughly speaking, time-dependent symmetries such that repeatedly commuting them with a suitable timeindependent symmetry yields an infinite hierarchy of *time-independent* commuting symmetries for the system in question, see e.g. [7] and references therein for further details), $\mathcal{R}^{i}(\mathcal{S}_{0})$ for (3) and $\mathfrak{R}^{i}(\mathcal{S}_{0})$ for (7), where i = 1, 2, ..., originate from the scaling symmetries for (3) and (7),

$$S_{0} = \begin{pmatrix} 3t(p_{xxx} + 9pqp_{x} + 3p^{2}q_{x}) + xp_{x} + p \\ 3t(q_{xxx} + 9pqq_{x} + 3q^{2}p_{x}) + xq_{x} + q \end{pmatrix} \text{ and} \\ S_{0} = \begin{pmatrix} 3t\left(u_{xxx} - \frac{3uv_{x}u_{xx}}{uv + c} - \frac{9u_{x}^{2}v_{x}}{uv + c} + \frac{3u^{2}v_{x}^{2}u_{x}}{(uv + c)^{2}}\right) + xu_{x} \\ 3t\left(v_{xxx} - \frac{3vu_{x}v_{xx}}{uv + c} - \frac{9v_{x}^{2}u_{x}}{uv + c} + \frac{3v^{2}u_{x}^{2}v_{x}}{(uv + c)^{2}}\right) + xv_{x} \end{pmatrix},$$

respectively.

Finally, (7) has nonlocal symmetries

$$\boldsymbol{G}_{1} = \begin{pmatrix} \sqrt{uv+c} \exp(-w_{1}/2) \\ 0 \end{pmatrix}, \quad \boldsymbol{G}_{2} = \begin{pmatrix} 0 \\ \sqrt{uv+c} \exp(w_{1}/2) \end{pmatrix},$$

obtained by differentiating Q_{-1} with respect to ω_1 and ω_2 , but these symmetries are annihilated by \Re and hence do not lead to new hierarchies of nonlocal symmetries.

It would be interesting to find out whether the systems in question possess nonlocal symmetries that do not belong to the above hierarchies, which is the form of solutions invariant under the nonlocal symmetries and whether these solutions could have any applications in nonlinear optics.

As a final remark, we note that because of the obvious symmetry of the complex sine-Gordon II system (2) under the interchange of x and y, all of the above results concerning the recursion operator, Hamiltonian and symplectic structures, and hierarchies of symmetries for (2) remain valid if we replace all x-derivatives with y-derivatives and vice versa and swap the operators D_x and D_y . Interestingly enough, the recursion operator $\tilde{\mathfrak{R}}$ obtained from \mathfrak{R} upon such an interchange proves to be inverse to \mathfrak{R} on symmetries of (2) up to a constant factor. More precisely, for any symmetry K of (2) we have

$$\widetilde{\mathfrak{R}}(\boldsymbol{K}) = k^2 c^4 \mathfrak{R}^{-1}(\boldsymbol{K}).$$

Taking into account our earlier results we see that the 'basic' hierarchy of (2) can be represented by a diagram of the form

$$\mathbf{u} \cdots \to \mathfrak{R}^{-2}(\boldsymbol{u}_y) \to \mathfrak{R}^{-1}(\boldsymbol{u}_y) \to \boldsymbol{u}_y \to \boldsymbol{u}_x = \mathfrak{R}(\boldsymbol{u}_y)/kc^2 \to \mathfrak{R}(\boldsymbol{u}_x) \to \mathfrak{R}^2(\boldsymbol{u}_x) \to \cdots,$$

and we guess that *all* symmetries presented at this diagram are *local*, i.e., they do not involve nonlocal variables. This is a fairly common situation for hyperbolic PDEs, see e.g. the discussion in [4].

Acknowledgments

The research of A.S. was partially supported by the Czech Grant Agency (GACR) under grant 201/04/0538, and by the Ministry of Education, Youth and Sports of the Czech Republic (MŠMT ČR) under grant MSM 4781305904.

References

- H. Aratyn, L.A. Ferreira, J.F. Gomes and A.H. Zimerman, The complex sine-Gordon equation as a symmetry flow of the AKNS hierarchy, J. Phys. A: Math. Gen. 33 (2000), L331–L337 (*Preprint* nlin.SI/0007002)
- [2] I. Bakas, Conservation laws and geometry of perturbed coset models, Int. J. Mod. Phys. A 9 (1994), 3443–3472; Q-H. Park, Deformed coset models from gauged WZW actions, Phys. Lett. B 328 (1994), 329–336

- [3] I. Bakas and J. Sonnenschein, On integrable models from pp-wave string backgrounds, J. High Energy Phys. 12 (2002), 049 (*Preprint* hep-th/0211257)
- [4] H. Baran, Can we always distinguish between positive and negative hierarchies?, J. Phys. A: Math. Gen. 38 (2005), L301–L306
- [5] I.V. Barashenkov and D.E. Pelinovsky, Exact vortex solutions of the complex sine-Gordon theory on the plane, Phys. Lett. B 436 (1998), 117–124 (*Preprint* hep-th/9807045)
- [6] I.V. Barashenkov, V.S. Shchesnovich, and R.M. Adams, Noncoaxial multivortices in the complex sine-Gordon theory on the plane, Nonlinearity 15 (2002), 2121–2146 (*Preprint nlin.SI/0202018*)
- [7] M. Błaszak, Multi-Hamiltonian Theory of Dynamical Systems (Springer, Heidelberg, 1998).
- [8] V.A. Brazhnikov, $\Phi^{(2)}$ perturbations of the WZW model, Nucl. Phys. **B 501** (1997), 685–710
- [9] D.K. Demskoi, S.Ya. Startsev, On construction of symmetries from integrals of hyperbolic partial differential systems, J. Math. Sci. 136 (2006), 4378–4374.
- [10] D.K. Demskoi, On a class of Liouville-type systems. Theor. Math. Phys. 141 (2004), 1509–1527
- [11] I. Dorfman, Dirac Structures and Integrability of Nonlinear Evolution Equations (John Wiley & Sons, Chichester, 1993).
- [12] V.A. Fateev, Integrable deformations in Z_N -symmetrical models of the conformal quantum field theory, Int. J. Mod. Phys. A 6 (1991), 2109–2132
- [13] A.S. Fokas and B. Fuchssteiner, Symplectic structures, their Bäcklund transformations and hereditary symmetries, Physica D 4 (1981/82), 47–66
- [14] M.V. Foursov, On integrable evolution equations in commutative and noncommutative variables, Ph.D. thesis (University of Minnesota, Minneapolis, 1999). Available online at http://www.irisa.fr/prive/foursov/articles/thesis.ps
- [15] M.V. Foursov, Classification of certain integrable coupled potential KdV and modified KdV-type equations, J. Math. Phys. 41 (2000), 6173-6185. Available online at http://www.irisa.fr/prive/foursov/articles/articlec.ps
- B.S. Getmanov, New Lorentz invariant systems with exact multi-soliton solutions, JETP Lett.
 25 (1977), 119–122

- [17] B.S. Getmanov, An integrable two-dimensional Lorentz-invariant nonlinear model of a complex scalar field (complex sine-Gordon II), Teor. Mat. Fiz. 48 (1981), 13–23
- [18] C. Gilson, J. Hietarinta, J. Nimmo, Y. Ohta, Sasa-Satsuma higher-order nonlinear Schrödinger equation and its bilinearization and multisoliton solutions. Phys. Rev. E 68 (2003) 016614 (*Preprint* nlin.SI/0304025)
- [19] R. Hirota, Exact envelope-soliton solutions of a nonlinear wave equation, J. Math. Phys. 14 (1973), 805–809.
- [20] R. Hirota, Bilinearization of soliton equations, J. Phys. Soc. Jap. 51 (1982), 323–331
- [21] N.H. Ibragimov, A.V. Aksenov, V.A. Baikov, V.A. Chugunov, R.K. Gazizov, A.G. Meshkov, *CRC handbook of Lie group analysis of differential equations*, vol.2, edited by N.H. Ibragimov, (CRC Press, Boca Raton, 1995).
- [22] Ayşe Karasu, Atalay Karasu, and S. Yu. Sakovich, A strange recursion operator for a new integrable system of coupled Korteweg-de Vries equations. Acta Appl. Math. 83 (2004), 85–94 (*Preprint* nlin.SI/0203036)
- [23] P. Kersten and J. Krasil'shchik "Complete integrability of the coupled KdV-mKdV system," in Lie groups, geometric structures and differential equations—one hundred years after Sophus Lie (Kyoto/Nara, 1999) (Math. Soc. Japan, Tokyo, 2002), pp.151–171 (preprint nlin.SI/0010041).
- [24] Y. Kivshar and G. Agrawal, Optical Solitons: From Fibers to Photonic Crystals (Academic Press, New York, 2003).
- [25] Y. Kodama, Optical solitons in a monomode fiber, J. Stat. Phys. **39** (1985), 597–614; Y. Kodama and A. Hasegawa, Nonlinear pulse propagation in a monomode dielectric guide, IEEE J. Quantum Electronics **QE-23** (1987), 510–524
- [26] F. Lund and T. Regge, Unified approach to strings and vortices with soliton solutions, Phys. Rev. D 14 (1976), 1524–1535
- [27] F. Magri, "A geometrical approach to nonlinear solvable equations," in Nonlinear evolution equations and dynamical systems (Proc. Meeting, Univ. Lecce, Lecce, 1979), Lect. Not. Phys. 120 (Springer, Berlin–New York, 1980), pp.233–263.

- [28] A.G. Meshkov, Hamiltonian and recursion operators for two-dimensional scalar fields Phys. Lett. A 170 (1992), 405–408
- [29] O.I. Mokhov, Symplectic forms on a loop space and Riemannian geometry Funct. Anal. Appl. 24 (1990), 247–249
- [30] A. Neveu and N. Papanicolaou, Integrability of the classical $[\bar{\psi}_i\psi_i]^2$ and $[\bar{\psi}_i\psi_i]^2 [\bar{\psi}_i\gamma_5\psi_i]^2$ interactions, Commun. Math. Phys. **58** (1978), 31–64
- [31] N. Olver and I.V. Barashenkov, Complex sine-Gordon II: a new algorithm for multivortex solutions on the plane, Theor. and Math. Phys. 144 (2005), 1223–1226 (*Preprint nlin.SI/0502048*)
- [32] P.J. Olver, Applications of Lie Groups to Differential Equations (Springer, New York, 1993).
- [33] K. Pohlmeyer, Integrable Hamiltonian systems and interactions through quadratic constraints Commun. Math. Phys. 46 (1976), 207–221
- [34] K. Porsezian, Soliton models in resonant and nonresonant optical fibers, Pramana 57 (2001) 1003–1039.
- [35] C. Rogers and W.K. Schief, Bäcklund and Darboux transformations (Cambridge University Press, Cambridge, 2002).
- [36] N. Sasa and J. Satsuma, New type of soliton solutions for a higher-order nonlinear Schrödinger equation, J. Phys. Soc. Japan 60 (1991), 409–417
- [37] W. Schief, Hidden integrability in ideal magnetohydrodynamics: The Pohlmeyer–Lund–Regge model, Physics of Plasmas 10 (2003), 2677–2685
- [38] S. Sciuto, Exterior calculus and two-dimensional supersymmetric models, Phys. Lett. B 90 (1980), 75–80
- [39] A. Sergyeyev, "On recursion operators and nonlocal symmetries of evolution equations," in Proc. Sem. Diff. Geom. edited by D. Krupka (Silesian University in Opava, Opava, 2000) pp.159–173 (Preprint nlin.SI/0012011)
- [40] A. Sergyeyev, A strange recursion operator demystified, J. Phys. A: Math. Gen. 38 (2005)
 L257–L262 (*Preprint* nlin.SI/0406032)

- [41] A. Sergyeyev, Why nonlocal recursion operators produce local symmetries: new results and applications, J. Phys. A: Math. Gen. 38 (2005), 3397–3407 (*Preprint nlin.SI/0410049*)
- [42] V.V. Sokolov and S.I. Svinolupov, Vector-matrix generalizations of classical integrable equations, Theor. Math. Phys. 100, 959–962 (1994).
- [43] V.V. Sokolov and T. Wolf, Classification of integrable polynomial vector evolution equations J.
 Phys. A: Math. Gen. 34 (2001), 11139–11148 (*Preprint nlin.SI/0611038*)
- [44] R.S. Tasgal and M.J. Potasek, Soliton solutions to coupled higher order nonlinear Schrödinger equations, J. Math. Phys. 33 (1992), 1208–1215
- [45] T. Tsuchida and T. Wolf, Classification of polynomial integrable systems of mixed scalar and vector evolution equations: I, J. Phys. A: Math. Gen. 38, 7691–7733 (2005) (Preprint nlin.SI/0412003)
- [46] N. Yajima and M. Oikawa, A class of exactly solvable nonlinear evolution equations, Progr. Theor. Phys. 54, 1576–1577 (1975).
- [47] A.V. Zhiber, V.V. Sokolov, Exactly integrable hyperbolic equations of Liouville type. Russian Math. Surveys 56 (2001), no.1, 61–101