Transformations of integrable hydrodynamic chains and their hydrodynamic reductions

Maxim V. Pavlov

Abstract

Hydrodynamic reductions of the hydrodynamic chain associated with dispersionless limit of 2+1 Harry Dym equation are found by the Miura type and reciprocal transformations applied to the Benney hydrodynamic chain.

Contents

1	Introduction	1
2	Modified Benney hydrodynamic chain	4
3	Dispersionless 2+1 Harry Dym equation	5
R	References	

keywords: hydrodynamic chain, reciprocal transformation, Miura type transformation, hydrodynamic reduction.

MSC: 35L40, 35L65, 37K10; PACS: 02.30.J, 11.10.E.

1 Introduction

Solutions of integrable hydrodynamic chains and 2+1 quasilinear systems can be found by the method of hydrodynamic reductions (see, for instance, [6], [8]). Suppose some hydrodynamic chain is given. Then one can seek hydrodynamic reductions for this hydrodynamic chain by the aforementioned method of hydrodynamic reductions or by $\bar{\partial}$ approach (see [2]), for instance. Thus, hydrodynamic reductions of a dispersionless limit of 2+1 Harry Dym equation can be found directly by the Hamiltonian approach applied for the Kupershmidt hydrodynamic chain (see [14] and [15]) or by $\bar{\partial}$ approach (see [18]). However, in this paper we concentrate on re-calculation of hydrodynamic reductions of hydrodynamic chains related by Miura type and reciprocal transformations. It means that we use already known hydrodynamic reductions of the Benney hydrodynamic chain (see [2], [11], [19]) for reconstruction of hydrodynamic reductions for a dispersionless limit of 2+1 Harry Dym equation. Thus, this paper is devoted to the construction of hydrodynamic reductions common for 2+1 quasilinear systems (dKP, dmKP, 2+1 Harry Dym) related by the Miura type and reciprocal transformations (see [3], [4], [5], [16], [17]).

Let us recall (see [7]) that the Benney hydrodynamic chain (see [1])

$$A_t^k = A_x^{k+1} + kA^{k-1}A_x^0, \qquad k = 0, 1, 2, \dots$$
(1)

satisfies the Gibbons equation

$$\lambda_t - \mu \lambda_x = \frac{\partial \lambda}{\partial \mu} \left[\mu_t - \partial_x \left(\frac{\mu^2}{2} + A^0 \right) \right], \qquad (2)$$

where the equation of the Riemann mapping is given by the asymptotic series

$$\lambda = \mu + \frac{A^0}{\mu} + \frac{A^1}{\mu^2} + \frac{A^2}{\mu^3} + \dots$$
(3)

The inverse asymptotic series

$$\mu = \lambda - \frac{\mathbf{H}_0}{\lambda} - \frac{\mathbf{H}_1}{\lambda^2} - \frac{\mathbf{H}_2}{\lambda^3} - \dots$$

yields an infinite series of polynomial conservation law densities $\mathbf{H}_k(A^0, A^1, ..., A^k)$, where the generating function of conservation laws is given by

$$\mu_t = \partial_x \left(\frac{\mu^2}{2} + A^0\right). \tag{4}$$

Since the transformation $\mathbf{H}_k = \mathbf{H}_k(A^0, A^1, ..., A^k)$ is invertible, the Benney hydrodynamic chain can be written in the conservative form

$$\partial_t \mathbf{H}_0 = \partial_x \mathbf{H}_1, \qquad \partial_t \mathbf{H}_n = \partial_x \left(\mathbf{H}_{n+1} - \frac{1}{2} \sum_{k=0}^{n-1} \mathbf{H}_k \mathbf{H}_{n-1-k} \right), \qquad n = 1, 2, \dots$$
(5)

Lets us apply the generating function of the Miura type transformations (see [12])

$$\mu = p + B^0$$

to the above series (see also [4], [17]). A deformation of the Riemann mapping determined by the asymptotic series

$$\lambda = p + B^0 + \frac{B^1}{p} + \frac{B^2}{p^2} + \frac{B^3}{p^3} + \dots$$
(6)

satisfies the modified Gibbons equation

$$\lambda_t - (p + B^0)\lambda_x = \frac{\partial\lambda}{\partial p} \left[p_t - \partial_x \left(\frac{p^2}{2} + B^0 p \right) \right],\tag{7}$$

where a dynamics of the coefficients B^k is given by the modified Benney hydrodynamic chain

$$B_t^k = B_x^{k+1} + B^0 B_x^k + k B^k B_x^0, \qquad k = 0, 1, 2, \dots$$
(8)

Remark: The comparison of two asymptotic series (3) and (6)

$$p + B^{0} + \frac{B^{1}}{p} + \frac{B^{2}}{p^{2}} + \frac{B^{3}}{p^{3}} + \dots = p + B^{0} + \frac{A^{0}}{p + B^{0}} + \frac{A^{1}}{(p + B^{0})^{2}} + \frac{A^{2}}{(p + B^{0})^{3}} + \dots$$

yields explicit polynomial Miura type transformations $A^k(B^0, B^1, ..., B^{k+1}), k = 0, 1, 2, ...$

The relationship between 2+1 Harry Dym equation and the KP equation was established in [16] (see also [3]). Let us recall this link between corresponding hydrodynamic chains (see [5]). The hydrodynamic chain connected with a dispersionless limit of 2+1 Harry Dym equation is (see, for instance, [3])

$$C_y^k = (C^{-1})^2 C_z^{k+1} + (k+1)C^{k+1}C^{-1}C_z^{-1}, \qquad k = -1, 0, 1, 2, \dots$$
(9)

A deformation of the Riemann mapping

$$\lambda = C^{-1}q + C^0 + \frac{C^1}{q} + \frac{C^2}{q^2} + \frac{C^3}{q^3} + \dots$$
(10)

is given by the Gibbons equation

$$\lambda_y - q(C^{-1})^2 \lambda_z = \frac{\partial \lambda}{\partial q} \left[q_y - \partial_z \left(\frac{q^2 (C^{-1})^2}{2} \right) \right].$$
(11)

This hydrodynamic chain has the first conservation law

$$\partial_y \frac{1}{C^{-1}} = \partial_z (-C^0).$$

Under the reciprocal transformation

$$dx = \frac{1}{C^{-1}}dz - C^0 dy, \qquad dt = dy$$
 (12)

the hydrodynamic chain (9) reduces to the hydrodynamic chain (8); the Gibbons equation (11) reduces to the Gibbons equation (7); the equation of the Riemann mapping (10) reduces to the equation of the Riemann mapping (6), where the generating function of the Miura type transformations is

$$p = C^{-1}q$$

and the Miura type transformations are (see [5])

$$B^k = C^k (C^{-1})^k, \quad k = 0, 1, 2, \dots$$

The Benney hydrodynamic chain is well known. Plenty hydrodynamic reductions (see for instance, [2], [11], [19]) are found many years ago. Below we show links with hydrodynamic reductions of hydrodynamic chains connected by the Miura type and reciprocal transformations with the Benney hydrodynamic chain.

2 Modified Benney hydrodynamic chain

For instance, the Zakharov hydrodynamic reductions (see [19])

$$a_t^i = \partial_x \left[\frac{(a^i)^2}{2} + A^0 \right], \qquad b_t^i = \partial_x (a^i b^i), \qquad A^0 = \sum_{n=1}^N b^n$$
 (13)

of the Benney moment chain (1) are connected with the equation of the Riemann surface

$$\lambda = \mu + \sum_{n=1}^{N} \frac{b^n}{\mu - a^n}.$$

Let us rewrite the above formula in the form

$$\lambda = \mu + \frac{b^i}{\mu - a^i} + \sum_{k \neq i} \frac{b^k}{\mu - a^k},\tag{14}$$

where the index i is **fixed**. Substituting the Taylor series

$$\mu^{(i)} = a^i + \frac{b^i}{\lambda} + \frac{c^i(\mathbf{a}, \mathbf{b})}{\lambda^2} + \frac{d^i(\mathbf{a}, \mathbf{b})}{\lambda^3} + \dots$$

in (4), one can obtain an infinite series of conservation laws at the vicinity of each puncture $\mu^{(i)} = a^i$. The Benney hydrodynamic chain (5) is determined for $k \ge 0$. Let us extend this hydrodynamic chain in the *negative* direction. Then the negative part of the Benney hydrodynamic chain is given by

$$\partial_t \mathbf{H}_{-1} = \partial_x \left[\frac{(\mathbf{H}_{-1})^2}{2} + \mathbf{H}_0 \right], \qquad \partial_t \mathbf{H}_{-k-1} = \frac{1}{2} \partial_x \left[\sum_{m=0}^k \mathbf{H}_{-m-1} \mathbf{H}_{m-k-1} \right], \qquad k = 1, 2, ...,$$

where $\mathbf{H}_{-1} \equiv a^i$, $\mathbf{H}_{-2} \equiv b^i$, and the above hydrodynamic type system

$$a_{t}^{k} = \partial_{x} \left(\frac{(a^{k})^{2}}{2} + \mathbf{H}_{-2} + \sum_{n \neq i} b^{n} \right), \qquad b_{t}^{k} = \partial_{x} (a^{k} b^{k}), \qquad k = 1, 2, \dots, N, \qquad k \neq i,$$

$$\partial_{t} \mathbf{H}_{-1} = \partial_{x} \left(\frac{(\mathbf{H}_{-1})^{2}}{2} + \mathbf{H}_{-2} + \sum_{n \neq i} b^{n} \right), \qquad \partial_{t} \mathbf{H}_{-2} = \partial_{x} (\mathbf{H}_{-1} \mathbf{H}_{-2})$$

is connected with the equation of the Riemann surface (see (14))

$$\tilde{\lambda} \equiv \lambda^{(i)} = \frac{\mathbf{H}_{-2}}{\mu - \mathbf{H}_{-1}} + \mu + \sum_{k \neq i} \frac{b^k}{\mu - a^k}.$$
(15)

Since the first moment of the modified Benney chain $B^0 \equiv \mathbf{H}_{-1}$, the equation of the Riemann surface for the corresponding hydrodynamic reduction

$$u_t^k = \partial_x \left(\frac{(u^k)^2}{2} + B^0 u^k \right), \qquad b_t^k = \partial_x [(B^0 + u^k) b^k], \qquad k = 1, 2, \dots, N, \qquad k \neq i,$$

$$B_t^0 = \partial_x \left(\frac{(B^0)^2}{2} + \mathbf{H}_{-2} + \sum_{n \neq i} b^n \right), \qquad \partial_t \mathbf{H}_{-2} = \partial_x (B^0 \mathbf{H}_{-2})$$

of the modified Benney hydrodynamic chain is given by

$$\lambda = \frac{\mathbf{H}_{-2}}{p} + B^0 + p + \sum_{k \neq i} \frac{b^k}{p - u^k},$$

where $u^k = a^k - B^0$; the generating function of the Miura type transformations is $p = \mu - B^0$.

3 Dispersionless 2+1 Harry Dym equation

Finally, let us apply the reciprocal transformation (see (12))

$$dz = \mathbf{H}_{-2}dx + B^0\mathbf{H}_{-2}dt, \qquad dy = dt \tag{16}$$

to the above hydrodynamic type system.

Then the equation of the Riemann surface for the corresponding hydrodynamic reduction

$$\bar{u}_y^k = \partial_z \left[\frac{(\bar{u}^k)^2}{2} (C^{-1})^2 \right], \qquad \bar{b}_y^k = \partial_z \left(C^{-1} \bar{u}^k \bar{b}^k \right), \qquad k = 1, 2, \dots, N, \qquad k \neq i,$$

$$C_y^0 = \left(1 + \sum_{n \neq i} \bar{b}^n \right) C^{-1} C_z^{-1} + (C^{-1})^2 \sum_{n \neq i} \bar{b}_z^n, \qquad C_y^{-1} = (C^{-1})^2 C_z^0$$

of a dispersionless limit of 2+1 Harry Dym equation is given by

$$\lambda = C^{-1}q + C^0 + \frac{1}{q} + \sum_{k \neq i} \frac{\bar{b}^k}{q - \bar{u}^k},$$

where $C^0 \equiv B^0$, $C^{-1} \equiv \mathbf{H}_{-2}$, $\bar{b}^k = b^k / \mathbf{H}_{-2}$, $\bar{u}^k = u^k / \mathbf{H}_{-2}$; and the generating function of the Miura type transformations is $p = qC^{-1}$.

Let us rewrite the equation of the Riemann surface (14) for the Zakharov hydrodynamic reduction (13) in the form (see [13])

$$\lambda = p \prod_{k=1}^{N} \frac{p - c^k}{p - u^k},$$

where $B^0 \equiv \mathbf{H}_{-1} = \sum (u^n - c^n)$. Application of the reciprocal transformation (16) yields *another* hydrodynamic reduction of a dispersionless limit of 2+1 Harry Dym equation

$$\bar{u}_y^k = \partial_x \left[\frac{(\bar{u}^k)^2}{2} (C^{-1})^2 \right], \qquad \bar{c}_y^k = \partial_x \left[\frac{(\bar{c}^k)^2}{2} (C^{-1})^2 \right], \qquad k = 1, \, 2, \, \dots, \, N,$$

where $C^{-1} \equiv \mathbf{H}_{-2} = \prod (u^n/c^n) \equiv \prod (\bar{u}^n/\bar{c}^n)$. The corresponding equation of the Riemann surface is given by

$$\lambda = q \prod_{k=1}^{N} \frac{1 - q/\bar{c}^k}{1 - q/\bar{u}^k}$$

Let us start from the waterbag reduction (see [9] and [10])

$$a_t^k = \partial_x \left[\frac{(a^k)^2}{2} + \sum_{n=1}^N \varepsilon_n a^n \right], \qquad k = 1, 2, ..., N$$

of the Benney moment chain (1)

$$\lambda = \mu - \varepsilon_i \ln(\mu - a^i) - \sum_{k \neq i} \varepsilon_k \ln(\mu - a^k), \qquad \sum \varepsilon_n = 0.$$

Let us identify $\mathbf{H}_{-1} \equiv a^i$, where the index *i* is **fixed**. Then the above hydrodynamic type system

$$a_t^k = \partial_x \left[\frac{(a^k)^2}{2} + \varepsilon_i \mathbf{H}_{-1} + \sum_{n \neq i} \varepsilon_n a^n \right], \qquad k = 1, 2, \dots, N, \qquad k \neq i,$$
$$\partial_t \mathbf{H}_{-1} = \partial_x \left[\frac{(A^{-1})^2}{2} + \varepsilon_i \mathbf{H}_{-1} + \sum_{n \neq i} \varepsilon_n a^n \right]$$

is connected with the equation of the Riemann surface (cf. (15))

$$\tilde{\lambda} \equiv \lambda^{(i)} = \frac{1}{\mu - \mathbf{H}_{-1}} \prod_{k \neq i} (\mu - a^k)^{-\varepsilon_k/\varepsilon_i} \exp \frac{\mu}{\varepsilon_i}.$$

Since the first moment of the modified Benney chain $B^0 \equiv \mathbf{H}_{-1}$, the equation of the Riemann surface for the corresponding hydrodynamic reduction

$$u_t^k = \partial_x \left(\frac{(u^k)^2}{2} + B^0 u^k \right), \qquad k = 1, 2, \dots, N, \qquad k \neq i,$$
$$B_t^0 = \partial_x \left(\frac{(B^0)^2}{2} + \sum_{n \neq i} \varepsilon_n u^n \right),$$

of the modified Benney chain is given by

$$\lambda = \frac{\mathbf{H}_{-2}}{p} e^{p/\varepsilon_i} \prod_{k \neq i} \left(1 - \frac{p}{u^k} \right)^{-\varepsilon_k/\varepsilon_i}.$$

Finally, let us apply the reciprocal transformation (16), where

$$\mathbf{H}_{-2} = e^{B^0/\varepsilon_i} \prod_{k \neq i} (u^k)^{-\varepsilon_k/\varepsilon_i}.$$

Then the equation of the Riemann surface for the corresponding hydrodynamic reduction

$$\bar{u}_y^k = \partial_z \left[\frac{(\bar{u}^k)^2}{2} (C^{-1})^2 \right], \qquad k = 1, 2, \dots, N, \qquad k \neq i,$$
$$C_y^{-1} = (C^{-1})^2 \partial_z \left(\sum_{n \neq i} \varepsilon_n \ln \bar{u}^n \right),$$

of the dispersionless limit of 2+1 Harry Dym equation is given by

$$\lambda = C^{-1}q - \sum_{n \neq i} \varepsilon_n \ln \frac{1 - \bar{u}^n/q}{\bar{u}^n}.$$

All other hydrodynamic reductions of the Benney moment chain can be re-calculated into hydrodynamic reductions of (9) in the same way.

Acknowledgement

I am grateful to the Institute of Mathematics in Taipei (Taiwan) where some part of this work has been done, and especially to Jen-Hsu Chang and Derchyi Wu for fruitful and stimulating discussions.

References

- [1] D.J. Benney, Some properties of long non-linear waves, Stud. Appl. Math., **52** (1973) 45-50.
- [2] L.V. Bogdanov, B.G. Konopelchenko, Symmetry constraints for dispersionless integrable equations and systems of hydrodynamic type, Phys. Lett. A, **330** (2004) 448–459.
- [3] Jen-Hsu Chang, Hydrodynamic Reductions of Dispersionless Harry Dym Hierarchy, J. Phys. A, Math. Gen., 38 No. 29 (2005) 6505-6515.
- [4] Jen-Hsu Chang, Ming-Hsien Tu, On the Miura Map between the dispersionless KP and dispersionless modified KP hierarchies, J. Math. Phys. 41 No. 8 (2000) 5391-5406.
- [5] Yu-Tung Chen, Ming-Hsien Tu, A Note on the Dispersionless Dym Hierarchy, Letters in Mathematical Physics, **63** (2003) 125–139.
- [6] E.V. Ferapontov, K.R. Khusnutdinova, On integrability of (2+1)-dimensional quasilinear systems, Comm. Math. Phys. 248 (2004) 187-206, E.V. Ferapontov, K.R. Khusnutdinova, The characterization of 2-component (2+1)-dimensional integrable systems of hydrodynamic type, J. Phys. A: Math. Gen. 37 No. 8 (2004) 2949 - 2963.
- [7] J. Gibbons, Collisionless Boltzmann equations and integrable moment equations, Physica D, 3 (1981) 503-511.
- [8] J. Gibbons, S.P. Tsarev, Reductions of Benney's equations, Phys. Lett. A, 211, (1996) 19-24. J. Gibbons, S.P. Tsarev, Conformal maps and reductions of the Benney equations, Phys. Lett. A, 258 (1999) 263-270.
- [9] J. Gibbons, L.A. Yu, The initial value problem for reductions of the Benney equations, Inverse Problems 16 No. 3 (2000) 605-618, L.A. Yu, Waterbag reductions of the dispersionless discrete KP hierarchy, J. Phys. A: Math. Gen., 33 (2000) 8127–8138.

- [10] Yu. Kodama, A method for solving the dispersionless KP equation and its exact solutions. Phys. Lett. A, **129** No. 4 (1988) 223-226. Yu. Kodama, A solution method for the dispersionless KP equation, Prog. Theor. Phys. Supplement. **94** (1988) 184.
- [11] I.M. Krichever, The averaging method for two-dimensional "integrable" equations, Funct. Anal. Appl. **22** No. 3 (1988) 200-213, *I.M. Krichever*, Spectral theory of two-dimensional periodic operators and its applications, Russian Math. Surveys **44** No. 2 (1989) 145-225. *I.M. Krichever*, The dispersionless equations and topological minimal models, Comm. Math. Phys., **143** No. 2 (1992) 415-429. *I.M. Krichever*, The τ -function of the universal Whitham hierarchy, matrix models and topological field theories, Comm. Pure Appl. Math. **47** (1994) 437-475.
- [12] B.A. Kupershmidt, Deformations of integrable systems, Proc. Roy. Irish Acad. Sect. A 83 No. 1 (1983) 45-74. B.A. Kupershmidt, Normal and universal forms in integrable hydrodynamical systems, Proceedings of the Berkeley-Ames conference on nonlinear problems in control and fluid dynamics (Berkeley, Calif., 1983), in Lie Groups: Hist., Frontiers and Appl. Ser. B: Systems Inform. Control, II, Math Sci Press, Brookline, MA, (1984) 357-378.
- M. V. Pavlov, Local Hamiltonian structures of Benney's system. Russian Phys. Dokl., 39 No. 9 (1994) 607-608. M. V. Pavlov, Exact integrability of a system of the Benney equations. Russian Phys. Dokl., 39 No. 11 (1994) 745-747.
- [14] M.V. Pavlov, The Kupershmidt hydrodynamic chains and lattices.
- [15] *M.V. Pavlov*, The Hamiltonian approach in the classification and the integrability of hydrodynamic chains.
- [16] C. Rogers, The Harry Dym equation in 2+1 dimensions: a reciprocal link with the Kadomtsev–Petviashvili equation, Phys. Lett. A, **120** No. 1 (1987) 15-18. W. Oevel, C. Rogers, Gauge Transformations and Reciprocal Links in 2+1 Dimensions, Reviews in Mathematical Physics, **5** (1993) 299-330.
- [17] Jiin-Chang Shaw, Ming-Hsien Tu, Canonical Miura maps between the modified KP and KP hierarchies, J. Phys. A, Math. Gen., 30 (1997) 4825-4833.
- [18] Derchyi Wu, Ming-Hsien Tu, Yu-Tung Chen, Jen-Hsu Chang, The ∂ approach to the dispersionless (2+1)-Harry Dym hierarchy, J. Phys. A, Math. Gen., 38 (2005) 6167-6181.
- [19] V.E. Zakharov, Benney's equations and quasi-classical approximation in the inverse problem method, Funct. Anal. Appl., 14 No. 2 (1980) 89-98. V.E. Zakharov, On the Benney's Equations, Physica 3D (1981) 193-200.