

# 3D-vortex labyrinths in the near field of solid-state microchip laser.

A.Yu.Okulov\*

*P.N.Lebedev Physical Institute of Russian Academy of Sciences  
Leninsky prospect 53, 119991 Moscow, Russia*

(Dated: January 15, 2007)

The usage of vortex labyrinths fields and Talbot lattices as optical dipole traps for neutral atoms is considered for the wavelength of trapping radiation in the range  $0.98 \div 2.79 \mu m$ . The square vortex lattices generated in high Fresnel number solid-state microchip lasers are studied as a possible realization. The distribution of light field is obtained via dynamical model based on Maxwell-Bloch equations for class-B laser, discrete Fox-Lee map with relaxation of inversion and static model based on superposition of copropagating Gaussian beams. The spatial patterns observed experimentally and obtained numerically are interpreted as nonlinear superposition of vortices with helicoidal phase dislocations. Such patterns are approximated analytically by a sum of array of vortex lines and additional parabolic subtrap. The separable optical trapping potential is proposed. The factorization of macroscopic wavefunction have led to analytical solution of Gross-Pitaevski equation for ensemble of quantum particles trapped in vortex labyrinth formed by spatially - periodic array of Laguerre-Gaussian beams.

PACS numbers: 03.75.Lm., 42.50.Vk., 42.60.Jf, 42.65.Sf., 47.32.Cc

## I. INTRODUCTION

Optical dipole traps for neutral atoms [1] are a subject of considerable interest. A certain amount of proposals of increasing complexity of trapping EM-field appeared recently. Apart from a relatively simple geometrical patterns like standing plane-wave gratings[2], evanescent wave mirrors[3], toroidal traps which utilize the intensity distribution of Laguerre-Gaussian (GL) beams in beam waist [4] a several proposals were made which utilize the arrays of Gaussian beams, both phase-locked [5, 6] and unlocked ones [7]. The interference inherent to phase-locking provides multiply connected configurations of intensity distribution, phase gradients and electromagnetic (EM) momentum density [8, 9]. The winding EM momentum distribution is the cause of the angular momentum transfer to macroscopic bodies (e.g. dielectric ball) [10] and to trapped BEC as well [11].

The EM field configuration under consideration is based upon properties of self-imaging optical fields[12, 13]. The difference between phase-locked array of zero-order Gaussian beams and phase-locked array of optical vortices (OV) obtained experimentally in the near field of solid-state microchip laser [14] is that the latter consists of array of parallel vortex lines with opposite circulations and topological charges  $\ell_{EM}$  (TC) (fig. 1) [5]. In contrast to earlier proposals where individual loading and addressing of trapping sites was considered [7] this OV array have sophisticated configuration of intensity [5] and EM momentum density of trapping field. Due to this configuration the BEC trapped in EM field might have macroscopic wavefunction of complex form composed of array of superfluid vortices (SFV). The cause of SFV formation is light-induced torque experienced by isolated resonant atom which interacts with Laguerre-Gaussian beam having phase singularity. It was show by Allen et al.[8] the value of torque in saturation limit is  $T = \hbar \ell_{EM} \Gamma$ . The origin of torque is due to nonzero azimuthal EM momentum component. The azimuthal Doppler shift corresponding to such motion had been onerved[9]. This torque might have an appreciable value even in nonresonant case, although it is significantly reduced by the multiple of  $(\Delta/\Gamma)^2$ , where  $\Delta$  - is detuning,  $\Gamma$  - is linewidth[8]. As a result an azimuthal component of EM momentum is transferred to atom in such a way that it will move around phase singularity - the direction of rotation is fixed by TC of trapping beam. This is the cause of circular motion of BEC confined in isolated toroidal trap.

In current paper the maximally transparent and simple analytical description presented which describes the transfer of angular momentum from optical vortex array (OVA) to trapped BEC. As a first step the distribution of light intensity and phase in the near field of solid-state microchip laser is computed. The configuration of intensity and phase of EM-field of OVA has complicated, multinode structure. In order to get analytic solution for macroscopic wavefunction  $\Psi$  trapped by such array, the special optical pancake-like potential (fig. 1) will be constructed. In this trapping potential the separation of variables in Gross-Pitaevsky equation[2] becomes possible. Next the procedure

---

\*Electronic address: okulov@sci.lebedev.ru; URL: <http://sites.lebedev.ru/okulov>

applied previously to elongated  $sech^2$ - profile optical trap [15] will be used. Due to special adjustment of potential it is possible to capture an analytic approximation for  $\Psi$  in the form of superposition of elementary equispaced vortices. The numerical modeling via split-step FFT technique supports this approximate formula: the small but unavoidable EM-torque existing in OV even at large detuning from resonance  $\Delta$  imposes the correlation between TC of BEC vortices and TC of the optical vortices.

The BEC angular momentum distribution is formed via periodic lattice of SFV with alternating topological charges. There is a remarkable contrast with superfluid in "rotating bucket" trap: while angular momenta of SFV in bucket are co-directed [16, 17] the BEC vortices trapped by OVA are counter-directed from site to site.

As a result in the net sum each angular momentum of SFV of positive topological charge is compensated by the angular momentum of SFV with negative charge and total angular momentum tends to zero. Meantime the mutual subtraction of angular momenta vectors of adjacent vortices in the net sum does not mean the mutual subtraction of rotational energies, which are the positive scalars. The macroscopic quantum state carries rotational kinetic energy of condensate containing  $N$  particles of mass  $m$  per unit volume, namely  $E_{rot} = N_{vortices} N \pi \hbar^2 \ln(b/a) \chi / m$ ,  $b$ -diameter of vortex core,  $a$ -interatomic distance,  $\chi$  - length of vortex [17]. This feature of macroscopic wavefunction  $\Psi$  trapped by vortex array might be useful from the point view of diminishing the decoherence induced by environment in topological quantum computing[18].

## II. SQUARE VORTEX LATTICES

Recent advances in controlling the dynamics of solid-state microchip lasers [14] offer the possibility of reliable control of spatiotemporal optical patterns. Compared to semiconductor lasers the host medium composed of dielectric crystal doped by neodymium  $Nd^{+3}$  or other rare earth ions ( $Er^{+3}$ ,  $Tm^{+3}$ ,  $Ho^{+3}$ ,  $Yb^{+3}$ ) have smaller gain and smaller self-phase modulation. Because of smaller nonresonant losses the heating of host crystal by radiation is not dramatic. The changes of the geometry and birefringence of host crystals, curvature and reflectivity of output couplers and spatial distribution of optical pumping give efficient control over mode structure. The square vortex lattices (SVL) observed in quasi plane parallel cavity[14] are of special interest. These lattices demonstrate high degree of spatial coherence : the relaxation oscillations of class-B high Fresnel number solid-state microchip laser [14] with Fresnel number in the range  $N_f \approx 100 \div 1000$  are characterized by single peak at frequency about  $(\sqrt{T_1 \tau_c})^{-1}$  (  $0.3 \div 1.2 MHz$  ). This is the firm evidence of single-longitudinal and single transverse mode behaviour. The former theoretical analysis was based on parabolic equation, resulted from adiabatic elimination of polarization from standard set of Maxwell-Bloch equations [19]:

$$\frac{\partial E(\vec{r}, t)}{\partial t} + \frac{E(\vec{r}, t)}{\tau_c} + \frac{i c}{2k} \Delta_{\perp} E(\vec{r}, t) = \frac{\sigma c N_0 L_a E(\vec{r}, t) (1 + i \delta \omega T_2)}{2 L_r (1 + \sigma T_1 c \epsilon_0 |E|^2 / \hbar \omega)} \quad (1)$$

where  $\tau_c$  - is photon lifetime in cavity,  $k=2\pi/\lambda$  - wavenumber,  $\sigma$  - stimulated emission crosssection,  $\delta \omega$  - detuning of lasing frequency from center of gain line,  $N_0$  - density of inverted resonant atoms per unit volume,  $T_1$  - inversion lifetime(longitudinal relaxation lifetime),  $L_a$  - thickness of active medium,  $L_r$  - length of resonator,  $c$  - speed of light,  $\epsilon_0$  - dielectric constant,  $\Delta_{\perp} = \nabla_{\perp}^2$  [20]. When finite gain linewidth  $T_2^{-1}$  is taken into account in the framework of Swift-Hohenberg equation the square vortex lattices were obtained numerically[21]:

$$\begin{aligned} & \frac{\partial E(\vec{r}, t)}{\partial t} + \frac{E(\vec{r}, t)}{(\tau_c + T_2)} + \frac{i \tau_c c}{2k(\tau_c + T_2)} \Delta_{\perp} E(\vec{r}, t) - \\ & \frac{T_2^2}{\tau_c(\tau_c + T_2)^2} \left( \frac{\tau_c c}{k} \Delta_{\perp} + \delta \omega T_2 \right)^2 E(\vec{r}, t) = \frac{\sigma c N_0 L_a E(\vec{r}, t) (1 + i \delta \omega T_2)}{2 L_r (1 + \sigma T_1 c \epsilon_0 |E|^2 / \hbar \omega)} \end{aligned} \quad (2)$$

The additional term with the square of transverse laplacian is responsible for transverse mode selection due to finite gain linewidth. An alternative model with discrete time step equal to  $\tau = 2L_r/c$  (time of bouncing of radiation between mirrors) utilizes the standard rate equations of class-B laser written at  $n$ -th step for electric field[5, 22]:

$$E_{n+1}(\vec{r}) = f(E_n(\vec{r})) = \frac{\sigma L_a N_n(\vec{r}) E_n(\vec{r}) (1 + i \delta \omega T_2)}{2} + E_n(\vec{r}), \quad (3)$$

inversion:

$$N_{n+1}(\vec{r}) = N_n(\vec{r}) + \left[ \frac{N_0(\vec{r}) - N_n(\vec{r})}{T_1} - \sigma N_n(\vec{r}) c \epsilon_0 |E|^2 / \hbar \omega \right] \frac{2 L_r}{c}, \quad (4)$$

and nonlocal integral mapping evaluating the field via fast Fourier transform at each timestep:

$$E_{n+1}(\vec{r}) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} K(\vec{r} - \vec{r}') f(E_n(\vec{r}')) d^2 r', \quad (5)$$

The kernel  $K$  for the nearly plane-parallel Fabry-Perot cavity of microchip laser with transverse filtering via aperture  $D(\vec{r}')$  has the form [12] :

$$K(\vec{r} - \vec{r}') = \frac{ikD(\vec{r}')}{2\pi L_r} \exp [ik(\vec{r} - \vec{r}')^2 / 2L_r]. \quad (6)$$

The similar spatiotemporal picture of transverse modes was obtained for 2D transversal field distribution. The following parameters were chosen in numerical simulation:  $T_1 = 2 \cdot 10^{-4} \text{ sec}$ ,  $L_r = 1 \text{ mm}$ , for  $Nd^{+3}$ -doped crystals  $\sigma = (1.2 - 0.6) \cdot 10^{-20} \text{ cm}^2$ ,  $N_0 = 10^{16} \text{ cm}^{-3}$ ,  $\delta\omega T_2 = 0.1$ .

In numerical evaluation of eq. (5) via split-step FFT method [23] the mesh size in  $X, Y$  plane was  $512 \cdot 512$  points. The "guard bands ratio" [26] was set equal to 8. The main part of the field  $\Psi$  was located inside central part of a mesh of  $64 \cdot 64$  size - the "image area". The tolerance of the energy spillover was kept within  $\epsilon_1 = 0.001$ . The "windowing" in wavenumber space after FFT at each timestep was performed by usage of "Fermi-Dirac" smoothed step function [23]. The dissipation inherent to "windowing" in split-step FFT method was compensated by nonlinear gain. The initial conditions were taken as multimode random field [23].

Just near lasing threshold the radiation mode has a distribution of intensity as rectangular grating of bright and dark spots : the latter are vortex cores. The nonlocal integral mapping proved to be successful in computation of the near field distribution as well. Quite unexpectedly in the most runs the parallel vortex lines were obtained (fig. 1) [5] rather than periodic array of bright and dark spots, typical to Talbot phenomenon[12]. The origin of parallel vortex lines is interpreted as nonlinear superposition of vortices with helicoidal phase dislocations. In the next sections the possibility of approximation of this nonlinear optical vortex lattice by linear equivalent will be outlined.

### III. TALBOT LATTICES

Consider the phase-locked rectangular lattice of zero-order Gaussian beams located at sites  $\vec{r}_{jx,jy}$  [12] separated by period  $p$ , where  $jx, jy$  - are discrete indices corresponding  $x$  and  $y$  coordinate of a given site. Let us assume for simplicity that polarization is linear, thus spin of light is zero. At  $z = 0$  plane the electric field  $E$  is given by expression:

$$E(\vec{r}, 0) = E_0 \sum_{jx,jy} \exp [-|\vec{r} - \vec{r}_{jx,jy}|^2 / (2d^2)] \quad (7)$$

After paraxial propagation of distance  $z$  the electric field  $E(\vec{r}, z)$  is transformed into:

$$E(\vec{r}, z) = E_0 \frac{i \exp [ikz]}{(1 - iz/kd^2)} \sum_{jx,jy} \exp [-\frac{|\vec{r} - \vec{r}_{jx,jy}|^2}{2d^2(1 - iz/(kd^2))}] \quad (8)$$

The constructive interference between adjacent beams produces periodic interference patterns in different  $z$ -spaced planes. The initial periodical pattern is reproduced at so-called Talbot-distances  $z_t = 2p^2/\lambda$ . In the intermediate planes  $z_t/4m$  the coarser lattices with periods  $p/m$  are produced[23, 24]. As a result a 3D lattice of bright spots is formed (fig. 2).

Each spot could serve as potential well for neutral atoms [1, 2, 5, 6], because intensity gradient will attract or repulse the atomic dipoles depending the sign of detuning of radiation frequency from resonance. At low frequencies (red-detuning) atomic dipole oscillates in-phase with trapping field and tries to align parallel to electric field. Thus potential energy of dipole  $U = -\vec{p} \cdot \vec{E}(\vec{r})/2$  is lower in local maxima of intensity and atoms are collected at bright spots. On the other hand at frequencies above the resonance (blue-detuning) atomic dipole oscillates out-of-phase and it has tendency to align anti-parallel to electric field. In this blue-detuned case the potential energy of dipole is higher in local maxima of intensity and atoms are repelled into dark regions.

Such geometry of trapping in integer and fractional Talbot planes based on superposition of co-propagating zeroth-order Gaussian beams was considered earlier[5] including possibility of manipulation of optical lattice geometry via mutual polarization [6] of constituting beams.

#### IV. ARTIFICIAL VORTEX LABYRINTH

Consider now the periodic array of Laguerre-Gaussian vortex beams with helicoidal phase dislocations (fig. 3):

$$E(\vec{r}, z=0) \approx E_0 \sum_{jx, jy} (|\vec{r} - \vec{r}_{jx, jy}|) \exp(-|\vec{r} - \vec{r}_{jx, jy}|^2/d^2) \exp(-|\vec{r}|^2/D^2) \exp[i \ell_{EM} \text{Arg}(\vec{r} - \vec{r}_{jx, jy}) + i \pi(jx + jy)]. \quad (9)$$

The topological charge  $\ell_{EM}$  is assumed to be unity, the neighbouring beams (components of the sum (9)) are  $\pi$ -shifted. The apodization function  $\exp(-|\vec{r}|^2/D^2)$  is added in order to suppress the maximum of interference pattern at the edge of array. The beams centers are placed in the centers of rectangular grid  $\vec{r}_{jx, jy}$  of period  $p$  whose axes are parallel to  $X, Y$ . The overlapping beams produce interference pattern formed by two arrays of bright and dark spots rotated at  $45^\circ$  angle with respect to initial array of LG beams. The dark spots are of two kinds: one lattice of spots coincides with lattice of initial vortices(9), the other one is produced by interference and it is shifted at distance  $p/\sqrt{2}$  along diagonal of initial lattice. The resulting interferogram has apparent  $45^\circ$  tilt compared to lattice of initial vortices (fig. 3).

The topological charges of dark spots (vortices) flip from one site to another. The interesting feature of this interference pattern is the distribution of angular momentum [13]. The initial array of LG beams carries unit circulation and corresponding angular momentum at each site. Such *chessboard-like* interferogram (fig. 3) contains additional array of vortices with alternative charges. The net angular momentum tends to be close to zero, because at the central part of array each positively directed TC  $\ell_{EM}$  is compensated by four adjacent negative ones having charge  $-\ell_{EM}$ . The elementary cell of such a lattice consists of two  $\pi$ -shifted initial vortices with co-directed charges  $\ell_{EM}$  located at diagonal and two counter-directed charges  $-\ell_{EM}$  ( $\pi$ -shifted too) placed at the other diagonal of cell. The period  $p$  of initial pattern was taken being equal to  $30\mu\text{m}$  while width of each initial beam was set  $w_{LG} = 22\mu\text{m}$  in order to provide significant mutual overlapping of vortices.

The longitudinal distribution of intensity of optical vortex array is composed of periodically spaced (with period  $p$ ) hollow tubes - vortex cores (fig. 4). Apart from Talbot gratings which are reproduced by diffraction at  $z_t = 2p^2/\lambda$  planes with corresponding period division in between planes the vortex array under consideration keeps its shape within Rayleigh range, i.e. at distances  $z < D^2/\lambda$ . Within this distance the diameter of cores is kept constant along  $z$  by virtue of interference of adjacent vortices, whose helical wavefronts are perfectly matched within elementary cell (fig. 5).

Consider first the interaction of individual atoms with single optical beam carrying topological charge. The gradient force will attract the "red"-detuned atomic dipole to the intensity maximum of an isolated first order Laguerre-Gaussian beam - "doughnut" beam (fig. 5), i.e. to the ring-shaped area around phase singularity. As a result the cloud of cold atoms will be accumulated near the maximum of intensity or "doughnut" as follows from variational solution of Gross-Pitaevsky equation (GPE) [2]:

$$i\hbar \frac{\partial \Psi(\vec{r}, t)}{\partial t} = -\frac{\hbar^2}{2m} \Delta \Psi(\vec{r}, t) + V_{ext}(\vec{r}) \Psi(\vec{r}, t) + \frac{4\pi\hbar^2 a}{m} \Psi(\vec{r}, t) |\Psi(\vec{r}, t)|^2, \quad (10)$$

with trapping potential  $V_{ext}$  of the form

$$V_{ext}(\vec{r}) = \frac{m\omega_z^2 z^2}{2} - \text{Re}[\alpha(\omega)] |E_0|^2 [r^2 \exp(-r^2/(2w_{LG}^2))] \quad (11)$$

where  $\alpha(\omega)$  is polarizability of atom[1],  $m$  - mass of particle,  $a$  - scattering length.

The approximate variational solution of GPE for ground state macroscopic wavefunction is Laguerre-Gaussian function [11]:

$$\Psi(\vec{r}, t) = |\vec{r}| \exp(-|\vec{r}|^2/(2w_r^2) - z^2/(2w_z^2) + i \ell_{BEC} \phi), \quad (12)$$

where  $\ell_{BEC}$  is the topological charge of this vortex  $\Psi$  solution. Hence the probability distribution  $|\Psi(\vec{r}, t)|^2$  of finding atom some near the point  $\vec{r} = (x, y, z)$  at the moment  $t$  is similar to the intensity distribution of trapping field as pointed out in [11]. More information could be obtained from the study of phase structure of wavefunction and comparison of topological charges of trapping field and BEC vortex. It is easy to realize using Madelung transform

$$\Psi(\vec{r}, t) = \sqrt{\rho(|\vec{r}|, \phi, t)} \exp(i\theta(|\vec{r}|, \phi, t)) \quad (13)$$

that there exists flow of "probability fluid" with velocity  $\vec{v}$  proportional (parallel) to the phase gradient lines:

$$v(\vec{r}, t) = \frac{\hbar}{m} \nabla \theta(\vec{r}, t) \quad (14)$$

Such "flow of wavefunction" occurs around the  $z$ -axis (beam axis). The "flow" described by (12) is potential and conservative as it should be for superfluid. This picture is complicated by vorticity of EM-momentum, inherent to LG-beams [9]. Strictly speaking the rotation of classical dipole around core is accelerated by nonconservative torque induced by azimuthal force circulating around vortex core. For two-level atom the value of torque  $T$  is given by following expression obtained by Babiker, Power and Allen [8]:

$$T = \hbar \ell_{EM} \Gamma \left[ \frac{I_L}{1 + I_L + \Delta^2/\Gamma^2} \right] \quad (15)$$

where  $I_L$  - normalized trapping field intensity,  $\Delta = \omega - \omega_0$  - detuning of trapping field frequency  $\omega$  from resonant frequency  $\omega_0$  of dipole. In saturation limit  $T$  is simply :  $T = \hbar \ell_{EM} \Gamma$ . In nonresonant case this torque reduces as  $\Gamma^2/\Delta^2$ . At trapping frequencies of  $Nd^{+3}$  laser  $\Delta = 10^{14} Hz$  and  $GaAs$  laser  $\Delta = 10^{13} Hz$  the  $D$ -line doublet of  $^{85}Rb$  with linewidth  $\Gamma = 5 \cdot 10^6 Hz$  [1] the torque is reduced by factor depending on laser intensity  $I_L$ . Thus due to trade-off between the saturation and detuning the torque exists both near and at large detuning from the resonance. Consequently the loop integral of the azimuthal force over circular trajectory around core  $\oint (\vec{F}_t \times \vec{r}) \cdot d\vec{l}$  is nonzero (fig. 5).

In classical picture because of this torque the dipole (or small dielectric ball) placed in the doughnut beam will move "upwards" the helicoidal phase staircase, i.e. it will rotate around LG-beam axis. The direction of rotation is determined by topological charge  $\ell_{EM}$  of the trapping beam (fig. 5). Qualitatively classical dipole is pushed by azimuthally periodic electric field - it happens because the phase of electric field oscillations at each point in circle around the center of beam is shifted with respect to neighbouring points. In the same way the small macroscopic object like glass ball placed at maximum of intensity of GL-beam will feel the "plane-wave-like" nonresonant pressure of light field. The associated azimuthal Doppler shift of the moving atom was observed experimentally recently [25]. Outside the resonance the origin of torque  $T$  is interpreted in similar way: the Poynting vector has component  $S(\vec{r})_t$  tangent to helix and local flux of electromagnetic momentum pushes dipole along phase gradient, i.e. in azimuthal direction. The local density of EM-momentum  $\vec{g} = \vec{S}/c^2$  is proportional to components of Poynting vector[9]:

$$\begin{aligned} S(\vec{r})_t &= \frac{\epsilon_0 \omega \ell_{EM} c^2}{r} |E(\vec{r})|^2 \\ S(\vec{r})_z &= \epsilon_0 c |E(\vec{r})|^2, \end{aligned} \quad (16)$$

where  $S(\vec{r})_t$  - is tangential component of Poynting vector,  $S(\vec{r})_z$  - is axial one,  $r$  - is the distance from optical vortex core,  $\omega$  - is frequency of trapping field.

Analogously because of tangential component of EM-momentum the big dielectric ball with radius larger than LG-beam core and comparable to "doughnut" radius placed in center of LG-beam would rotate with acceleration because of the such torque and corresponding transfer of the angular momentum[10].

When loaded in optical vortex lattice (fig. 6) the atomic dipole will move around the adjacent vortex core with acceleration. The radius of rotation will increase until the dipole will approach the separatrix of the velocity fields. Next after certain amount of rotations around the vortex core it could jump to another vortex using bright spots between vortices as bridge (fig. 6). As mentioned earlier the speed of rotation is increased due to light-induced torque and radius of rotation is increased as well until dipole would come to another attractive core. The migration of a dipole from one vortex to another is assisted by bright regions which serve as channels connecting vortices (fig. 3). This qualitative picture is complicated by azimuthally inhomogeneous distribution of intensity around each core.

Our aim now is to show that just described classical motion have quantum mechanical counterpart. The transfer of angular momentum results in the specific form of macroscopic wavefunction  $\Psi$  maintaining coherence all over trapping array:

$$\begin{aligned} \Psi(\vec{r}) &\approx \sum_{jx, jy} (|\vec{r} - \vec{r}_{jx, jy}|) \exp(-|\vec{r} - \vec{r}_{jx, jy}|^2/d^2) \\ &\exp(-|\vec{r}|^2/D^2) \exp[i \ell_{BEC} \text{Arg}(\vec{r} - \vec{r}_{jx, jy}) + i \pi(jx + jy)]. \end{aligned} \quad (17)$$

Next section presents the method of solution of Gross-Pitaevsky equation with vortex trapping field  $E(\vec{r})$  in the form (9).

## V. SEPARABLE VORTEX ARRAY POTENTIAL FOR BEC

In order to get closed form solution for macroscopic  $\Psi$  it is worth to mention that azimuthal accelerating force has very small value, falling as  $\Delta^{-2}$  under detuning from resonance[8]. Next let us introduce optical potential  $V_{ext}$  as a square modulus of trapping field  $E(\vec{r}_\perp, z)$ [1]. The torque  $T$  will be taken into account as a "selection" rule for choosing distribution of topological charges  $l_{BEC}$  in resulting solution.

It was shown recently [15] that Gross-Pitaevsky equation[2] in 3D-configuration:

$$i\hbar \frac{\partial \Psi(\vec{r}, t)}{\partial t} = -\frac{\hbar^2}{2m} \Delta \Psi(\vec{r}, t) + V_{ext}(\vec{r}) \Psi(\vec{r}, t) + \frac{4\pi\hbar^2 a(\vec{B})}{m} \Psi(\vec{r}, t) |\Psi(\vec{r}, t)|^2, \quad (18)$$

admits the application of standard method of separation of variables widely used for solution of *linear* partial differential equation, e.g. in quantum mechanics. The separation of variables means that wavefunction is factorized:

$$\Psi(\vec{r}, t) = \Psi_\perp(\vec{r}_\perp, t) \Psi_\parallel(z, t) \quad (19)$$

provided the Hamiltonian is the sum of two components. First component depends on a longitudinal variable  $z$  and a second component depends upon transverse variables  $\vec{r}_\perp$ .

Following to [15] in order to separate variables and factorize the wavefunction let us choose trapping potential in the following form, as a sum of components depending on longitudinal coordinate  $z$  and transverse coordinates  $\vec{r}_\perp$  separately:

$$V_{ext}(\vec{r}_\perp, z) = V_z + V_\perp = \frac{m\omega_z^2 z^2}{2} - \text{Re}[\alpha(\omega)] |E(\vec{r}_\perp)|^2 + \frac{m\omega_\perp^2 |\vec{r}_\perp|^2}{2} \quad (20)$$

where  $\alpha(\omega)$  is atomic polarizability[1] :

$$\alpha(\omega) = 6\pi\epsilon_0 c^3 \frac{\Gamma/\omega_0^2}{(\omega_0^2 - \omega^2 - i(\omega^3/\omega_0^2)\Gamma)} \quad (21)$$

The  $\alpha(\omega)$  is assumed to be real due to large negative ("red") detuning from atomic resonance at frequency  $\omega_0 = 2.4 \cdot 10^{15} \text{ Hz}$ . The imaginary part of denominator under detuning  $\Delta = \omega - \omega_0$  for  $^{85}\text{Rb}$  atoms trapped by EM-field at  $\lambda = 1.06 \div 0.808 \mu\text{m}$  is  $(3 \div 4.7) \cdot 10^8$  times smaller than real part, so the permittivity of atom  $\alpha(\omega)$  is real with good accuracy [1]. The trapping field  $E(\vec{r}_\perp)$  is assumed to be periodic function of transverse variables  $\vec{r}_\perp = (x, y)$ , composed of LG beams placed at the nodes  $(jx, jy)$  of rectangular grid of period  $p$  :

$$E(\vec{r}_\perp) \approx E_0 \sum_{jx, jy} \exp(-|\vec{r}|^2/D^2) (|\vec{r}_\perp - \vec{r}_{\perp jx, jy}|) \exp[(\vec{r}_\perp - \vec{r}_{\perp jx, jy})^2/d^2] + i\ell_{EM}\phi + i\pi(jx + jy) \quad (22)$$

The additional parabolic well with frequency  $\omega_\perp$  is introduced in (20) to get analytical solution for one trapping vortex. In order to avoid the interference between different trapping beams the usage of different carrier frequencies is recommended for longitudinal parabolic well  $\frac{m\omega_z^2 z^2}{2}$ , vortex array beam  $E(\vec{r})$  and parabolic subtrap in (20). The characteristic scales of potential in longitudinal direction and transverse direction are chosen to form "pancake trap" :  $\omega_z \gg \omega_\perp$  (fig. 1). The opposite case of elongated trap with  $\omega_z \ll \omega_\perp$  and "solitonic" longitudinal potential  $V_z \approx \text{sech}^2(z)$  was considered earlier using analogous procedure [15].

The longitudinal part of wavefunction  $\Psi_\parallel(z, t)$  is obtained as a ground state of 1D harmonic oscillator:

$$i\hbar \frac{\partial \Psi_\parallel}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \Psi_\parallel}{\partial z^2} + \frac{m\omega_z^2 z^2}{2} \Psi_\parallel \quad (23)$$

$$\Psi_{||} = (\frac{m\omega_z}{\pi\hbar})^{1/4} \exp[-m\omega_z z^2/(2\hbar) - i\omega_z t] \quad (24)$$

The transverse part of wavefunction  $\Psi_{\perp}(\vec{r}_{\perp}, t)$  is to be obtained by solving "transverse" GPE:

$$i\hbar \frac{\partial \Psi_{\perp}}{\partial t} = -\frac{\hbar^2}{2m} \Delta_{\perp} \Psi_{\perp} + V_{\perp}(\vec{r}_{\perp}) \Psi_{\perp} + \frac{4\pi\hbar^2 a(\vec{B})}{m} \Psi_{\perp} |\Psi_{\perp}|^2 [\int_{-\infty}^{\infty} |\Psi_{||}(z, t)|^4 dz] / [\int_{-\infty}^{\infty} |\Psi_{||}(z, t)|^2 dz], \quad (25)$$

where

$$V_{\perp} = \frac{m\omega_{\perp}^2 |\vec{r}_{\perp}|^2}{2} - \text{Re}[\alpha(\omega)] |E(\vec{r}_{\perp})|^2 \quad (26)$$

Because of normalization

$$\int_{-\infty}^{\infty} |\Psi_{||}(z, t)|^4 dz = 1/2 \quad \text{and} \quad \int_{-\infty}^{\infty} |\Psi_{||}(z, t)|^2 dz = 1 \quad (27)$$

The following 2D GPE results from separation of variables for "pancake" trap:

$$i\hbar \frac{\partial \Psi_{\perp}}{\partial t} = -\frac{\hbar^2}{2m} \Delta_{\perp} \Psi_{\perp} + V_{\perp}(\vec{r}_{\perp}) \Psi_{\perp} + \frac{2\pi\hbar^2 a(\vec{B})}{m} \Psi_{\perp} |\Psi_{\perp}|^2 \quad (28)$$

The scattering length  $a$  as a function of magnetic field is:

$$a(\tilde{\mathbf{B}}) = a_{bg} (1 + \frac{\Delta_B}{B - B_R}) \quad (29)$$

where  $\Delta_B$  is a width of Feshbach resonance,  $B_R$  - resonant magnetic field,  $a_{bg}$ -background scattering length[2].

Formally the separation of variables is applicable each time when Hamiltonian is a sum of components depending on different groups of variables, but this method have additional physical meaning for asymmetric potentials. The examples are elongated in  $z$ -direction trap [15] and "pancake" 2D trap (fig. 1) as in current case. The dynamics of  $\Psi$  in 2D traps was considered in large amount of papers, including the geometries of periodic potentials, Bessel lattices etc. In current case the SFV lattice under consideration has some features, qualitatively described above in discussion of classical motion of dipole around phase singularity.

The continuous transfer of angular momentum from optical vortex to BEC with wavefunction  $\Psi$  might induce the vortices in initially nonrotating BEC. Because the direction of rotation of classical particle is determined by distribution of TC of optical vortices  $\ell_{EM}$  of trapping beam the distribution of TC's in quantum superfluid lattice  $\ell_{BEC}$  will be set identical to those of trapping field.

The possible solution of equation (28) presumes the identical spatial distributions for fields  $\Psi_{\perp}(\vec{r}_{\perp}, t)$  and  $E(\vec{r}_{\perp})$ . It means that  $\Psi$  is also a sum of a LG functions with alternating topological charges  $\ell_{BEC}$  (see eq.(17)). Unfortunately the effective diameter of the core  $b$  is much larger than effective size of LG beam bottleneck (4-10  $\mu m$ ). But this discrepancy does affect the basic features of solution because of logarithmic dependence of vortex energy on vortex dimensions [17].

The correlation  $K$  of two complex spatially inhomogeneous fields  $\Psi_{\perp}(\vec{r}_{\perp}, t)$  and  $E(\vec{r}_{\perp})$  is expected to be unit:

$$K = \frac{|\int \Psi_{\perp} E^*(\vec{r}_{\perp}) d^2 \vec{r}_{\perp}|^2}{[\int |\Psi_{\perp}|^2 d^2 \vec{r}_{\perp}][\int |E(\vec{r}_{\perp})|^2 d^2 \vec{r}_{\perp}]} = 1 \quad (30)$$

The key point is in adjusting the parameters in  $\Psi_{\perp}(\vec{r}_{\perp}, t)$  and  $E(\vec{r}_{\perp})$  in such a way that two last terms in (25) would cancel each other. This might happen when following condition is imposed upon the coefficients:

$$\text{Re}[\alpha(\omega)] |E_0|^2 = \frac{2\pi\hbar^2 a(\vec{B})}{m} \quad (31)$$

The mutual compensation of these two terms could be achieved via tuning of magnetic field  $\vec{B}$  near Feshbach resonance.

Consider first the case of single vortex trap collocated with a single parabolic subtrap:

$$V_{\perp} = \frac{m \omega_{\perp}^2 r^2}{2} - \text{Re}[\alpha(\omega)] |E_0|^2 r^2 \exp[-r^2/d^2] \quad (32)$$

The following exact solution for 2D harmonic oscillator is known :

$$\Psi_{\perp} = \sqrt{\frac{2}{\pi}} \left[ \frac{m \omega_{\perp}}{\hbar} \right]^{3/2} r \exp \left[ -\frac{m \omega_{\perp}^2 r^2}{2\hbar} + i \phi \ell_{BEC} - i 2 \omega_{\perp} t \right] \quad (33)$$

which is similar to variational solution for "transverse" wavefunction (12) [11]. Note in our case the exact wavefunction of the transversal GPE found, rather than variational one (12). The stability analysis will be published elsewhere. The angular momentum per particle is given by:

$$\langle \hat{\ell} \rangle = \int \int \Psi_{\perp}^* (-i\hbar) \frac{\partial \Psi_{\perp}}{\partial \phi} d^2 r_{\perp} = \hbar \quad (34)$$

where  $\phi$  is azimuthal angle (fig. 5). Again SFV carries angular momentum  $\hbar$  per particle and kinetic energy of the whole vortex line  $E_{kin} = N\pi\hbar^2 \chi \ln(\tilde{b}/\tilde{a})/m$  [17]. Consider now the trapping of BEC by of phase-locked Gaussian-Laguerre beams placed at the nodes  $i, j$  of rectangular grid of period  $p$  ( see Eq. 9).

Let us assume for simplicity that optical wavelength  $\lambda$  is equal to De-Broglie wavelength  $\lambda_{db}$ :

$$\lambda = \lambda_{db} = \frac{\hbar}{\sqrt{2mk_B T}} \quad (35)$$

The corresponding BEC temperature for  $^{85}\text{Rb}$  atoms is:

$$T = 1.7^{-7} K \quad (36)$$

After imposing compensation condition (27) the residual part of transverse GPE (28) corresponds to free-space evolution:

$$i\hbar \frac{\partial \Psi_{\perp}}{\partial t} = -\frac{\hbar^2}{2m} \Delta_{\perp} \Psi_{\perp} + \frac{m \omega_{\perp}^2 r^2}{2} \Psi_{\perp} \quad (37)$$

The free space propagation (fig. 1) equation for EM-field will be of similar form:

$$\frac{\partial E}{\partial z} = -\frac{i}{2k} \Delta_{\perp} E + \frac{k r^2}{2f_{cavity}} E \quad (38)$$

where  $f_{cavity}$  is effective focal length of the laser cavity (fig. 1) induced by thermal lens collocated with inhomogeneity of optical pumping.

Because of linearity of equation and superposition principle the solution of ( 38 ) will be the sum of free-space modes including zero-order Gaussian functions, Gaussian-Hermit or Laguerre-Gaussian modes located at sites  $\vec{r}_{jx,jy}$  [12] separated by period  $p$ . The axes of such optical array are parallel to  $X, Y$ . In order to solve numerically eq. (28) the split-step FFT method [23] was used. The mesh size in  $X, Y$  plane was  $512 \cdot 512$  points, the "guard bands ratio" [26] was chosen equal to 8. So the main part of the field  $\Psi$  was located inside central part of a mesh of  $64 \cdot 64$  size - the "image area". The tolerance of the energy spillover was kept within  $\epsilon_1 = 0.0001$ . The windowing in wavenumber space after FFT at each timestep was performed by usage of "Fermi-Dirac" smoothed step function [23]. The dissipation inherent to split-step FFT method have led to decrease of total "amount of particles"  $\int \int |\Psi|^2 dx dy$  within "image area" at a speed of  $10^{-3}$  per time step. The special initial conditions of "preselected" SFV array in the form (17) superimposed upon homogeneous background gave the spatial distribution of "transverse" wavefunction  $\Psi_{\perp}$



well correlated ( $K \approx 0.7$ ) with the OVA array distribution - the array of phase-locked Gaussian-Laguerre wavepackets of the first order :

$$\Psi_{\perp} = \frac{2}{\pi} \left[ \frac{m\omega_{\perp}}{\hbar} \right]^{3/2} \sum_{jx,jy} (|\vec{r} - \vec{r}_{jx,jy}|) \exp(-|\vec{r}|^2/D^2) \exp\left[-\frac{m\omega_{\perp}^2 |\vec{r} - \vec{r}_{jx,jy}|^2}{2\hbar}\right] + i \ell_{BEC} \text{Arg}(\vec{r} - \vec{r}_{jx,jy}) + i \pi(jx + jy) - i 2 \omega_{\perp} t \quad (39)$$

The numerical solution for  $\Psi_{\perp}$  of GPE proved to be very close to the linear combination of Gaussian-Laguerre functions. Nevertheless this solution takes into account the interference between wavefunctions of "subvortices", because the "doughnut" radius is set to be a bit less than distance between lattice nodes. The arising interference pattern is well correlated with the interference pattern produced by Laguerre-Gaussian OVA  $|E(r_{\perp})|^2$  (fig. 3) with the same geometrical parameters and wavelength  $\lambda$ . This solution is stable with respect to small harmonic perturbations.

Each SFV carries angular momentum  $\hbar$  per particle and rotational kinetic energy  $E_{kin} = N\pi\hbar^2 \chi \ln(\tilde{b}/\tilde{a})/m$ ,  $\tilde{b}$  - diameter of vortex core,  $\tilde{a}$  - interatomic distance,  $\chi$  - length of vortex which is roughly equal to thickness of "pancake" [17]. The energy associated with superfluid vortices is of order  $10^{-(19 \div 20)} J$  or  $0.1 \div 1 eV$  at density of dilute  $^{85}Rb$  gas  $N \approx 10^{16 \div 18} cm^{-3}$ . In contrast to superfluid in rotating bucket where angular momenta of vortices are co-directed [16], the BEC vortices trapped by optical vortex array are counter-directed from site to site. Consequently in the net sum of angular momenta each vortice of positive topological charge is compensated by the term with negative charge and total angular momentum tends to zero. Nevertheless the mutual subtraction of angular momenta (vectorial) of adjacent vortices in the net sum does not mean the mutual subtraction of rotational energies, which are the positive scalars. The ground state carry substantial amount of rotational kinetic energy of condensate which contains  $N$  particles of mass  $m$  per unite volume, namely  $E_{rot} = N_{vortices} N \pi \hbar^2 \chi \ln(\tilde{b}/\tilde{a})/m$ .

## VI. CONCLUSION

The optical vortex arrays emitted by solid-state microchip laser are analysed from the point of view of application to optical dipole traps. The numerical modeling gave the field distributions highly correlated with experimentally observed ones. The array of phase-locked Gaussian-Laguerre beams equispaced at the nodes of rectangular lattice proved to be a reasonable approximation for experimental and numerical results as well. For macroscopic wavefunction  $\Psi$  of BEC trapped in such complex optical field the analytical solutions of Gross-Pitaevsky equation were found based on separation of variables and mutual compensation of vortex component of external trapping field via nonlinear term of GPE. The obtained wavefunctions have perfect correlation with trapping field, including distribution of topological charges, which form "antiferromagnetic-like" lattice. Within framework of this particular model the "antiferromagnetic" lattice of BEC vortices carriers total angular momentum close to zero while net rotational kinetic energy of SFV lattice tends to be equal to the sum of rotational energies of vortices.

The field of classical velocities, i.e. the field of phase gradients, obtained via Madelung transform, forms labyrinth structure. In classical picture the trapped particle moves across the vortex array labyrinth jumping from one whirl to another.

In quantum picture represented via analytic solution of GPE for one trapping vortex and numerical solution for optical vortex array the coherent macroscopic wavefunction extends all over OV trapping array with transverse spatial dimension of several hundred microns. The complex field of velocities, rotational energy and high degree of correlation of SFV wavefunction with OV trapping field promise more resistance to decoherence.

The qualitative analytic solution supports the basic feature related to trapping of macroscopic particles and to BEC trapping: the transfer of OAM from EM-field to superfluid. The proposed OVA trap might be interesting from the point of view of studies of quantum-classical correspondence.

The mechanism of imposing the topological charges to BEC vortices by means of manipulating the vorticity of trapping optical array could result in demonstration of macroscopic quantum interference phenomena. Evidently there are 4 possible topologically equivalent combinations of parameters of solution (39) of OV charges ( $\ell_{EM} = \pm 1$ ) and their relative phases ( $\pm \pi$ ). Thus there exist 4 wavefunctions  $\Psi$  having an identical probability  $|\Psi|^2$  distribution and different orientation of SFL vortices with respect to physical axes  $X, Y$  of trapping setup, characterised by  $\ell_{BEC}$  and their relative phases. The transformation of the one  $\Psi$  into another one having different phase structure is equivalent to  $90^\circ$  rotation around  $z$  - axis.

---

[1] R.Grimm, M.Weidemuller and Yu.B.Ovchinnikov., "Optical dipole traps for neutral atoms" *Adv.At.Mol.Opt.Phys.* **42**, 95 (2000).

- [2] L. Pitaevskii and S. Stringari., *"Bose-Einstein Condensation"*, Clarendon Press, Oxford, 2003.  
F. Dalfovo, S. Giorgini, S. Stringari, L.P. Pitaevskii.,  
*Rev. Mod. Phys.* **71**, 463 (1999).
- [3] Balykin, V.I., Letokhov, V.S., Ovchinnikov, Yu.B., and Sidorov A.I., *Phys. Rev. Lett.* **60**, 2137 (1988),
- [4] E. M. Wright, J. Arlt and K. Dholakia., *"Toroidal Optical Dipole Traps for Atomic Bose-Einstein Condensates using Laguerre-Gaussian Beams"*, *Phys. Rev. A* **63**, 013608 (2001).
- [5] A.Yu. Okulov., *"3D-configuration of the vortex lattices in microchip laser cavity"*,  
QCMC-2004, AIP Conference Proceedings, **734**, p.366 (2004).  
A.Yu. Okulov., *"3D structure of the vortex lattices in solid-state microchip laser"*, Bulletin Lebedev Physical Institute, **9**,  
p.3, Sept. (2003).
- [6] Yu.B. Ovchinnikov., *Phys. Rev.*, **73A**, 033404 (2006).
- [7] R. Dumke, M. Volk, T. Muther, F.B.J. Buchkremer, G. Birkel and W. Ertmer., *Phys. Rev. Lett.* **89**, 097903 (2002).
- [8] M. Babiker, W. L. Power, and L. Allen., *Phys. Rev. Lett.* **73**, 1239 (1994).
- [9] V. Garcés-Chavez, D. McGloin, M. J. Padgett, W. Dultz, H. Schmitzer, and K. Dholakia., *Phys. Rev. Lett.* **91**, 093602 (2003).
- [10] M.J. Friese, J. Euger, H. Rubinstein-Dunlop., *Phys. Rev. A* **54**, 1543 (1996).  
N.R. Heckenberg, M.E.J. Friese, T.A. Nieminen and H. Rubinsztein-Dunlop., *"Mechanical Effects of Optical Vortices"*.  
pp 75-105, in M. Vasnetsov (ed), *Optical Vortices*, Nova Science Publishers (1999).
- [11] J. Tempere and J. T. Devreese, E. R. I. Abraham., *Phys. Rev.*, **64A**, 023603 (2002).
- [12] A.Yu. Okulov., *"Two-dimensional periodic structures in nonlinear resonator"*, *JOSA B* **7**, p.1045, (1990).
- [13] J. Courtial, R. Zambrini, M.R. Dennis, M. Vasnetsov., *"Angular momentum of optical vortex arrays"*, *Optics Express* **14**, p.938 (2006).
- [14] Y.F. Chen, Y.P. Lan., *Phys. Rev.*, **64A**, 063807 (2001).,  
Y. F. Chen, Y. P. Lan., *Phys. Rev.*, **65**, 013802 (2001).
- [15] R. Fidele, P.K. Shukla, S. De Nicola, M.A. Manko, V.I. Manko, F.S. Cataliotti. *JETP Lett.*, **80**, 8, p.609-613 (2004).
- [16] J.R. Abo-Shaer, C. Raman, J.M. Vogels and W. Ketterle., *Phys. Rev. Lett.* **88**, 070409 (2002).
- [17] R.P. Feynman, *"Statistical mechanics"*, Ch.11, Reading, Massachusetts (1972).
- [18] A.Yu. Kitaev., LANL e-print quant-ph/ 9707021, <http://arxiv.org>, (1997).
- [19] A.F. Suchkov., *JETP* **22**, 1026 (1965).
- [20] A.E. Siegman., *"Lasers"*. University Science Books, Mill Valley, California (1986).
- [21] K. Staliunas., *Phys. Rev. A* **48**, 1573 (1993). K. Staliunas, C.O. Weiss., *"Nonstationary vortex lattices in large-aperture class B lasers"*, *JOSA B* **12**, 1142 (1995).
- [22] F. Hollinger, Chr. Jung, and H. Weber., *"Simple mathematical model describing multitransversal solid-state laser"*, *JOSA B* **7**, 1013 (1990).
- [23] A.Yu. Okulov., *"Scaling of diode-pumped solid-state lasers via self-imaging"*, *Opt. Comm.*, **99**, p.350-354 (1993).,  
*J. Mod. Opt.*, **38**, N.10, p.1887 (1991).
- [24] A.Yu. Okulov., *"Gain-mode correlation in diode-pumped solid-state laser"*, *Optics and Spectroscopy.*, **77**, N6, p.985 (1994).
- [25] V. Garcés-Chavez, K. Volke-Sepulveda, S. Chavez-Cerda, W. Sibbett and K. Dholakia., *Phys. Rev. A* **66**, 063402 (2002).
- [26] E.A. Sziklas, A.E. Siegman., *"Mode calculations in unstable resonators with flowing saturable gain. 2: Fast Fourier transform method"*, *Appl. Opt.*, **14**, 1874 (1975).

### List of Figure Captions

Fig.1. Conceptual view of the near field optical trap. Upper plot: Transverse (in XOY plane) distribution of intensity in the near field of solid-state microchip laser [5]. Middle: The longitudinal scale extends to 6 Talbot lengths. Z - axis is directed along optical axis of microchip laser resonator(bottom) . Additional tightly confined parabolic well keeping BEC cloud localized in z - direction is depicted via potential  $V(z) \approx (z - z_0)^2$ . Such potential is assumed to be superimposed by the other microchip laser beam with cylindrical focusing lens at slightly different wavelength of radiation from the range  $0.98 \div 2.79 \mu m$ , in order to avoid interference. Fresnel number has value of  $N_f \approx N_{vortices}^2 = 64$ .

Fig.2.Diffractive self-imaging of two-dimensional lattice of 8x8 Gaussian beams with period  $p = 28 \mu m$  . The longitudinal cross-section at  $y = 0$  plane of intensity distribution  $I(x, y, z)$  presented. The lattice is self-reproduced at Talbot distance, spatial period division occurs at quarter Talbot distances and central lobe forms outside the Rayleigh range.

Fig.3.Intensity distribution in transverse plane of artificial Laquerre-Gaussian vortex array. Vortices(dark holes) are at nodes of rectangular *chessboard – like* grid. Letters  $\ell_{EM}$  denote vortices with alternating topological charges, changing sign from one site of lattice to another.

Fig.4.Longitudinal section of vortex array in the near field in  $x$ - $z$  plane at  $y = 0$  section. The vortex lines are parallel, the topological charges  $\ell_{EM}$  are flipping from one vortex line to another. The wavelength  $\lambda = 1 \mu m$ , period  $p$  of lattice both in  $x$  and  $y$  directions is  $30 \mu m$ .

Fig.5. a)Helicoidal phase surface of Gaussian-Laquerre beam. Atomic dipole having "red" detuning moves along phase gradient. Trajectory is located at maximum of intensity.  $\vec{S}_t$  is the component of Pointing vector tangent to helix. The major component of  $\vec{S}_z$  is directed parallel to beam propagation, i.e. along Z-axis. b)Helicoidal phase surface of Laquerre-Gaussian beam array. The elementary cell consists of four vortices. The adjacent vortices have alternating topological charges  $\ell_{EM}$  and alternating angular momenta. In classical picture the atomic dipole moves along phase gradient of a given vortex next it jumps to another one.

Fig.6.The argument of macroscopic wavefunction  $\Psi$  and corresponding field of velocities obtained via Madelung transform . The superfluid vortices form the lattice with alternating topological charges  $\ell_{BEC} = \pm 1$ . The elementary cell consists of four BEC vortices whose locations are identical to the vortices of trapping EM-field. Horizontal pair has the same charges  $\ell_{BEC}$  which are  $\pi$ -shifted with respect to each other, the vertical pair have  $\pi$ -shifted  $-\ell_{BEC}$  charges. The alternating charges  $\ell_{BEC}$  make the field of velocities continuous.

Fig.7.The distribution of the square modulus of macroscopic wavefunction  $\Psi$  obtained via numerical modeling. The location of adjacent counter-rotating vortices  $\ell_{BEC}$  shown.




FIG. 1: Conceptual view of the near field optical trap. Upper plot: Transverse (in XOY plane) distribution of intensity in the near field of solid-state microchip laser [5]. Middle: The longitudinal scale extends to 6 Talbot lengths. Z - axis is directed along optical axis of microchip laser resonator(bottom) . Additional tightly confined parabolic well keeping BEC cloud localized in  $z$  - direction is depicted via potential  $V(z) \approx (z - z_0)^2$ . Such potential is assumed to be superimposed by the other microchip laser beam with cylindrical focusing lens at slightly different wavelength of radiation from the range  $0.98 \div 2.79 \mu m$ , in order to avoid interference. Fresnel number has value of  $N_f \approx N_{vortices}^2 = 64$ .




FIG. 2: Diffractive self-imaging of two-dimensional lattice of  $8 \times 8$  Gaussian beams with period  $p = 28 \mu m$ . The longitudinal cross-section at  $y = 0$  plane of intensity distribution  $I(x, y, z)$  presented. The lattice is self-reproduced at Talbot distance, spatial period division occurs at quarter Talbot distances and central lobe forms outside the Rayleigh range.




FIG. 3: Intensity distribution in transverse plane of artificial Laquerre-Gaussian vortex array. Vortices (dark holes) are at nodes of rectangular *chessboard*–like grid. Letters  $l_{EM}$  denote vortices with alternating topological charges, changing sign from one site of lattice to another.




FIG. 4: Longitudinal section of vortex array in the near field in  $x$ - $z$  plane at  $y = 0$  section. The vortex lines are parallel, the topological charges  $\ell_{EM}$  are flipping from one vortex line to another. The wavelength  $\lambda = 1\mu m$ , period  $p$  of lattice both in  $x$  and  $y$  directions is  $30\mu m$




FIG. 5: a) Helicoidal phase surface of Laquerre-Gaussian beam. Atomic dipole having "red" detuning moves along phase gradient. Trajectory is located at maximum of intensity.  $\vec{S}_t$  is the component of Pointing vector tangent to helix. The major component of  $\vec{S}_z$  is directed parallel to beam propagation, i.e. along  $Z$ -axis. b) Helicoidal phase surface of Laquerre-Gaussian beam array. The elementary cell consists of four vortices. The adjacent vortices have alternating topological charges  $\ell_{EM}$  and alternating angular momenta. In classical picture the atomic dipole moves along phase gradient of a given vortex next it jumps to another one.




FIG. 6: The argument of macroscopic wavefunction  $\Psi$  and corresponding field of velocities obtained via Madelung transform. The superfluid vortices form the lattice with alternating topological charges  $\ell_{BEC} = \pm 1$ . The elementary cell consists of four BEC vortices whose locations are identical to the vortices of trapping EM-field. Horizontal pair has the same charges  $\ell_{BEC}$  which are  $\pi$ -shifted with respect to each other, the vertical pair have  $\pi$ -shifted  $-\ell_{BEC}$  charges. The alternating charges  $\ell_{BEC}$  make the field of velocities continuous.




FIG. 7: The distribution of the square modulus of macroscopic wavefunction  $\Psi$  obtained via numerical modeling. The location of adjacent counter-rotating vortices  $\ell_{BEC}$  shown.