# Weak compressibility of surface wave turbulence

# MARIJA VUCELJA<sup>1</sup> AND ITZHAK FOUXON<sup>1,2</sup>

<sup>1</sup>Physics of Complex Systems, Weizmann Institute of Science, Rehovot, 76100, Israel <sup>2</sup>Racah Institute of Physics, Hebrew University of Jerusalem, Jerusalem, 91904, Israel

(Received 1 October 2018)

We study the growth of small-scale inhomogeneities of the density of particles floating in weakly nonlinear, small-amplitude, surface waves. Despite the amplitude smallness, the accumulated effect of the long-time evolution may produce strongly inhomogeneous distribution of the floaters: density fluctuations grow exponentially with a small but finite exponent. We show that the exponent is of sixth or higher order in wave amplitude. As a result, the inhomogeneities do not form within typical time-scales of the natural environment. We conclude that turbulence of surface waves is weakly compressible and alone it cannot be a realistic mechanism of the clustering of matter on liquid surfaces.

### 1. Introduction

Clustering of matter on the surface of lakes and pools and of oil slicks and seaweeds on the sea surface is well-known empirically while there is no theory that describes it. Since surface flows are compressible even for incompressible fluids, such theory should be based on the general description of the development of density inhomogeneities in a compressible flow. Important characteristics of the formation of small-scale inhomogeneities is minus the sum of the Lyapunov exponents of the flow,  $\lambda$ . It gives the asymptotic logarithmic growth rate of the density on fluid particle trajectories at large times. The rate  $\lambda$  is minus the average value of the velocity divergence seen by a fluid particle, and it is always non-negative because contracting regions with negative divergence have more particles and hence larger statistical weight, see Balkovsky et al. (2001); Ruelle (1996, 1997, 1999). We note that  $\lambda$  is also the production rate of the Gibbs entropy, so that the condition  $\lambda \geqslant 0$  can be regarded as an analogue of the second law of thermodynamics for the dissipative dynamics, see Ruelle (1996, 1997, 1999); Falkovich & Fouxon (2003, 2004). For a generic flow one has  $\lambda > 0$  and the asymptotic density becomes a singular measure, the so-called Sinai-Ruelle-Bowen measure, see e.g. Dorfman (1999). For floaters this means that they form a multi-fractal structure on the surface, see Yu et al. (1991); Falkovich et al. (2001); Balkovsky et al. (2001); Ruelle (1999); Falkovich & Fouxon (2003, 2004); Bec et al. (2004); Balk et al. (2004); Eckhardt & Schumacher (2001) for theory and Ramshankar et al. (1990); Sommerer & Ott (1993); Sommerer (1996); Nameson et al. (1996); Schroder et al (1996); Cressman & Goldburg (2003); Denissenko et al. (2006); Bandi et al. (2006) for experiments. This structure is the attractor of the two-dimensional dissipative dynamics obeyed by the particles on the surface, and it evolves constantly for time-dependent flows, see e.g. Dorfman (1999); Ott (2002). The Kaplan-Yorke dimension of the attractor,  $D_{KY} = 1 + \lambda_1/|\lambda + \lambda_1|$ , is between one and two, assuming that the principal Lyapunov exponent  $\lambda_1$  is positive, see Ott (2002).

Surface flows are generic compressible flows for which the Eulerian compressibility, measured by the dimensionless ratio  $C = \langle (\partial_i v_i)^2 \rangle / \langle (\partial_i v_i)^2 \rangle$ , is of order one, cf. Boffetta et al. (2006a). Here angular brackets stand for the spatial average, v is the floaters velocity field, and C changes from zero for incompressible flow to one for potential flow. One expects  $\lambda \sim \lambda_1$  from  $C \sim 1$ , so that the deviation of  $D_{KY}$  from the surface dimension 2 is also of order one,  $2 - D_{KY} \sim 1$ . The expectation holds true for the flow on the surface of three-dimensional turbulence. Performing numerical simulations with the full 3d Navier-Stokes equations, Boffetta et al. (2006a) found  $C \approx 0.5$ ,  $D_{KY} \approx 1.15$  and observed strong clustering on the surface, see also Cressman et al. (2004); Boffetta et al. (2006b); Eckhardt & Schumacher (2001). However, underwater turbulence is relatively rare in the natural environment (due to stable stratification), and it is important to consider other cases of the surface flows, of which the small-amplitude surface waves are probably the most wide-spread. Despite the amplitude smallness, a small-but-finite  $\lambda$  produces a large effect over time-scales of order  $1/\lambda$  and larger. Thus to estimate the role of surface waves in the formation of the floaters inhomogeneities in the natural environment, one needs to know how small  $\lambda$  is. In this Article we show that  $\lambda$  is of sixth or higher order in wave amplitude. Let us note that for surface waves the degree of compressibility C is due to linear waves, which produce potential flow with C=1. Thus one could expect that the estimates  $\lambda \sim \lambda_1$  and  $2 - D_{KY} \sim 1$  would hold for surface waves like they do for the underwater turbulence. We show that in fact, under some natural non-degeneracy assumptions described in the Discussion, for surface waves  $\lambda \ll \lambda_1$ .

The calculation of  $\lambda$  for random waves in different situations was considered in Balk et al. (2004); Vucelja et al. (2006). For surface waves, in the linear order in the wave amplitude, the particles move periodically, see e.g. Batchelor (2002), hence there is no net clustering. Thus analysis of clustering of the floaters demands the account of the nonlinear effects. Balk et al. (2004) assumed a linear relation between the velocity field of the floaters and the wave amplitudes, and considered a Gaussian ensemble of non-interacting waves. The nonlinearity in this case comes from expressing the Lagrangian objects in terms of the Eulerian ones. It was shown that  $\lambda$  vanishes in the fourth order in the wave amplitude for longitudinal waves, whose dispersion relation does not allow the same frequency for two different wave-vectors (e.g. sound, gravity, capillary waves). On the other hand, the lowest order non-vanishing contribution into  $\lambda_1$  was shown to be always of the fourth order in the wave amplitude. Under the same assumption of the linear relation between the velocity and the wave amplitudes, Vucelja et al. (2006) demonstrated that the account of the wave interactions does not change the conclusion of Balk et al. (2004) on the vanishing of  $\lambda$  in the fourth order in the wave amplitude.

The results above are inconclusive as far as surface waves are concerned, for which the relation between the velocity and the amplitudes is nonlinear due to a small but finite curvature of the surface, see Zakharov (1966, 1968). For this case a separate calculation of  $\lambda$  is needed. Here we provide such a calculation. We consider weakly nonlinear surface waves and show that neither the wave interactions, nor the nonlinear relation between the velocity and the amplitudes, create non-zero sum of the Lyapunov exponents up to the fourth order in the wave amplitudes. The main tool of our analysis is a recently derived Green-Kubo type formula for the sum of the Lyapunov exponents, see Falkovich & Fouxon (2004, 2003). This formula expresses  $\lambda$  in terms of the correlations of the flow divergence in the particle frame. It describes the interplay between the particle motion and the local flow compressions in accumulating density inhomogeneities which become pronounced as a result of the long-time evolution. The slowness of the particle drift from its initial position allows us to express the correlations in terms of the Eulerian correlation functions of the velocity, which we further evaluate by a lengthy,

but straightforward calculation, cf. Vucelja et al. (2006). The following text is organized as follows. In the next Section we introduce the expression for  $\lambda$  in terms of the Eulerian correlations of the surface flow velocity, valid up to the fourth order in the wave amplitude. To perform the calculation one needs to express the velocity in terms of the normal coordinates. This is dealt with in Section 3. The calculation of the various terms occurring in  $\lambda$  is performed in Section 4. Discussion finishes the article.

#### 2. The sum of the Lyapunov exponents

Behavior of the density  $n(t, \boldsymbol{x})$  in a velocity field  $\boldsymbol{v}(t, \boldsymbol{x})$  is governed by the continuity equation  $\partial_t n + \nabla \cdot (\boldsymbol{v}n) = 0$ , see Batchelor (2002). Introducing Lagrangian particle trajectories by the equation  $\partial_t \boldsymbol{X}(t, \boldsymbol{x}) = \boldsymbol{v}[t, \boldsymbol{X}(t, \boldsymbol{x})]$  with the initial condition  $\boldsymbol{X}(0, \boldsymbol{x}) = \boldsymbol{x}$ , we may write the solution for the density as  $n[t, \boldsymbol{X}(t, \boldsymbol{x})] = n(0, \boldsymbol{x}) \exp[-\int_0^t w[t', \boldsymbol{X}(t', \boldsymbol{x})]dt']$ . Here we introduced  $w(t, \boldsymbol{x}) \equiv \nabla \cdot \boldsymbol{v}(t, \boldsymbol{x})$ . We characterize the growth of spatial inhomogeneities by the asymptotic logarithmic growth rate  $\lambda$  at large times, defined by

$$\lambda = \lim_{t \to \infty} \frac{1}{t} \ln \left[ \frac{n[t, \boldsymbol{X}(t, \boldsymbol{x})]}{n(0, \boldsymbol{x})} \right] = -\lim_{t \to \infty} \frac{1}{t} \int_0^t w[t', \boldsymbol{X}(t', \boldsymbol{x})] dt'. \tag{2.1}$$

The above limit is well-defined, Dorfman (1999), and it gives minus the sum of the Lyapunov exponents of the flow v(t, x). It was shown in Falkovich & Fouxon (2004, 2003) that if v(t, x) is a random, spatially homogeneous, stationary flow, then one has

$$\lambda = \int_0^\infty dt \langle w(0, \boldsymbol{x}) w[t, \boldsymbol{X}(t, \boldsymbol{x})] \rangle. \tag{2.2}$$

We shall apply the above formula to the case where  $\boldsymbol{v}(t,\boldsymbol{x})$  is the two-dimensional velocity field governing the motion of the floaters in a (quasi-)stationary ensemble of weakly nonlinear surface waves sustained by some forcing, see Zakharov et~al. (1992). We first use the amplitude smallness to express the Lagrangian correlation function in (2.2) in terms of the velocity correlation functions given in the Eulerian frame. We follow Vucelja et~al. (2006) who considered (2.2) in the case of arbitrary low-amplitude waves. For dispersion relation  $\Omega_{\boldsymbol{k}}$ , considering packets with both the wavenumber and the width of order k, the correlation time of  $\boldsymbol{w}$  can be estimated as  $\Omega_{\boldsymbol{k}}^{-1}$  and the correlation length as  $k^{-1}$ . The particle deviation from the initial position,  $\boldsymbol{X}(t,\boldsymbol{x})-\boldsymbol{x}$ , during the period,  $t\simeq\Omega_{\boldsymbol{k}}^{-1}$ , is  $\epsilon=kv/\Omega_{\boldsymbol{k}}\ll 1$  times smaller than  $k^{-1}$  which allows to expand (2.2) near  $\boldsymbol{x}$ . Performing the expansion to order  $\epsilon^4$  we find

$$\lambda \approx \lambda_2 + \lambda_3 + \lambda_4 \,, \tag{2.3}$$

$$\lambda_2 \equiv \int_0^\infty dt \langle w(0)w(t)\rangle \,, \quad \lambda_3 \equiv \int_0^\infty dt \int_0^t dt_1 \left\langle w(0) \frac{\partial w(t)}{\partial x^\alpha} v^\alpha(t_1) \right\rangle \,, \tag{2.4}$$

$$\lambda_4 \equiv \int_0^\infty dt \int_0^t dt_1 \int_0^{t_1} dt_2 \left\langle w(0) v_\beta(t_2) \left( \frac{\partial w(t)}{\partial x_\alpha} \frac{\partial v_\alpha(t_1)}{\partial x_\beta} + \frac{\partial^2 w(t)}{\partial x_\alpha \partial x_\beta} v_\alpha(t_1) \right) \right\rangle. (2.5)$$

Here and below we suppress the spatial coordinate x over which the averaging is performed. The expansion above was introduced in Vucelja et~al.~(2006). We mention that all contributions  $\lambda_i$  are of fourth or higher order in wave amplitude, see below. To use the above formula to find  $\lambda$  to order  $\epsilon^4$ , we need to establish the expression for the surface flow v to order  $\epsilon^3$ .

### 3. The velocity field of the floaters

The velocity field that governs the evolution of the floaters coordinates  $\mathbf{r} = (x, y)$  in the horizontal plane, has the following form

$$\mathbf{v}(\mathbf{r},t) = \left(\frac{\partial \phi(\mathbf{r},z,t)}{\partial x} \left[z = \eta(\mathbf{r},t)\right], \frac{\partial \phi(\mathbf{r},z,t)}{\partial y} \left[z = \eta(\mathbf{r},t)\right]\right), \tag{3.1}$$

where  $\eta(\mathbf{r},t)$  is the surface elevation and  $\phi(\mathbf{r},z,t)$  is the velocity potential,  $\mathbf{v} = \nabla \phi$ . Zakharov (1966, 1968) showed that the system of weakly interacting surface waves is a Hamiltonian system with canonically conjugate coordinates  $\eta(\mathbf{r},t)$  and  $\psi(\mathbf{r},t) \equiv \phi(\mathbf{r},\eta(\mathbf{r},t),t)$ . Up to the order  $\epsilon^3$ , the surface flow of the floaters is given by

$$v^{\alpha} = i \int \frac{d\mathbf{k}_1}{(2\pi)^2} e^{i\mathbf{k}_1 \cdot \mathbf{r}} k_1^{\alpha} \psi_1 - i \int \frac{d\mathbf{k}_{12}}{(2\pi)^4} e^{i(\mathbf{k}_1 + \mathbf{k}_2) \cdot \mathbf{r}} |k_1| k_2^{\alpha} \psi_1 \eta_2$$

$$(3.2)$$

$$-\frac{i}{2} \int \frac{d\mathbf{k}_{123}}{(2\pi)^6} e^{i(\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3) \cdot \mathbf{r}} |k_1| \left( |k_1| k_2^{\alpha} + |k_1| k_3^{\alpha} - 2\sqrt{k_1^2 + k_2^2} k_3^{\alpha} \right) \psi_1 \eta_2 \eta_3 , \quad (3.3)$$

cf. Zakharov (1968) formula (1.8). We introduced shorthand notations  $\eta_i(t) = \eta(\mathbf{k}_i, t)$ ,  $\psi_i(t) = \psi(\mathbf{k}_i, t)$  and  $d\mathbf{k}_{ijl...} = d\mathbf{k}_i d\mathbf{k}_j d\mathbf{k}_l...$  Note that the velocity field on the surface  $\mathbf{v}(\mathbf{r}, t)$  is neither potential, nor solenoidal. We now return to the expression for the sum of the Lyapunov exponents (2.3).

# 4. Calculation of the sum of the Lyapunov exponents

In this Section we provide the calculation of  $\lambda$  to the fourth order in wave amplitude, based on the calculation of the different contributions  $\lambda_i$ , see (2.3)-(2.5). Some parts of this analysis deal with the same objects as those considered in Balk *et al.* (2004); Vucelja *et al.* (2006), however our analysis is different and it uses specific properties of surface waves. Below we provide a more detailed calculation, as compared to the short articles Balk *et al.* (2004) and Vucelja *et al.* (2006).

# 4.1. Contribution of the four-point correlation functions

We first consider the contribution  $\lambda_4$  to  $\lambda$ . To the fourth order in wave amplitude we may assume Gaussian non-interacting waves and use Wick's theorem to decouple the averages. Employing identities like  $\langle v_{\alpha}(t_1)\partial_{\alpha}\partial_{\beta}w(t)\rangle = -\langle (\partial_{\beta}v_{\alpha}(t_1))(\partial_{\alpha}w(t))\rangle$ , that follow by integration by parts, one finds

$$\lambda_4 = -\int_0^\infty dt \int_0^t dt_1 \int_0^{t_1} dt_2 \left[ \left\langle w(0) \frac{\partial w(t)}{\partial x_\alpha} \right\rangle \left\langle v_\alpha(t_1) w(t_2) \right\rangle + \left\langle w(0) \frac{\partial v_\alpha(t_1)}{\partial x_\beta} \right\rangle \left\langle w(t) \frac{\partial v_\beta(t_2)}{\partial x_\alpha} \right\rangle$$

$$+ \left\langle \frac{\partial w(0)}{\partial x_{\alpha}} \frac{\partial w(t)}{\partial x_{\beta}} \right\rangle \left\langle v_{\alpha}(t_{1}) v_{\beta}(t_{2}) \right\rangle + \left\langle w(t_{2}) \frac{\partial w(t)}{\partial x_{\alpha}} \right\rangle \left\langle v_{\alpha}(t_{1}) w(0) \right\rangle \right]. \tag{4.1}$$

Here we do not assume isotropy of the waves. Isotropy would make terms like  $\langle v_{\alpha}(t_1)w(t_2)\rangle$  vanish. In the considered order,  $\mathbf{v} = \nabla \psi$  is a potential field and spectral representation of the pair-correlation function gives

$$\langle \psi(0)\psi(t)\rangle = \int \frac{d\mathbf{k}}{(2\pi)^2} E(\mathbf{k}) \cos\left(\Omega_{\mathbf{k}} t\right) \Rightarrow \langle v_{\alpha}(0)v_{\beta}(t)\rangle = \int \frac{d\mathbf{k}}{(2\pi)^2} k_{\alpha} k_{\beta} E(\mathbf{k}) \cos\left(\Omega_{\mathbf{k}} t\right),$$

and similar expressions for other correlation functions in (4.1). Note that the potentiality of surface waves, holding in the Gaussian approximation, makes the velocity spectrum

vanish at k=0 even if  $E(k=0)\neq 0$  [cf. Balk et al. (2004); Vucelja et al. (2006)]. We find

$$\lambda_{4} = \int \frac{d\mathbf{k} d\mathbf{q} E(\mathbf{k}) E(\mathbf{q})}{(2\pi)^{4}} \int_{0}^{\infty} dt \int_{0}^{t} dt_{1} \int_{0}^{t_{1}} dt_{2} \left[ k^{4} q^{2} (\mathbf{k} \cdot \mathbf{q}) \sin \left[ \Omega_{\mathbf{k}} t \right] \sin \left[ \Omega_{\mathbf{q}} (t_{1} - t_{2}) \right] - k^{2} q^{2} (\mathbf{k} \cdot \mathbf{q})^{2} \cos \left[ \Omega_{\mathbf{k}} t_{1} \right] \cos \left[ \Omega_{\mathbf{q}} (t_{1} - t_{2}) \right] - k^{4} (\mathbf{k} \cdot \mathbf{q})^{2} \cos \left[ \Omega_{\mathbf{k}} t \right] \cos \left[ \Omega_{\mathbf{q}} (t_{1} - t_{2}) \right] + k^{4} q^{2} (\mathbf{k} \cdot \mathbf{q}) \sin \left[ \Omega_{\mathbf{k}} (t - t_{2}) \right] \sin \left[ \Omega_{\mathbf{q}} t_{1} \right] \right].$$

$$(4.2)$$

To calculate the time-integrals we represent the products of trigonometric functions above as sums or differences of cosine functions and use

$$\int_{0}^{\infty} dt \int_{0}^{t} dt_{1} \int_{0}^{t_{1}} dt_{2} \cos(at + bt_{1} + ct_{2}) = -\frac{\pi\delta(a)}{b(b+c)} + \frac{\pi\delta(a+b)}{bc} - \frac{\pi\delta(a+b+c)}{c(b+c)}, \quad b \neq -c,$$

$$\int_{0}^{\infty} dt \int_{0}^{t} dt_{1} \int_{0}^{t_{1}} dt_{2} \cos(at - bt_{1} + bt_{2}) = -\frac{\pi\delta'(a)}{b} + \frac{\pi\delta(a)}{b^{2}} - \frac{\pi\delta(a-b)}{b^{2}}.$$
(4.3)

All terms which are supported only at the zero frequency in the frequency representation  $(\delta$ -functions or their derivatives) are also supported only at the zero wavenumber, since the dispersion relation of surface waves vanishes at k=0 only. As a result, due to the presence of positive powers of k in (4.2), these terms vanish (similarly to the vanishing of the velocity spectrum at k=0 shown above). It is then easy to see that the first and the fourth terms in  $\lambda_4$  (having the same dependence on the wavevectors) cancel each other, while the second and the third terms give

$$\lambda_4 = \int \frac{d\mathbf{k}d\mathbf{q}}{(2\pi)^4} E(\mathbf{k}) E(\mathbf{q}) k^2 (\mathbf{k} \cdot \mathbf{q})^2 (k^2 - q^2) \left( \frac{\pi \delta(\Omega_{\mathbf{k}} - \Omega_{\mathbf{q}})}{2\Omega_{\mathbf{q}}^2} \right) . \tag{4.4}$$

Since for surface waves the equality of the frequencies of two waves implies the equality of their wavelengths, then the above terms cancel each other,  $\lambda_4 = 0$ . This reproduces in a simple way the result of Balk *et al.* (2004) which says that in the Gaussian approximation  $\lambda$  vanishes for potential waves having the property that the equality of the frequencies implies the equality of the wavelengths.

## 4.2. Contribution of the two-point correlation function

In the next two subsections we calculate those terms in  $\lambda_2$  and  $\lambda_3$  that can be found directly using Wick's theorem. After that we calculate the terms involving the wave interactions. We first consider  $\lambda_2$  in (2.3). Using the explicit form of the surface velocity (3.3) we get

$$2\lambda_{2} = \int \frac{k_{1}^{2}k_{2}^{2}d\mathbf{k}_{12}dt}{(2\pi)^{4}} \langle \psi_{1}(0)\psi_{2}(t)\rangle - 2\int \frac{d\mathbf{k}_{123}dt}{(2\pi)^{6}} |k_{1}|k_{3}^{2}(\mathbf{k}_{1} \cdot \mathbf{k}_{2} + k_{2}^{2}) \langle \psi_{1}(0)\eta_{2}(0)\psi_{3}(t)\rangle$$

$$-\int \frac{d\mathbf{k}_{1234}dt}{(2\pi)^{8}} \langle \psi_{1}(0)\eta_{2}(0)\eta_{3}(0)\psi_{4}(t)\rangle k_{4}^{2}|k_{1}| \left( |k_{1}|(\mathbf{k}_{1} \cdot \mathbf{k}_{2} + k_{2}^{2} + \mathbf{k}_{2} \cdot \mathbf{k}_{3}) + \left( |k_{1}| - 2\sqrt{k_{1}^{2} + k_{2}^{2}} \right) (\mathbf{k}_{1} \cdot \mathbf{k}_{3} + \mathbf{k}_{2} \cdot \mathbf{k}_{3} + k_{3}^{2}) \right) + \int \frac{d\mathbf{k}_{1234}dt}{(2\pi)^{8}} \langle \psi_{1}(0)\eta_{2}(0)\psi_{3}(t)\eta_{4}(t)\rangle$$

$$\times |k_{1}||k_{3}|(\mathbf{k}_{1} \cdot \mathbf{k}_{2} + k_{2}^{2})(\mathbf{k}_{3} \cdot \mathbf{k}_{4} + k_{4}^{2})$$

$$(4.5)$$

We note that the third term above, that can be decomposed by Wick's theorem, vanishes because it is supported at the zero frequency  $\omega_4$  imposing  $k_4 = 0$ . The last term can also

be analyzed using Wick's theorem. Noting that the pair-correlations are supported at the zero sum of the involved wavenumbers, we find that the last term equals

$$\int \frac{d\mathbf{k}_{1234}dt}{(2\pi)^8} \left[ \langle \psi_1(0)\psi_3(t)\rangle \langle \eta_2(0)\eta_4(t)\rangle + \langle \psi_1(0)\eta_4(t)\rangle \langle \eta_2(0)\psi_3(t)\rangle \right] 
\times |k_1||k_3|(\mathbf{k}_1 \cdot \mathbf{k}_2 + k_2^2)(\mathbf{k}_3 \cdot \mathbf{k}_4 + k_4^2).$$
(4.6)

To establish the expressions for the pair-correlation functions we pass to the normal coordinates  $a(\mathbf{k},t)$ :

$$\eta(\mathbf{k},t) = \sqrt{\frac{|k|}{2\Omega_{\mathbf{k}}}} \left[ a(\mathbf{k},t) + a^*(-\mathbf{k},t) \right], \ \psi(\mathbf{k},t) = -i\sqrt{\frac{\Omega_{\mathbf{k}}}{2|k|}} \left[ a(\mathbf{k},t) - a^*(-\mathbf{k},t) \right]. (4.7)$$

In the Gaussian approximation the pair correlation functions are given by

$$\langle a^{*}(\boldsymbol{k},t)a(\boldsymbol{k}',0)\rangle = (2\pi)^{2}\delta(\boldsymbol{k}-\boldsymbol{k}')n(\boldsymbol{k})\exp\left[i\Omega_{\boldsymbol{k}}t\right], \quad \langle a(\boldsymbol{k},t)a(\boldsymbol{k}',0)\rangle = 0, \tag{4.8}$$

$$\langle \psi(\boldsymbol{k},t)\psi(\boldsymbol{k}',0)\rangle = \frac{\Omega_{\boldsymbol{k}}(2\pi)^{2}\delta(\boldsymbol{k}+\boldsymbol{k}')}{2k}\left[n(\boldsymbol{k})\exp\left(-i\Omega_{\boldsymbol{k}}t\right) + n(-\boldsymbol{k})\exp\left(i\Omega_{-\boldsymbol{k}}t\right)\right],$$

$$\langle \eta(\boldsymbol{k},t)\eta(\boldsymbol{k}',0)\rangle = \frac{k(2\pi)^{2}\delta(\boldsymbol{k}+\boldsymbol{k}')}{2\Omega_{\boldsymbol{k}}}\left[n(\boldsymbol{k})\exp\left(-i\Omega_{\boldsymbol{k}}t\right) + n(-\boldsymbol{k})\exp\left(i\Omega_{-\boldsymbol{k}}t\right)\right],$$

$$\langle \psi(\boldsymbol{k},t)\eta(\boldsymbol{k}',0)\rangle = \frac{(2\pi)^{2}\delta(\boldsymbol{k}+\boldsymbol{k}')}{2i}\left[n(\boldsymbol{k})\exp\left(-i\Omega_{\boldsymbol{k}}t\right) - n(-\boldsymbol{k})\exp\left(i\Omega_{-\boldsymbol{k}}t\right)\right]. \tag{4.9}$$

Using the correlation functions above and noting the vanishing of the terms containing  $\delta$ -functions supported at the zero frequency, we can write (4.6) as

$$\begin{split} &\int \frac{d\boldsymbol{k}_{12}}{(2\pi)^3} \Bigg[ k_1^2 (\boldsymbol{k}_1 \cdot \boldsymbol{k}_2 + k_2^2)^2 \left( \frac{\Omega_1 k_2}{4k_1 \Omega_2} \right) \left[ n_1 n_{-2} \delta(\Omega_1 - \Omega_{-2}) + n_{-1} n_2 \delta(\Omega_{-1} - \Omega_2) \right] \\ &- \frac{1}{4} |k_1| |k_2| (\boldsymbol{k}_1 \cdot \boldsymbol{k}_2 + k_2^2) (\boldsymbol{k}_1 \cdot \boldsymbol{k}_2 + k_1^2) \left[ n_1 n_{-2} \delta(\Omega_1 - \Omega_{-2}) + n_{-1} n_2 \delta(\Omega_{-1} - \Omega_2) \right] \Bigg] = 0, \end{split}$$

where we introduced the shorthand notation:  $n(\pm \mathbf{k}_i, t) = n_{\pm i}(t)$  and  $\Omega_{\pm \mathbf{k}_i} = \Omega_{\pm i}$ . We use that for surface waves  $\Omega_{\mathbf{k}}$  is an increasing function of  $|\mathbf{k}|$ . We also use that  $\delta$ -functions imply  $\Omega_1 = \Omega_2$  and  $k_1^2 = k_2^2$ . We find that  $\lambda_2$  can be written as

$$\lambda_2 = \int \frac{d\mathbf{k}_{12}dt}{(2\pi)^4} \left[ \frac{k_1^2 k_2^2}{2} \langle \psi_1(0)\psi_2(t) \rangle - \int \frac{d\mathbf{k}_3 |k_1| k_3^2 (\mathbf{k}_1 \cdot \mathbf{k}_2 + k_2^2)}{(2\pi)^d} \langle \psi_1(0)\eta_2(0)\psi_3(t) \rangle \right]. \quad (4.10)$$

The calculation of the above terms demands the account of the interactions. We now consider the terms in  $\lambda_3$  that can be calculated by the direct use of Wick's theorem.

4.3. Contribution of the three-point correlation function

We consider  $\lambda_3$  from (2.3). Using the expression (3.3) for the velocity, we obtain

$$\lambda_{3} = -\int \frac{d\mathbf{k}_{123}}{(2\pi)^{6}} k_{1}^{2} k_{2}^{2} (\mathbf{k}_{2} \cdot \mathbf{k}_{3}) \int_{0}^{\infty} dt \int_{0}^{t} dt_{1} \langle \psi_{1}(0)\psi_{2}(t)\psi_{3}(t_{1}) \rangle$$

$$+ \int \frac{d\mathbf{k}_{1234}}{(2\pi)^{8}} \int_{0}^{\infty} dt \int_{0}^{t} dt_{1} \left[ |k_{1}| (\mathbf{k}_{1} \cdot \mathbf{k}_{2} + k_{2}^{2}) k_{3}^{2} (\mathbf{k}_{3} \cdot \mathbf{k}_{4}) \langle \psi_{1}(0)\eta_{2}(0)\psi_{3}(t)\psi_{4}(t_{1}) \rangle$$

$$+ k_{1}^{2} |k_{2}| (\mathbf{k}_{2} \cdot \mathbf{k}_{3} + k_{3}^{2}) (\mathbf{k}_{2} \cdot \mathbf{k}_{4} + \mathbf{k}_{3} \cdot \mathbf{k}_{4}) \langle \psi_{1}(0)\psi_{2}(t)\eta_{3}(t)\psi_{4}(t_{1}) \rangle$$

$$+ k_{1}^{2} k_{2}^{2} (\mathbf{k}_{2} \cdot \mathbf{k}_{4}) |k_{3}| \langle \psi_{1}(0)\psi_{2}(t)\psi_{3}(t_{1})\eta_{4}(t_{1}) \rangle \right]. \tag{4.11}$$

One can use Wick's theorem for the last three terms. Employing the identity

$$\int_0^\infty dt \int_0^t dt_1 \exp[i\omega_1 t + i\omega_2 t_1] = \frac{i\pi[\delta(\omega_1) - \delta(\omega_1 + \omega_2)]}{\omega_2},\tag{4.12}$$

one finds that part of the obtained terms contain  $\delta$ -functions supported at the zero frequency and they vanish because of the vanishing of the integrand there. Let us consider the rest of the terms. For the first of the fourth-order terms in (4.11) one finds

$$\int \frac{d\mathbf{k}_{12}}{(2\pi)^4} \frac{\Omega_1 \pi}{4k_1} (\mathbf{k}_1 \cdot \mathbf{k}_2) \left\{ |k_1|^3 (\mathbf{k}_1 \cdot \mathbf{k}_2 + k_2^2) \left[ \frac{n_{-1} n_2 \delta(\Omega_{-1} - \Omega_2)}{\Omega_2} + \frac{n_{-2} n_1 \delta(\Omega_{-2} - \Omega_1)}{\Omega_{-2}} \right] - |k_1| k_2^2 (\mathbf{k}_1 \cdot \mathbf{k}_2 + k_2^2) \left[ \frac{n_{-1} n_2 \delta(\Omega_{-1} - \Omega_2)}{\Omega_{-1}} + \frac{n_{-2} n_1 \delta(\Omega_{-2} - \Omega_1)}{\Omega_1} \right] \right\} = 0,$$

where the equality of the frequencies implies the equality of the wavelengths. Analogously, for the second Gaussian term one finds

$$\int \frac{d\mathbf{k}_{23}}{(2\pi)^4} \frac{\Omega_2 \pi}{4k_2} \left\{ |k_2|^3 (\mathbf{k}_2 \cdot \mathbf{k}_3 + k_3^2)^2 \left[ \frac{n_{-3} n_2 \delta(\Omega_{-3} - \Omega_2)}{\Omega_{-3}} + \frac{n_{-2} n_3 \delta(\Omega_{-2} - \Omega_3)}{\Omega_3} \right] - |k_2| k_3^2 (\mathbf{k}_2 \cdot \mathbf{k}_3 + k_3^2) (\mathbf{k}_2 \cdot \mathbf{k}_3 + k_2^2) \left[ \frac{n_{-3} n_2 \delta(\Omega_{-3} - \Omega_2)}{\Omega_2} + \frac{n_{-2} n_3 \delta(\Omega_{-2} - \Omega_3)}{\Omega_{-2}} \right] \right\} = 0.$$

Finally, the third Gaussian term contains only  $\delta$ -functions supported at the zero frequencies, so it also produces zero.

#### 4.4. Contribution of the interaction terms

As a result of the calculation in the previous sections, adding equations (4.10) and (4.11), one can write  $\lambda$  as a sum over terms which calculation involves the wave interactions. These terms have a special property important for the following analysis: all the terms contain a field at the zero frequency. Using Fourier representation over the frequency one finds

$$\lambda = \frac{1}{2} \int \frac{k_1^2 k_2^2 d\mathbf{k}_{12} d\omega}{(2\pi)^5} \langle \psi_1(\omega) \psi_2(\omega = 0) \rangle$$

$$- \int \frac{d\mathbf{k}_{123} d\omega_{12}}{(2\pi)^8} |k_1| k_3^2 (\mathbf{k}_1 \cdot \mathbf{k}_2 + k_2^2) \langle \psi_1(\omega_1) \eta_2(\omega_2) \psi_3(\omega = 0) \rangle$$

$$- \int \frac{d\mathbf{k}_{123} d\omega_{123}}{(2\pi)^9} k_1^2 k_2^2 (\mathbf{k}_2 \cdot \mathbf{k}_3) \langle \psi_1(\omega_1) \psi_2(\omega_2) \psi_3(\omega_3) \rangle \frac{i\pi [\delta(\omega_2) - \delta(\omega_1)]}{\omega_3}, \quad (4.13)$$

where we introduced shorthand notation  $d\omega_{ijl} = d\omega_i d\omega_j d\omega_l$ ... and used in the last term the identity (4.12) and the proportionality of the correlation function to  $\delta(\omega_1 + \omega_2 + \omega_3)$ . We observe that all the terms contain  $\psi(\mathbf{k}, \omega = 0)$ . We assume that the forcing sustaining the stationary wave turbulence vanishes at the zero frequency [note that the first term on the RHS above vanishes in the Gaussian approximation independently of this assumption, cf. Balk et al. (2004) and Vucelja et al. (2006)] and express  $\psi(\mathbf{k}, \omega = 0)$  with the help of higher order terms. Substituting the resulting expressions into the correlation functions will allow the use of Wick's decomposition.

#### 4.4.1. The expression for $\psi(\mathbf{k}, \omega = 0)$

To calculate  $\lambda$  we need to know  $\psi(\mathbf{k}, \omega = 0)$  to the third order in wave amplitude. Consider the dynamical equation on the surface elevation  $\eta$ , see Batchelor (2002); Zakharov

(1968).

$$\frac{\partial \eta}{\partial t} = \frac{\partial \phi}{\partial z} [z = \eta] - \nabla \eta \nabla \phi [z = \eta]. \tag{4.14}$$

To order  $\epsilon^2$ , the equation reads

$$\frac{\partial \eta}{\partial t} = \frac{\partial \phi_0}{\partial z} [z = 0] + \eta \frac{\partial^2 \phi_0}{\partial z^2} [z = 0] + \frac{\partial \phi_1}{\partial z} - \nabla \eta \nabla \phi_0 [z = 0] + O(\eta^3), \quad (4.15)$$

where  $\phi_0$  and  $\phi_1$  are the terms of the expansion of the potential with respect to the surface elevation, see Zakharov (1968). Using the expressions for  $\phi_i$ , see Zakharov (1968), performing the Fourier transform over the space and time coordinates, and setting the frequency  $\omega = 0$ , we find

$$0 = |k|\psi(\mathbf{k}, \omega = 0) + \int \frac{d\mathbf{k}_1 d\omega}{(2\pi)^3} k_1^2 \psi(\mathbf{k}_1, \omega) \eta(\mathbf{k} - \mathbf{k}_1, -\omega) - \int \frac{d\mathbf{k}_1 d\omega}{(2\pi)^3} |k| |k_1| \psi(\mathbf{k}_1, \omega)$$
$$\times \eta(\mathbf{k} - \mathbf{k}_1, -\omega) + \int \frac{d\mathbf{k}_1 d\omega}{(2\pi)^3} \mathbf{k}_1 \cdot (\mathbf{k} - \mathbf{k}_1) \psi(\mathbf{k}_1, \omega) \eta(\mathbf{k} - \mathbf{k}_1, -\omega) + O(\eta^3). \tag{4.16}$$

It follows that

$$\psi(\mathbf{k},\omega=0) = \int \frac{d\mathbf{k}_1 d\omega}{(2\pi)^3} \left( |k_1| - \frac{\mathbf{k}_1 \cdot \mathbf{k}}{|k|} \right) \psi(\mathbf{k}_1,\omega) \eta(\mathbf{k} - \mathbf{k}_1, -\omega) + O(\eta^3).$$
 (4.17)

The physical meaning of the above representation is that the zero frequency field arises due to the nonlinear interactions only, and in the lowest order it can be represented as a result of a single wave scattering. The above formula is sufficient to calculate the interaction terms containing the products of three fields.

#### 4.4.2. Interaction terms containing the products of three fields

To calculate the interaction terms of the third order we use (4.17), Wick's decomposition and Fourier transformed version of (4.9). The second term in (4.13) can be written with the help of (4.17) as

$$-\int \frac{d\mathbf{k}_{1234}d\omega_{123}}{(2\pi)^{11}} |k_{1}|k_{3}^{2}(\mathbf{k}_{1} \cdot \mathbf{k}_{2} + k_{2}^{2}) \left( |k_{4}| - \frac{\mathbf{k}_{4} \cdot \mathbf{k}_{3}}{|k_{3}|} \right) \langle \psi(\mathbf{k}_{1}, \omega_{1}) \eta(\mathbf{k}_{2}, \omega_{2}) \psi(\mathbf{k}_{4}, \omega_{3})$$

$$\eta(\mathbf{k}_{3} - \mathbf{k}_{4}, -\omega_{3}) \rangle = -\int \frac{d\mathbf{k}_{12}d\omega_{1}}{(2\pi)^{7}} |k_{1}| (\mathbf{k}_{1} + \mathbf{k}_{2})^{2} (\mathbf{k}_{1} \cdot \mathbf{k}_{2} + k_{2}^{2}) \left( |k_{1}| - \frac{\mathbf{k}_{1} \cdot (\mathbf{k}_{1} + \mathbf{k}_{2})}{|\mathbf{k}_{1} + \mathbf{k}_{2}|} \right)$$

$$\left( \frac{\Omega_{1}|k_{2}|}{4\Omega_{2}|k_{1}|} \right) [n_{1}\delta(\omega_{1} + \Omega_{1}) + n_{-1}\delta(\omega_{1} - \Omega_{-1})] [n_{2}\delta(-\omega_{1} + \Omega_{2}) + n_{-2}\delta(\omega_{1} + \Omega_{-2})]$$

$$+ \frac{1}{4} \int \frac{d\mathbf{k}_{12}d\omega_{1}}{(2\pi)^{7}} |k_{1}| (\mathbf{k}_{1} + \mathbf{k}_{2})^{2} (\mathbf{k}_{1} \cdot \mathbf{k}_{2} + k_{2}^{2}) \left( |k_{2}| - \frac{\mathbf{k}_{2} \cdot (\mathbf{k}_{1} + \mathbf{k}_{2})}{|\mathbf{k}_{1} + \mathbf{k}_{2}|} \right)$$

$$\times [n_{1}\delta(\omega_{1} + \Omega_{1}) - n_{-1}\delta(\omega_{1} - \Omega_{-1})] [n_{-2}\delta(\omega_{1} + \Omega_{-2}) - n_{2}\delta(\omega_{1} - \Omega_{2})], \tag{4.18}$$

where to show that the remaining contraction vanishes, one should use that  $\langle \psi(\mathbf{k}, \omega = 0) \rangle$ is representable as an integral over  $\langle \psi(\mathbf{k},t)\eta(\mathbf{k}',t)\rangle$  which in the Gaussian approximation vanishes by (4.9). Using  $\Omega_{-k} = \Omega_k$  and noting that the terms supported at  $\Omega_k = 0$ vanish, one may rewrite the above as

$$-\frac{1}{4} \int \frac{d\mathbf{k}_{12}}{(2\pi)^7} |k_1| (\mathbf{k}_1 + \mathbf{k}_2)^2 (\mathbf{k}_1 \cdot \mathbf{k}_2 + k_2^2) \left( |k_1| - \frac{\mathbf{k}_1 \cdot (\mathbf{k}_1 + \mathbf{k}_2)}{|\mathbf{k}_1 + \mathbf{k}_2|} \right) \left[ n_1 n_{-2} \delta(\Omega_1 - \Omega_{-2}) + n_{-1} n_2 \delta(\Omega_2 - \Omega_{-1}) \right] + \frac{1}{4} \int \frac{d\mathbf{k}_{12}}{(2\pi)^7} |k_1| (\mathbf{k}_1 + \mathbf{k}_2)^2 (\mathbf{k}_1 \cdot \mathbf{k}_2 + k_2^2) \left( |k_2| - \frac{\mathbf{k}_2 \cdot (\mathbf{k}_1 + \mathbf{k}_2)}{|\mathbf{k}_1 + \mathbf{k}_2|} \right)$$

8

$$[n_1 n_{-2} \delta(\Omega_1 - \Omega_{-2}) + n_{-1} n_2 \delta(\Omega_2 - \Omega_{-1})] = \frac{1}{4} \int \frac{d\mathbf{k}_{12}}{(2\pi)^7} |k_1| (\mathbf{k}_1 + \mathbf{k}_2)^2 (\mathbf{k}_1 \cdot \mathbf{k}_2 + k_2^2)$$

$$\left(\frac{(\mathbf{k}_1 - \mathbf{k}_2) \cdot (\mathbf{k}_1 + \mathbf{k}_2)}{|\mathbf{k}_1 + \mathbf{k}_2|}\right) \left[ n_1 n_{-2} \delta(\Omega_1 - \Omega_{-2}) + n_{-1} n_2 \delta(\Omega_2 - \Omega_{-1}) \right] = 0,$$
(4.19)

where the last term contains  $k_1^2 - k_2^2 = 0$ . Let us now consider the last term in (4.13) that can be written as

$$i\pi \int \frac{d\mathbf{k}_{123}d\omega_{23}}{\omega_3(2\pi)^9} k_1^2 k_2^2 \mathbf{k}_3 \cdot (\mathbf{k}_2 - \mathbf{k}_1) \langle \psi(\mathbf{k}_1, \omega = 0) \psi(\mathbf{k}_2, \omega_2) \psi(\mathbf{k}_3, \omega_3) \rangle. \tag{4.20}$$

Substituting (4.17) we find

$$i\pi \int \frac{d\mathbf{k}_{1234} d\omega_{234} k_1^2 k_2^2 \mathbf{k}_3 \cdot (\mathbf{k}_2 - \mathbf{k}_1)}{\omega_3 (2\pi)^{12}} \left( |k_4| - \frac{\mathbf{k}_4 \cdot \mathbf{k}_1}{|k_1|} \right) \langle \eta(\mathbf{k}_1 - \mathbf{k}_4, -\omega_4) \psi(\mathbf{k}_2, \omega_2) \psi(\mathbf{k}_3, \omega_3)$$

$$\times \psi(\mathbf{k}_4, \omega_4) \rangle = i\pi \int \frac{d\mathbf{k}_{23} d\omega_3}{\omega_3 (2\pi)^8} (\mathbf{k}_2 + \mathbf{k}_3)^2 k_2^2 \mathbf{k}_3 \cdot (2\mathbf{k}_2 + \mathbf{k}_3) \left( |k_3| - \frac{\mathbf{k}_3 \cdot (\mathbf{k}_2 + \mathbf{k}_3)}{|\mathbf{k}_2 + \mathbf{k}_3|} \right) \frac{\Omega_3}{4i |k_3|}$$

$$[n_2 \delta(\omega_3 - \Omega_2) - n_{-2} \delta(\omega_3 + \Omega_{-2})] [n_3 \delta(\omega_3 + \Omega_3) + n_{-3} \delta(\omega_3 - \Omega_{-3})] + i\pi \int \frac{d\mathbf{k}_{23} d\omega_3}{\omega_3 (2\pi)^8}$$

$$\times (\mathbf{k}_2 + \mathbf{k}_3)^2 k_2^2 \mathbf{k}_3 \cdot (2\mathbf{k}_2 + \mathbf{k}_3) \left( |k_2| - \frac{\mathbf{k}_2 \cdot (\mathbf{k}_2 + \mathbf{k}_3)}{|\mathbf{k}_2 + \mathbf{k}_3|} \right) \left( \frac{\Omega_2}{4i |k_2|} \right)$$

$$\times [n_2 \delta(\omega_3 - \Omega_2) + n_{-2} \delta(\omega_3 + \Omega_{-2})] [n_3 \delta(\omega_3 + \Omega_3) - n_{-3} \delta(\omega_3 - \Omega_{-3})],$$

$$(4.21)$$

where the term involving the contraction  $\langle \eta(\mathbf{k}_1 - \mathbf{k}_4, -\omega_4) \psi(\mathbf{k}_4, \omega_4) \rangle$  corresponding to  $\langle \psi(\mathbf{k}, \omega = 0) \rangle$ , vanishes as was shown in the analysis of the previous term, where we omitted the terms supported at the zero frequency. Taking the integral over  $\omega_3$ , omitting the terms supported at the zero frequency, we find

$$i\pi \int \frac{d\mathbf{k}_{23}}{(2\pi)^8} (\mathbf{k}_2 + \mathbf{k}_3)^2 k_2^2 \mathbf{k}_3 \cdot (2\mathbf{k}_2 + \mathbf{k}_3) \left( |k_3| - \frac{\mathbf{k}_3 \cdot (\mathbf{k}_2 + \mathbf{k}_3)}{|\mathbf{k}_2 + \mathbf{k}_3|} \right)$$

$$\left[ \frac{n_2 n_{-3} \delta(\Omega_2 - \Omega_{-3})}{4i |k_3|} + \frac{n_{-2} n_3 \delta(\Omega_{-2} - \Omega_3)}{4i |k_3|} \right] + i\pi \int \frac{d\mathbf{k}_{23}}{(2\pi)^8}$$

$$\times (\mathbf{k}_2 + \mathbf{k}_3)^2 k_2^2 \mathbf{k}_3 \cdot (2\mathbf{k}_2 + \mathbf{k}_3) \left( |k_2| - \frac{\mathbf{k}_2 \cdot (\mathbf{k}_2 + \mathbf{k}_3)}{|\mathbf{k}_2 + \mathbf{k}_3|} \right)$$

$$\left[ -\frac{n_{-2} n_3 \delta(\Omega_3 - \Omega_{-2})}{4i |k_2|} - \frac{n_2 n_{-3} \delta(\Omega_2 - \Omega_{-3})}{4i |k_2|} \right] = 0, \tag{4.22}$$

where zero is obtained in the same way as with the previous term. The result of this subsection is that  $\lambda$  can be written as

$$\lambda = \frac{1}{2} \int \frac{k_1^2 k_2^2 d\mathbf{k}_{12}}{(2\pi)^4} \int \frac{d\omega}{2\pi} \langle \psi(\mathbf{k}_1, \omega) \psi(\mathbf{k}_2, \omega = 0) \rangle. \tag{4.23}$$

#### 4.4.3. Interaction term containing the product of two fields

To calculate the RHS of (4.23) we note that it is sufficient to know  $\psi(\mathbf{k}_1, \omega)$  at an arbitrarily small but finite  $\omega$  where the forcing is again negligible and one use the dynamic equation (4.14) without the force. The Fourier transform of (4.15) taken now at a small but finite frequency gives

$$i\omega\eta(\mathbf{k},\omega) = |k|\psi(\mathbf{k},\omega) + \int \frac{d\mathbf{k}_1 d\omega_1}{(2\pi)^3} k_1^2 \psi_1(\omega_1) \eta(\mathbf{k} - \mathbf{k}_1, \omega - \omega_1) - \int \frac{d\mathbf{k}_1 d\omega_1}{(2\pi)^3} |k| |k_1| \psi_1(\omega)$$

$$\times \eta(\mathbf{k} - \mathbf{k}_1, \omega - \omega_1) + \int \frac{d\mathbf{k}_1 d\omega_1}{(2\pi)^3} \mathbf{k}_1 \cdot (\mathbf{k} - \mathbf{k}_1) \psi_1(\omega) \eta(\mathbf{k} - \mathbf{k}_1, \omega - \omega_1) + O(\eta^3). \tag{4.24}$$

which produces

$$\psi(\mathbf{k},\omega) = -\frac{i\omega\eta(\mathbf{k},\omega)}{|\mathbf{k}|} + \int \frac{d\mathbf{k}_1 d\omega_1}{(2\pi)^3} \left( |k_1| - \frac{\mathbf{k}_1 \cdot \mathbf{k}}{|\mathbf{k}|} \right) \psi(\mathbf{k}_1,\omega_1) \eta(\mathbf{k} - \mathbf{k}_1,\omega - \omega_1) + O(\eta^3).$$

It follows that the (4.23) can be written as

$$\lambda = \frac{1}{2} \int \frac{k_1^2 k_2^2 d\mathbf{k}_{12}}{(2\pi)^4} \int \frac{d\omega}{2\pi} \left\langle \left[ -\frac{i\omega\eta(\mathbf{k}_1,\omega)}{|\mathbf{k}_1|} + \int \frac{d\mathbf{k}_3 d\omega_1}{(2\pi)^3} \left( |\mathbf{k}_3| - \frac{\mathbf{k}_3 \cdot \mathbf{k}_1}{|\mathbf{k}_1|} \right) \psi(\mathbf{k}_3,\omega_1) \right. \\ \times \eta(\mathbf{k}_1 - \mathbf{k}_3,\omega - \omega_1) + O(\eta^3) \right] \psi(\mathbf{k}_2,\omega = 0) \right\rangle = \frac{1}{2} \int \frac{k_1^2 k_2^2 d\mathbf{k}_{12}}{(2\pi)^4} \int \frac{d\omega_1}{2\pi} \left\langle \left[ \int \frac{d\mathbf{k}_3 d\omega_3}{(2\pi)^3} \right] \right. \\ \times \left( |\mathbf{k}_3| - \frac{\mathbf{k}_3 \cdot \mathbf{k}_1}{|\mathbf{k}_1|} \right) \psi(\mathbf{k}_3,\omega_3) \eta(\mathbf{k}_1 - \mathbf{k}_3,\omega_1 - \omega_3) + O(\eta^3) \right] \psi(\mathbf{k}_2,\omega = 0) \right\rangle, \tag{4.25}$$

where we used that  $\omega \langle \eta(\mathbf{k}_1, \omega) \psi(\mathbf{k}_2, \omega = 0) \rangle \propto \omega \delta(\omega) = 0$ . Finally, using that  $\psi(\mathbf{k}, \omega = 0)$  is by itself of order  $\epsilon^2$ , see (4.17), taking the integral over  $\omega_3$  and omitting the terms supported at the zero frequency, we find  $\lambda$  to the order  $\epsilon^4$ 

$$\lambda = \frac{1}{8} \int \frac{d\mathbf{k}_{13}}{(2\pi)^3} (\mathbf{k}_1 + \mathbf{k}_3)^4 \left( |k_3| - \frac{\mathbf{k}_3 \cdot (\mathbf{k}_1 + \mathbf{k}_3)}{|\mathbf{k}_1 + \mathbf{k}_3|} \right) [n_3 n_{-1} \delta(\Omega_{-1} - \Omega_3)$$

$$+ n_{-3} n_1 \delta(\Omega_1 - \Omega_{-3})] \left( |k_3| - |k_1| - \frac{(\mathbf{k}_3 - \mathbf{k}_1) \cdot (\mathbf{k}_1 + \mathbf{k}_3)}{|\mathbf{k}_1 + \mathbf{k}_3|} \right) = 0.$$
(4.26)

The above can be easily verified by noting that the integrand can be reduced to an expression proportional to  $k_1^2 - k_3^2$ . Thus we obtain that the sum of Lyapunov exponents for weakly interacting surface waves is zero in the fourth order in the wave amplitude.

# 5. Discussion

We have shown that the sum of the Lyapunov exponents for surface wave turbulence vanishes in the fourth order in wave amplitude. Using the approximate Gaussianity of the statistics, it is easy to see that the leading order term in  $\lambda$  is of the sixth order in wave-amplitude, or higher. Therefore we have derived that  $\lambda \lesssim \Omega_{\bf k} \epsilon^6$ . For waves with a typical period of the order of seconds and with not too small  $\epsilon=0.1$ , we find that the time-scale  $1/\Omega_{\bf k}\epsilon^6$  is of the order of weeks. Thus, even if there is no degeneracy in the sixth order and  $\lambda \sim \Omega_{\bf k}\epsilon^6$ , the formation of the inhomogeneities would occur at the time-scale of weeks and larger. It is unlikely that a low-amplitude wave turbulence would persist for such time. Thus we expect the turbulence of small-amplitude surface waves to have negligible effect on the formation of the floaters inhomogeneities in realistic situations. Let us stress that this weak compressibility of surface waves holds in the sense of the long-time action of the flow on the particles, while the characteristic value of the ratio of the surface flow divergence to the curl is of order one.

Let us consider some estimates for the Lyapunov exponents of the floaters velocity field. For the non-interacting Gaussian surface waves,  $\lambda_1$  is non-zero in the fourth order in wave amplitude, see Balk *et al.* (2004), while  $\lambda$  is non-zero in the sixth order, Balk *et al.* (2004); Stepanov (2006). The nonlinear wave interactions and the nonlinearity of the velocity-amplitude relation add to  $\lambda_1$  additional terms of the fourth order in the wave amplitude and higher. It is highly implausible that these terms produce exact cancelation

of  $\lambda_1$  in the fourth order - such cancelation would depend on the precise form of both the interactions and the velocity-amplitude relation. Moreover, positive Lyapunov exponent and Lagrangian chaos hold even for rather simple Eulerian flows, see e.g. Bohr & Hansen (1996), so no degeneracy for  $\lambda_1$  is expected that would lead to an exact cancellation in the fourth order. Therefore we expect that  $\lambda_1$  for surface waves is of the fourth order in the wave amplitude. Similarly, we expect no exact cancelation of the non-vanishing Gaussian terms for  $\lambda$ , in the sixth order in the wave amplitude, see also below. Then we have the following order of magnitude estimates:  $\lambda_1 \propto \Omega_k \epsilon^4$  and  $\lambda \propto \Omega_k \epsilon^6$ . It follows that surface wave turbulence is also weakly compressible in the sense of the ratio  $\lambda/\lambda_1 \ll 1$ , which is of the second order in the wave amplitude. The Kaplan-Yorke dimension of the particles attractor on the surface is close to the space dimension:  $D_{KY} \approx 2 - \lambda/\lambda_1 \approx 2$ . For dimensionless exponents  $\tilde{\lambda}_1 \equiv \lambda_1/\Omega_k$  and  $\tilde{\lambda} \equiv \lambda/\Omega_k$  we find the order of magnitude relation  $\tilde{\lambda} \sim \tilde{\lambda}_1^{3/2}$  holding at small wave amplitudes. The above estimates are supported by numerical simulations performed by Umeki (1992) in a similar problem with standing surface waves. It was shown there that both  $\lambda \ll \lambda_1$  and  $2 - D_{KY} \ll 1$  hold. Moreover, the numerical values of the dimensionless exponents  $\tilde{\lambda}$ ,  $\tilde{\lambda}_1$  found there, can be easily seen to be in agreement with the relation  $\tilde{\lambda} \sim \tilde{\lambda}_1^{3/2}$ . The box-counting dimension of the particles attractor on the surface was found very close to  $D_{KY}$ , which we expect to hold for wave turbulence as well. The detailed calculations of the exact expressions for  $\lambda_1$  and  $\lambda$  are subjects for future work.

We believe that our conclusion on the weak clustering in surface waves with a small amplitude is an important step in identifying possible reasons for the clusters of the floaters observed on liquid surfaces ubiquitously. Our results suggest that other mechanisms need to be explored, such as the wave breaking and Langmuir circulation, see e.g. Thorpe (2005), the interplay of waves and currents, see Vucelja *et al.* (2006), and others.

We are indebted to G. Falkovich for constant help and useful discussions. We are grateful to M. G. Stepanov for helpful discussions. M. V. thanks S. Nazarenko for a useful remark. This work was supported by the Israeli Science Foundation.

#### REFERENCES

- Balk, A. M., Falkovich, G. & Stepanov, M. G. 2004 Growth of density inhomogeneities in a flow of wave turbulence. *Phys. Rev. Lett.* **92**, 244504.
- Balkovsky, E., Falkovich, G. & Fouxon, A. 2001 Intermittent distribution of inertial particles in turbulent flows. *Phys. Rev. Lett.* **86**, 2790.
- Bandi, M. M., Goldburg, W. I. & Jr, G. R. Cressman 2006 Measurement of entropy production rate in compressible turbulence. *Europhys. Lett.* **76**, 595–601.
- BATCHELOR, G. K. 2002 An Introduction to Fluid Dynamics. Cambridge: Cambridge University. BEC, J., GAWEDZKI, K. & HORVAI, P. 2004 Multifractal clustering in compressible flows. Phys. Rev. Lett. 92, 224501.
- BOFFETTA, G., DAVOUDI, J., ECKHARDT, B. & SCHUMACHER, J. 2006a Lagrangian tracers on a surface flow: The role of time correlations. *Phus. Rev. Lett.* **93**, 134501.
- BOFFETTA, G., DAVOUDI, J. & LILLO, F. DE 2006b Multifractal clustering of passive tracers on a surface flow. *Europhys. Lett.* **74**, 62.
- Bohr, T. & Hansen, J.L. 1996 Chaotic particle motion under linear surface waves. *Chaos* 6, 554–563.
- Cressman, J. R., Davoudi, J., Goldburg, W. & Schumacher, J. 2004 Eulerian and lagrangian studies in surface flow turbulence. *New J. Physics* **6**, 53.
- CRESSMAN, J. R. & GOLDBURG, W. 2003 Compressible flow: Turbulence at the surface. *J. Stat. Phys.* **113**, 875–883.
- Denissenko, P., Falkovich, G. & Lukaschuk, S. 2006 How waves affect the distribution of particles that float on a liquid surface. *Phys. Rev. Lett.* **97**, 244501.

DORFMAN, J. 1999 Introduction to Chaos in Nonequilibrium Statistical Mechanics. Cambridge Univ. Press.

ECKHARDT, B. & SCHUMACHER, J. 2001 Turbulence and passive scalar transport in a free-slip surface. *Phys. Rev. E* **64**, 016314.

FALKOVICH, G. & FOUXON, A. 2003 nlin.cd/0312033.

Falkovich, G. & Fouxon, A. 2004 Entropy production and extraction in dynamical systems and turbulence. *New J. Physics* **6**.

FALKOVICH, G., GAWĘDZKI, K. & VERGASSOLA, M. 2001 Particles and fields in fluid turbulence. Rev. Mod. Phys. 73, 913.

Nameson, A., Antonsen, T. & Ott, E. 1996 Power law wave number spectra of fractal particle distributions advected by flowing fluids. *Phys. Fluids* 8, 2426–2434.

Ott, E. 2002 Chaos in Dynamical Systems. Cambridge: Cambridge University Press.

RAMSHANKAR, R., BERLIN, D. & GOLLUB, J. 1990 Transport by capillary waves. part i. particle trajectories. *Phys. Fluids A* 2, 1955–1965.

Ruelle, D. 1996 Positivity of entropy production in nonequilibrium statistical mechanics. J. Stat. Phys. 85, 1–23.

Ruelle, D. 1997 Positivity of entropy production in the presence of a random thermostat. *J. Stat. Phys.* **86**, 935–990.

Ruelle, D. 1999 Smooth dynamics and new theoretical ideas in nonequilibrium statistical mechanics. J. Stat. Phys. 95, 393–468.

Schroder et al, E. 1996 Relative particle motion in capillary waves. *Phys. Rev. Lett.* **76**, 4717.

Sommerer, J. C. 1996 Experimental evidence for power-law wave number spectra of fractal tracer distributions in a complicated surface flow. *Phys. Fluids* 8, 2441–2446.

Sommerer, J. C. & Ott, E. 1993 Particles floating on a moving fluid: A dynamically comprehensible physical fracta. *Science* **259**, 335–339.

Stepanov, M. G. 2006 private communication.

THORPE, S. A. 2005 The Turbulent Ocean. Cambridge Univ. Press.

UMEKI, M. 1992 Lagrangian motion of fluid particles induced by three-dimensional standing surface waves. Phys. Fluids A 4, 1968–1978.

Vucelja, M., Falkovich, G. & Fouxon, I. 2006 Clustering of matter in waves and currents. submitted to Phys. Rev. Lett. .

Yu, L., Ott, E. & Chen, Q. 1991 Fractal distribution of floaters on a fluid surface and the transition to chaos for random maps. *Physica D* 53, 102–124.

Zakharov, V. E. 1966 PhD thesis, in Russian, Institute for Nuclear Physics (Physics and Mathematics), Novosibirsk.

Zakharov, V. E. 1968 Stability of periodic waves of finite amplitude on the surface of a deep fluid (note a sign misprint in (1.8)). J. Appl. Mech. Tech. Phys. 9, 190–194.

ZAKHAROV, V. E., L'VOV, V. & FALKOVICH, G. 1992 Kolmogorov spectra of turbulence. Berlin: Springer-Verlag.