

# Creation and Decay of $\eta$ -Mesic Nuclei

G.A. Sokol<sup>1</sup>, T.A. Aibergenov, A.V. Koltsov, A.V. Kravtsov,  
 Yu.I. Krutov, A.I. L'vov, L.N. Pavlyuchenko,  
 V.P. Pavlyuchenko, S.S. Sidorin

*P.N. Lebedev Physical Institute, Leninsky Prospect 53, Moscow 117924, Russia*

## Abstract.

First experimental results on photoproduction of  $\eta$ -mesic nuclei are analyzed. In an experiment performed at the 1 GeV electron synchrotron of the Lebedev Physical Institute, correlated  $\pi^+n$  pairs arising from the reaction

$$\gamma + {}^{12}\text{C} \rightarrow N + \eta(A-1) \rightarrow N + \pi^+ + n + (A-2)$$

and flying transversely to the photon beam have been observed. When the photon energy exceeds the  $\eta$ -meson production threshold, a distribution of the  $\pi^+n$  pairs over their total energy is found to have a peak in the subthreshold region of the internal-conversion process  $\eta p \rightarrow \pi^+n$  which signals about formation of  $\eta$ -mesic nuclei.


The idea that a bound state of the  $\eta$ -meson and a nucleus (the so-called  $\eta$ -mesic nucleus) can exist in Nature was put forward long ago by Peng [1] who relied on the first estimates of the  $\eta N$  scattering length  $a_{\eta N}$  obtained by Bhalerao and Liu [2]. Owing to  $\text{Re } a_{\eta N} > 0$ , an average attractive potential exists between slow  $\eta$  and nucleons. This can result in binding  $\eta A$  systems, provided the life time of  $\eta$  in nuclei is long enough [3]. Modern calculations [4,5] predict a rather strong  $\eta N$  attraction which is sufficient for binding  $\eta$  in all nuclei with  $A \geq 4$ .

The very first experiments on searching for the  $\eta$ -mesic nuclei performed at BNL [6] and LAMPF [7] gave negative results. Meantime, studies of the reactions  $p(d, {}^3\text{He})\eta$  [8,9],  ${}^{18}\text{O}(\pi^+, \pi^-){}^{18}\text{Ne}$  [10], and  $\vec{d}(d, {}^4\text{He})\eta$  [11] suggest that a quasi-bound  $\eta A$  state is formed in these reactions [12,13].

In the present work we report on first results concerning formation of  $\eta$ -mesic nuclei in photoreactions. A very efficient trigger for searching for  $\eta$ -mesic nuclei [15] consists in detecting decay products of the  $\eta$ -mesic nucleus, viz. a  $\pi N$  pair produced in the reaction  $\eta N \rightarrow \pi N$  inside the nucleus. Here  $\eta$  itself is produced at an earlier stage, in the reaction  $\gamma N \rightarrow \eta N$  in our case. Both these reactions are mediated by the  $S_{11}(1535)$  resonance which affects also a propagation of the

<sup>1</sup>) Talk given at CIPANP-2000 (May 22-28, 2000, Quebec). E-mail: gsokol@x4u.lebedev.ru

intermediate  $\eta$  in the medium (via multiple  $\eta N$  rescattering) and leads to capturing slow  $\eta$  into a bound state (Fig. 1). Formation of the bound state of the  $\eta$  and the nucleus becomes possible when the momentum of the produced  $\eta$  is small (typically less than 150 MeV/c). This requirement suggests photon energies  $E_\gamma = 650\text{--}850$  MeV as most suitable for creating  $\eta$ -nuclei.




**FIGURE 1.** Mechanism of creation and decay of  $\eta$ -mesic nuclei.

$\pi N$  pairs emerging from  $\eta$ -mesic nucleus decays have an opening angle  $\langle\theta_{\pi N}\rangle = 180^\circ$  and specific kinetic energies of their components (though smeared by the Fermi motion),  $\langle E_\pi\rangle \simeq 300$  MeV,  $\langle E_n\rangle \simeq 100$  MeV. Among four possible isotopic combinations  $\pi^+n$ ,  $\pi^-p$ ,  $\pi^0n$ ,  $\pi^0p$  the first one is quite suitable for measuring energies of the particles.

Accordingly, in an experiment performed at the 1 GeV electron synchrotron of the Lebedev Physical Institute, correlated  $\pi^+n$  pairs arising from the reaction

$$\gamma + {}^{12}\text{C} \rightarrow N + \eta(A-1) \rightarrow N + \pi^+ + n + (A-2) \quad (1)$$


have been searched for. An experimental setup (Fig. 2) consisted of a carbon target  $\varnothing 4\text{ cm} \times 4\text{ cm}$  and two time-of-flight scintillator spectrometers having a time resolution of  $\delta\tau \simeq 0.1$  ns. A plastic anticounter A of charged particles (of the 90% efficiency), placed in front of the neutron detectors, and  $dE/dx$  layers, placed between start and stop detectors in the pion spectrometer, were used for a better identification of particles.



**FIGURE 2.** Layout of the experimental setup. Shown also time-of-flight spectra in the  $\pi$  (left) and  $n$  (right) spectrometers.

Strategy of measurements was as follows. Two bremsstrahlung-beam energies were used,  $E_{\gamma\text{max}} = 650$  MeV and 850 MeV, i.e., well below and well above  $\eta$


production threshold on the free nucleon which is 707 MeV. The first, “calibration” run was performed at 650 MeV with the spectrometers positioned at angles  $\theta_n = \theta_\pi = 50^\circ$  around the beam. In that run, this was a quasi-free photoproduction  $\gamma p \rightarrow \pi^+ n$  which dominated the observed yield of the  $\pi^+ n$  pairs. Then, at the same “low” energy 650 MeV, the spectrometers were positioned at  $\theta_n = \theta_\pi = 90^\circ$  (the “background” run). In such a kinematics, the quasi-free production did not contribute and the observed count was presumably dominated by double-pion production. At last, the third run (the “effect+background” run) was performed at the same  $90^\circ/90^\circ$  position, however with the higher photon beam energy of 850 MeV, at which  $\eta$  mesons are produced too.



**FIGURE 3.** Raw  $\pi^+ n$  event distributions over the particle velocities  $\beta_n$  and  $\beta_\pi$  for the “calibration” run (a), “background” run (b), and “effect+background” run (c).

In accordance with measured velocities of particles detected by the spectrometers, all candidates to the  $\pi^+ n$  events were separated into three classes: fast-fast (FF), fast-slow (FS), and slow-slow (SS) events. The FF events mostly correspond to  $\pi^0 \pi^0$  production which results in hitting detectors by photons or  $e^+/e^-$ . The FS events mostly emerge from the  $\pi^+ n$  pairs. Comparing yields and time spectra in these runs (and, in particular, using the SS events for extrapolating and subtracting a background), we have found a clear excess of the FS events which appeared when the photon energy exceeded  $\eta$  production threshold. The total cross section


of photoproduction of such excess pairs, averaged over the photon energy range 650–850 MeV, was found to be about  $\sigma_{\text{tot}}(\pi^+n) \simeq 10 \mu\text{b}$ . See Ref. [14] for more details. In the present work a further analysis of the excess FS events is done and their energy characteristics are determined.




**FIGURE 4.** Corrected two-dimensional distributions over the velocities  $\beta$  of the  $\pi^+n$  events with the end-point energy of the bremsstrahlung spectrum  $E_{\gamma\text{max}} = 850$  and  $650$  MeV.

In order to find kinetic energies of the neutron and pion, the velocities  $\beta_i = L_i/ct_i$  of both the particles have to be determined. They are subject to fluctuations stemming from errors  $\delta t_i$  and  $\delta L_i$  in the time-of-flight  $t_i$  and the flight base  $L_i$ . Such fluctuations are clearly seen in the case of the ultra-relativistic FF events which have experimentally observed velocities close but not equal to 1 (see Fig. 3). Therefore, an experimental  $\beta$ -resolution of the setup can be directly inferred from the FF events. Then, using this information and applying an inverse-problem statistical method described in Ref. [16], one can unfold the experimental spectrum, obtain a smooth velocity distribution in the physical region  $\beta_i < 1$  (Fig. 4), and eventually find a distribution of the particle's kinetic energies  $E_i = M_i[(1 - \beta_i^2)^{-1/2} - 1]$ . Finding  $E_i$ , we introduced corrections related with average energy losses of particles in absorbers and in the detector matter. It is worth to say that the number of the  $\pi^+n$  FS events visibly increases when the photon beam energy becomes sufficient for producing  $\eta$  mesons.

Of the most interest is the distribution of the  $\pi^+n$  events over their total energy  $E_{\text{tot}} = E_n + E_\pi$ , because creation and decay of  $\eta$ -mesic nuclei is expected to produce a relatively narrow peak in  $E_{\text{tot}}$  of the width  $\sim 50\text{--}70$  MeV (see, e.g., [14,17]). Such a peak was indeed observed: see Fig. 5, in which an excess of the FS events appears when the photon energy exceeds the  $\eta$ -production threshold. Subtracting a smooth background, we have found a 1-dimensional energy distribution of the  $\pi^+n$  events presumably coming from (bound)  $\eta$  decaying in the nucleus, see Fig. 6. The experimental width of this distribution is about 100 MeV, including the apparatus resolution. Its center lies by  $\Delta E = 40$  MeV below the energy excess  $m_\eta - m_\pi = 408$  MeV in the reaction  $\eta N \rightarrow \pi N$ , and it is well below the position of the  $S_{11}(1535)$  resonance too. Up to effects of binding of protons annihilated in



**FIGURE 5.** Distribution over the total kinetic energy of the  $\pi^+n$  pairs for the “effect+background” run (the left panel) and for the “background” run (the right panel) obtained after unfolding the raw spectra.




**FIGURE 6.** Distribution over the total kinetic energy of the  $\pi^+n$  pairs after subtraction of the background. Arrows indicate threshold in the reaction  $\eta N \rightarrow \pi N$ , i.e. 408 MeV, and the weighted center of the histogram. For a comparison, a product of free-particle cross sections of  $\gamma N \rightarrow \eta N$  and  $\eta N \rightarrow \pi N$  [5] is shown with the dashed line (in arbitrary units).

the decay subprocess  $\eta p \rightarrow \pi^+n$ , the value  $\Delta E$  characterizes the binding energy of  $\eta$  in the nucleus. The width of that peak is determined both by the width of the  $\eta$ -bound state and by the Fermi motion.

Whereas the fixed opening angle  $\theta_{\pi n} = 180^\circ$  chosen in the kinematics with  $\theta_n = \theta_\pi = 90^\circ$  selects  $\pi^+n$  pairs carrying a low total momentum in the direction of the photon beam, an independent check of the transverse momentum  $p_\perp = p_\pi - p_n$  is meaningful. The corresponding distribution is shown in Fig. 7. On the top of a background, there is a narrower peak in  $p_\perp$  having a width compatible with the Fermi momentum of nucleons in the nucleus.

In conclusion, an excess of correlated  $\pi^+n$  pairs with the opening angle  $\langle \theta_{\pi N} \rangle = 180^\circ$  has been experimentally observed when the energy of photons exceeded the  $\eta$ -production threshold. A distribution of the pairs over their total kinetic energy was found to have a peak lying below threshold of the elementary process  $\pi N \rightarrow \eta N$ .



**FIGURE 7.** Distribution over the total transverse momentum  $p_{\perp}$  of  $\pi^{+}n$  pairs for the “effect+background” run (the left panel) and for the “calibration” run (the right panel).

A narrow peak is also found in the pair’s distribution over their total transverse momentum. All that suggests that these  $\pi^{+}n$  pairs arise from creation and decay of captured bound  $\eta$  in the nucleus, i.e., they arise through the stage of formation of an  $\eta$ -mesic nucleus.

The work was supported by RFBR grant 99-02-18224.

## REFERENCES

1. J.C. Peng, AIP Conference Proceedings **133**, 255 (1985).
2. R.S. Bhalerao and L.C. Liu, *Phys.Rev.Lett.* **54**, 865 (1985).
3. L.C. Liu and Q. Haider, *Phys. Rev. C* **34** (1986) 1845.
4. M. Batinić et al., *Phys. Rev. C* **57** (1998) 1004.
5. A.M. Green and S. Wycech, *Phys. Rev. C* **55** (1997) R2167.
6. R.E. Chrien et al., *Phys. Rev. Lett.* **60** (1988) 2595.
7. B.J. Lieb and L.C. Liu, LAMPF Progress Report, LA-11670-PR (1988).
8. J. Berger et al., *Phys. Rev. Lett.* **61** (1988) 919.
9. B. Mayer et al., *Phys. Rev. C* **53** (1996) 2068.
10. J.D. Johnson et al., *Phys. Rev. C* **47** (1993) 2571.
11. N. Willis et al., *Phys. Lett.* **B 406** (1997) 14.
12. C. Wilkin, *Phys. Rev. C* **47** (1993) R938.
13. L. Kondratyuk et al., Proc. Int. Conf. “Mesons and Nuclei at Intermediate Energies”, Dubna 1994 (World Scientific, Singapore, Eds. M.Kh. Khanhasayev and Zh.B. Kurmanov), p. 714.
14. G.A. Sokol et al., *Fizika B* **8** (1998) 81.
15. G.A. Sokol and V.A. Tryasuchev, *Kratk. Soobsh. Fiz.* [Sov. Phys. – Lebedev Institute Reports] **4** (1991) 23.
16. V.P. Pavlyuchenko, *Vopr. Atom. Nauki i Tekhn, ser. Tekh. Fiz. Eksp.* **1/13** (1983) 39.
17. A.I. L’vov, nucl-th/9810054 and Proc. 7th Int. Conf. “Mesons and Light Nuclei”, Pruhonice, Prague, Czech Republic, 1998 (World Scientific, eds. J. Adam et al.), p. 469.