

Near threshold electroproduction of the ω meson at $Q^2 \approx 0.5 \text{ GeV}^2$

P. Ambrozewicz,^{1,2,*} J. Mitchell,² J. Dunne,^{2,†} P. Markowitz,^{3,2} C. J. Martoff,¹ J. Reinhold,^{4,*} B. Zeidman,⁴ D. J. Abbott,² A. Ahmidouch,^{5,6,‡} C. S. Armstrong,⁷ J. Arrington,⁴ K. A. Assamagan,^{8,5} K. Bailey,⁴ O. K. Baker,^{5,2} S. Beedoe,⁹ H. Breuer,¹⁰ R. Carlini,² J. Cha,⁵ G. Collins,¹⁰ C. Cothran,¹¹ W. J. Cummings,⁴ S. Danagoulian,^{9,2} D. Day,¹¹ F. Duncan,¹⁰ D. Dutta,^{12,§} T. Eden,⁵ R. Ent,² L. Ewell,^{10,¶} H. T. Fortune,¹³ H. Gao,⁴ D. F. Geesaman,⁴ P. Gueye,⁵ K. K. Gustafsson,^{10,**} J.-O. Hansen,^{4,††} W. Hinton,⁵ C. E. Keppel,^{5,2} A. Klein,¹⁴ D. Koltenuk,^{13,‡‡} D. J. Mack,² R. Madey,^{5,6} D. G. Meekins,^{7,††} H. Mkrtchyan,¹⁵ R. M. Mohring,^{10,§§} S. K. Mtingwa,⁹ G. Niculescu,^{5,¶¶} I. Niculescu,⁵ T. G. O'Neill,^{4,***} D. Potterveld,⁴ J. W. Price,¹⁶ B. A. Raue,³ P. Roos,¹⁰ G. Savage,⁵ R. Sawafta,^{9,2} R. E. Segel,¹² S. Stepanyan,¹⁷ V. Tadevosyan,¹⁷ L. Tang,^{5,2} B. P. Terburg,¹⁸ S. Wood,² C. Yan,² and B. Zihlmann^{11,†††}

¹Temple University, Philadelphia, Pennsylvania 19122

²Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606

³Florida International University, Miami, Florida 33199

⁴Argonne National Laboratory, Argonne, Illinois 60439

⁵Hampton University, Hampton, Virginia 23668

⁶Kent State University, Kent, Ohio 44242

⁷College of William and Mary, Williamsburg, Virginia 23187

⁸Brookhaven National Laboratory, Upton, New York 11973

⁹North Carolina A & T State University, Greensboro, North Carolina 27411

¹⁰University of Maryland, College Park, Maryland 20742

¹¹University of Virginia, Charlottesville, Virginia

¹²Northwestern University, Evanston, Illinois 60201

¹³University of Pennsylvania, Philadelphia, Pennsylvania 19104

¹⁴Old Dominion University, Norfolk, Virginia

¹⁵Yerevan Physics Institute, Yerevan, Armenia 375036

¹⁶Rensselaer Polytechnic Institute, Troy, New York 12180

¹⁷Yerevan Physics Institute, Yerevan, Armenia 375306

¹⁸University of Illinois, Champaign-Urbana, Illinois 61801

(Dated: November 21, 2018)

Electroproduction of the ω meson was investigated in the $^1\text{H}(e, e'p)\omega$ reaction. The measurement was performed at a 4-momentum transfer $Q^2 \approx 0.5 \text{ GeV}^2$. Angular distributions of the virtual photon-proton center-of-momentum cross sections have been extracted over the full angular range. These distributions exhibit a strong enhancement over t -channel parity exchange processes in the backward direction. According to a newly developed electroproduction model, this enhancement provides significant evidence of resonance formation in the $\gamma^*p \rightarrow \omega p$ reaction channel.

PACS numbers: 25.30.Rw, 25.30.Dh, 13.60.Le

I. INTRODUCTION

There are only few measurements of the cross section for electroproduction of light vector mesons in the near threshold regime [1, 2]. These experiments, carried out at DESY, despite suffering from very low statistics revealed that different mechanisms contribute to production of the ρ^0 and ω mesons in this region. The data for both the energy dependence and angular distribution of ρ^0 meson electroproduction were found to be consistent

*presently at Florida International University, Miami, Florida 33199

†presently at Mississippi State University, Starkville, Mississippi 39762

‡presently at North Carolina A & T State University, Greensboro, North Carolina 27411

§presently at Duke University, Durham, North Carolina 27708

¶presently at University of Michigan Medical Center, Ann Arbor, Michigan 48109

**presently at University of Helsinki, Helsinki, Finland

††presently at Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606

‡‡presently at MIT Lincoln Laboratory, Lexington, Massachusetts 02420

§§presently at Millennium Cell, Inc., Eatontown, NJ 07724

¶¶presently at James Madison University, Harrisonburg, Virginia

22807

***presently at Sun Microsystems, Inc. Mountain View, California 94043

†††presently at University of Gent, B-9000 Gent, Belgium

with a vector meson dominance (VMD) model described by t -channel particle exchange with natural or unnatural parity. This production mechanism is represented by the t -channel diagrams of Fig. 1. Diffractive scattering, interpreted as t -channel Pomeron exchange in the language of Regge theory, is the dominant process in the natural parity exchange mechanism above the traditional resonance region. Near the ω production threshold, because of the appreciable relative decay width $\Gamma_{\omega \rightarrow \pi^0 \gamma} (\sim 8\%)$, t -channel unnatural parity exchange, mediated by the exchange of the π^0 meson, can make significant, even dominant, contributions to ω electroproduction.

A VMD-based model [3], which includes both of these




FIG. 1: Vector meson dominance t -channel contributions: (a) diffractive scattering - natural parity exchange, (b) π^0 exchange - unnatural parity exchange.

mechanisms fails, however, to reproduce the electroproduction data near threshold [2]. It was found that the strength of the total cross section at threshold is much larger than that predicted for the t -channel exchange contributions. This enhancement was associated with the non-peripheral component of the total cross section corresponding to large t or, equivalently, backward scattering angles. Theoretical models based on t -channel exchange predict a strongly forward peaked angular distribution of the cross section that monotonically decreases with increasing angle. The results presented in this paper substantially differ from this prediction. Such discrepancies were suggested by other earlier measurements which, as in Ref. [2], found disagreements in the energy dependence of the total cross section [4, 5]. More recent theoretical models address this by including s -channel and u -channel contributions to compensate for the additional strength at threshold.

The data for the present analysis were acquired in Hall C at the Thomas Jefferson National Accelerator Facility (Jefferson Lab) during an experiment designed to study electroproduction of strangeness via $^1\text{H}(e, e'K^+)\Lambda(\Sigma)$ [6]. Part of the background in the kaon electroproduction experiment were moderately inelastic $e'p$ events rejected in the analysis by kaon particle identification. These $e'p$ events, analyzed in the present work, provide the largest, to date, available data set on ω meson electroproduction.

This work reports on a measurement of the differential cross section for electroproduction of ω mesons ob-

served in the $^1\text{H}(e, e'p)\omega$ reaction near threshold at four-momentum transfer $Q^2 \approx 0.5 \text{ GeV}^2$. The detailed analysis can be found in Ref. [7].

II. EXPERIMENT

The experiment was conducted in Hall C at Jefferson Lab. The layout of the instrumentation is indicated in Fig. 2. Data were taken using 3.245 GeV electrons impinging on a 4.36-cm long target cell [8, 9]. Liquid hydrogen circulating through the cell was cooled in a heat exchanger by 15 K gaseous helium and kept at a temperature of $(19 \pm 0.2) \text{ K}$ and a pressure of 24 psia.

The experiment used the High Momentum Spectrom-




FIG. 2: Top view of Hall C. **Q** and **D** denote quadrupole and dipole magnets respectively.

eter (HMS) to detect scattered electrons. Its geometrical acceptance of $\sim 6.8 \text{ msr}$ was defined by an octagonal aperture in a 6.35-cm thick tungsten collimator. Before being detected, the electrons traversed the magnetic field of four superconducting magnets; three quadrupoles followed by a dipole. A pair of drift chambers at the focal plane of the spectrometer was used to determine the electron momentum while a threshold gas Čerenkov detector and Pb-glass calorimeter provided particle identification at both hardware (trigger) and software levels. Arrays of segmented scintillator hodoscopes were used to form the trigger and provide time-of-flight (ToF) measurements. All of the ω data were taken with an HMS spectrometer central angle of 17.20° and a central momentum of 1.723 GeV. This choice defined the virtual photon flux centered at 17.67° from the beam direction, and the four-momentum transfer $Q^2 \approx 0.5 \text{ GeV}^2$.

TABLE I: Central values of the hadron arm momentum p_0 , angular setting θ_0 , as well as the corresponding virtual photon proton separation $\theta_{\gamma p}$, and virtual photon ω meson CM angle θ^* .

| p_0 (GeV) | θ_0 (deg) | $\theta_{\gamma p}$ (deg) | θ^* (deg) |
|-------------|------------------|---------------------------|------------------|
| 1.077 | 17.67 | 0.00 | 180 |
| | 22.00 | 4.33 | 155 |
| | 26.50 | 8.78 | 135 |
| | 31.00 | 13.3 | 115 |
| 0.929 | 17.67 | 0.00 | 180 |
| | 22.00 | 4.33 | 130 |
| | 26.50 | 8.78 | 110 |
| | 31.00 | 13.3 | 95 |
| | 35.00 | 17.3 | 85 |
| 0.650 | 17.67 | 0.00 | 0 |
| | 22.00 | 4.33 | 15 |
| | 26.50 | 8.78 | 25 |

The Short Orbit Spectrometer (SOS) was set to detect positively charged particles (π^+ , K^+ or p) and served as the hadron arm in the experiment. An octagonal aperture in a 6.35-cm thick tungsten collimator defined the SOS solid angle acceptance to be roughly 7.5 msr. Hadrons were detected after passing through the magnetic field of three resistive magnets; a quadrupole and two dipoles with opposite bending directions. A detector package similar to that of the HMS allowed for momentum determination (multi-wire drift chambers) and particle identification (segmented hodoscope arrays and Čerenkov detectors). Having fixed the electron arm position and momentum, the angular θ_0 and momentum p_0 setting of the hadron arm was varied to access different scattering angles θ^* in the hadron (γ^*p) center-of-momentum (CM) system. These spectrometer settings, which corresponded to increasing virtual photon proton angular separation $\theta_{\gamma p}$ in the lab, allowed complete coverage for the ω scattering angles θ^* with respect to the virtual photon direction in the CM frame, particularly backward of 60° . The data taken for the forward angles suffered from very low statistics. All the settings are presented in Table I. Figure 3 shows the full kinematic coverage of the data set in conjunction with the available acceptance. The closed curves in this figure are contours of constant invariant mass W and the radial lines are contours of constant scattering angle θ^* in the hadron CM frame. Open circles are at 20 MeV and 5 degree increments, respectively. The plot was generated for the ω mass, 0.782 GeV, and $Q^2 = 0.5 \text{ GeV}^2$. It is evident from this plot that a finite acceptance in proton lab momentum can produce cuts in which the range of accepted W is a strong function of θ^* . These correlations were accounted for in the extraction of the differential cross sections from the data.




FIG. 3: Total kinematic coverage. Straight lines define the acceptance of the experimental apparatus for all the kinematic settings.

III. DATA ANALYSIS

Inelastic electron-proton final states were relatively easy to identify. Electrons were well separated from pions at the trigger level and final purification was achieved by using cuts on detector responses from the HMS gas Čerenkov detector and the Pb-glass calorimeter. Protons were selected using two types of scintillator timing information, time-of-flight (ToF) and coincidence time. In the SOS, the ToF was measured between two pairs of segmented hodoscope arrays separated by 1.76 m. In addition, relative coincidence time was measured between the hadron and electron arm scintillator arrays. The top plots in Fig. 4 show typical distributions of ToF velocity, β_{ToF} , and coincidence time. The relatively large momentum acceptance, $\pm 20\%$ of the central setting (Table I), resulted in a variation of velocity with momentum (manifested as an asymmetry in the proton β_{ToF} distribution, see Fig. 4 top left). This, together with the associated pathlength variations, required corrections to the coincidence time to account for deviations from the central trajectory. The corrected coincidence time distribution (Fig. 4 top right) clearly shows the 2 ns radio frequency (RF) microstructure of the electron beam. This structure was essential in the proton identification and accidental background removal. Real coincidence events, $e'p$ pairs coming from the same interaction point, form a prominent peak at -4.5 ns. The remaining peaks are formed by random coincidences.

The final sample of protons was selected by requiring the corrected coincidence time to be within the three RF peaks centered on the true coincidence peak and by employing a cut, for improved selectivity, on the difference between ToF velocity β_{ToF} and the velocity calculated using the measured proton momentum β_p . This combi-




FIG. 4: **Top left:** Velocity distribution from the time-of-flight measurement (β_{ToF}) for the real coincidence time cut shown in the bottom plot and described in the text. **Top right:** Distribution of the corrected coincidence time for protons. The estimated random coincidence contribution is overlaid on top of the coincident proton peak. Protons were selected using the ToF cut shown in the bottom plot. **Bottom:** Typical spectrum of the difference in the velocities as determined by the time-of-flight technique and proton momentum versus the corrected coincidence time.

nation of cuts allowed the retention of those protons that underwent interactions in the SOS detector hut. These events form a shoulder that extends from the proton coincident peak toward negative values of $\beta_{\text{ToF}} - \beta_p$ (Fig. 4 bottom).

Random coincidences, also present beneath the true coincidence peak (Fig. 4 top right), contributed a background in the final data sample (Fig. 7). These were averaged and removed by selecting a sample of random coincidences from five RF peaks (the selection procedure is shown in the bottom of Fig. 4). The random-subtracted distribution for any physics quantity was then obtained by subtracting the corresponding distribution for real and random samples, weighted by a 3:5 ratio to account for the differing numbers of peaks in the respective samples.

The kinematics of the ω channel for a fixed target is diagrammatically shown in Fig. 5. Kinematic quantities




FIG. 5: Fixed target ${}^1\text{H}(e, e'p)X$ scattering process. Here, as well as in the text below, the energy and three-momentum transfer ν and \vec{q} are given by $\nu = E - E'$ and $\vec{q} = \vec{k} - \vec{k}'$.

characterizing the process can be expressed employing the notation of Fig. 5:

$$Q^2 = -[(E, \vec{k}) - (E', \vec{k}')]^2 \stackrel{m_e \rightarrow 0}{\approx} 4EE' \sin^2(\theta_{e'}/2) \quad (1)$$

$$W^2 = [(M_p, \vec{0}) + (\nu, \vec{q})]^2 = M_p^2 + 2M_p\nu - Q^2, \quad (2)$$

$$t = [(M_p, \vec{0}) - (E_p, \vec{p})]^2 = 2M_p(M_p - E_p), \quad (3)$$

$$M^2 = [(M_p, \vec{0}) + (E, \vec{k}) - (E', \vec{k}') - (E_p, \vec{p})]^2 = W^2 + M_p^2 - 2E_p(M_p + \nu) + 2|\vec{q}||\vec{p}| \cos \theta_{\gamma p}, \quad (4)$$

where $\theta_{e'}$ is the laboratory electron scattering angle and $\theta_{\gamma p}$ is the proton scattering angle with respect to the virtual photon direction. Q^2 is square of the four-momentum transfer to the target, W is the invariant mass of the virtual photon-proton system, t is the squared four-momentum transfer to the proton, and M is the mass of the system of undetected particles.

Reconstruction of the missing mass, performed accord-




FIG. 6: Light vector meson electroproduction. The histogram shows events for all accepted momenta for one setting. Note the presence of the pseudoscalar η meson signal.

ing to Eqn. (4), reveals a spectrum with a strong ω meson signal atop a complicated background (Fig. 6). The

data were corrected for trigger inefficiency ($<1\%$), track reconstruction inefficiencies ($\sim 10\%$), particle ID inefficiencies ($\sim 2\%$), and computer and electronic dead times ($\sim 5\%$). In the CM system, the virtual photon cross section for ω production $d\sigma_v/d\Omega^*$ is given in terms of the conventional two-particle coincidence cross section

$$\frac{d\sigma}{dp_{e'}d\Omega_{e'}d\Omega^*} = \Gamma_T \frac{d\sigma_v}{d\Omega^*}, \quad (5)$$

where Γ_T is the virtual photon flux. The virtual photon cross section can be decomposed into transverse (σ_T), longitudinal (σ_L), and interference terms (σ_{TT} , σ_{LT}), such that

$$\frac{d\sigma_v}{d\Omega^*} = \sigma_U + \varepsilon \cos 2\phi^* \sigma_{TT} + \sqrt{\frac{\varepsilon(\varepsilon+1)}{2}} \cos \phi^* \sigma_{LT}, \quad (6)$$

where $\sigma_U = \sigma_T + \varepsilon \sigma_L$, ε is the virtual photon polarization parameter, and ϕ^* is the relative angle between the electron scattering plane and hadron production plane.

The biggest challenge in cross section extraction was the separation of the data into the physics backgrounds and ω meson production (Fig. 7). This was accomplished by using a Monte Carlo program to simulate both processes, the dominant background as well as ω production. The background was modeled as a combination of two processes, electroproduction of the neutral ρ meson and multi-pion production.

Production of the ρ was assumed to be purely diffrac-




FIG. 7: Missing mass distribution for $^1\text{H}(e, e'p)X$ showing the decomposition into a peak for ω and the background.

tive [10],

$$\frac{d\sigma}{d\Omega^*dM} = \left(\frac{M_\rho}{M}\right)^n B_\rho(M) D e^{bt'}, \quad (7)$$

where $t' = t - t_{\min}$, with t_{\min} being the momentum transfer when the scattering occurs along the virtual photon direction. In the above expression, coefficients D and b are Q^2 and W dependent to account for their variation near threshold and D , at $Q^2 = 0$, corresponds to the photoproduction cross section. The skewness of the ρ meson

shape, apparent from other experiments, was accounted for by using the Ross-Stodolsky parameterization [11] (in Eqn. (7) first factor on the right-hand side) with the exponent $n = 5.2$ coming from a fit to the DESY data [1]. For both the background and the ω meson, the mass distributions were generated according to a fixed width relativistic Breit-Wigner distribution

$$B_v(M) = \frac{M_v^2 \Gamma_v^2}{(M^2 - M_v^2)^2 + M_v^2 \Gamma_v^2}, \quad (8)$$

where v is ρ or ω with $M_\omega = 781.94$ MeV, $\Gamma_\omega = 8.43$ MeV, $M_\rho = 768.1$ MeV, and $\Gamma_\rho = 150.7$ MeV [12].

The multi-pion processes were collectively modeled as a Lorentz invariant electroproduction phase space for two-body production of a fictitious particle with arbitrary mass M . This term is meant to account for all physically allowed reactions (W is well above the $\pi\pi$ threshold) that result in more than three particles (including the electron and proton) in the final state. The flatly distributed low yield of events (approximately 2% at most settings) coming from the aluminum walls of the liquid hydrogen target were also treated as a part of the phase space background. The phase space was simulated by

$$\frac{d\sigma}{d\Omega^*dM} = \frac{1}{32\pi^2} \left(\frac{p^*}{q^*}\right) \frac{M}{W^2}, \quad (9)$$

where q^* and p^* are the initial and final momenta in the CM frame, respectively.

The production of the ω meson was simulated with a cross section assumed to be t -channel unnatural parity exchange

$$\frac{d\sigma}{d\Omega^*dM} = B_\omega(M) (\sigma_T^\pi + \varepsilon \sigma_L^\pi), \quad (10)$$

with σ_T^π , σ_L^π being the transverse and longitudinal parts of the corresponding cross section [3]. Within this model, the longitudinal contribution σ_L^π is insignificant because it is an order of magnitude smaller than σ_T^π for the kinematic regime of the experiment. Natural parity exchange was neglected because it is also roughly one order of magnitude smaller than σ_T^π within this regime. Similarly neglected were the nearly vanishing contributions from the interference terms σ_{TT} and σ_{LT} . This amounts to modeling the total cross section using only the largest contribution. The Monte Carlo program simulated finite target effects (multiple scattering and ionization energy losses), acceptance corrections, and radiative processes. The radiative corrections were modeled after the approximations from Ref. [13]. They were accounted for by altering the incident and scattered electron kinematics and applying loop and vertex corrections which modify the cross section, but do not modify the missing mass distribution. Having simulated all the processes for each kinematic setting, the data and Monte Carlo events were binned in CM




FIG. 8: Missing mass M and invariant mass W distributions broken down into individual contributions for two different momentum settings but the same angular settings. Solid circles with error bars are the data. The shaded histogram is the full Monte Carlo fit. The dotted line histogram corresponds to the resulting phase space yield, the solid line histogram to the ω yield, and the dashed line histogram to the ρ^0 yield.

scattering angle θ^* . Finally, a binned maximum likelihood fit was performed simultaneously in missing mass, W , Q^2 , t , and θ^* . The approach incorporated in the fit was developed by R. Barlow [14]. The likelihood function accounted for fluctuations in the data and Monte Carlo distributions due to finite statistics. Its maximization allowed the search for the overall strengths, p_i , of each process modeled, so that the resulting yields for each bin satisfy the relation

$$Y_{\text{DATA}} = Y_{\text{MC}} = p_1 Y_\omega + p_2 Y_\rho + p_3 Y_{\text{phsp}}. \quad (11)$$

Results of the fitting process for the high momentum setting of the hadron arm, $p_0^{\text{SOS}} = 1.077$ GeV, and the intermediate momentum setting, $p_0^{\text{SOS}} = 0.929$ GeV, for the same angular setting of $\theta_{\gamma\text{P}} = 4.33^\circ$, are shown in Fig. 8 and Fig. 9. Figures 10 and 11 show the result of summation of the fits for all θ^* bins within these two hadron arm settings, respectively, thus reflecting the goodness of the fit. Performing the fit allowed separation of the raw data into the Monte Carlo determined




FIG. 9: Distributions of four-momentum transfers Q^2 and t for the same settings as described in the caption of Fig. 8. The cut-off at low values of $-t$ reflects the proton momentum cut applied in the analysis since the proton energy is directly proportional to t in the fixed target regime (see Eqn. 3).

background, consisting of the ρ meson and phase-space contributions, and the ω meson signal, thus obtaining the data yields (Fig. 12). Subsequently, the differential virtual photon cross section was computed by scaling the model cross section by the data yield Y_{DATA}^ω ($\approx p_1 Y_\omega$), normalized to the simulated yield

$$\frac{d\sigma_v}{d\Omega^*} = \frac{Y_{\text{DATA}}^\omega}{Y_{\text{MC}}^\omega} \left(\frac{d\sigma}{d\Omega^*} \right)_{\text{MC}}. \quad (12)$$

The Monte Carlo yield Y_{MC}^ω was evaluated by integrating the model cross section, $d\sigma/d\Omega^*dM$ over the entire acceptance of the apparatus and binning the result in the CM scattering angle. For any θ_i^* bin, this process can be expressed as

$$Y_{\text{MC}}^\omega = \int_{\mathcal{A}(\theta_i^*)} \Gamma_T \mathcal{R} \frac{d\sigma}{d\Omega^*dM} dQ^2 dW d\phi_{e'} d\Omega^* dM, \quad (13)$$

where \mathcal{R} represents the multiplicative part of the radiative corrections and $\mathcal{A}(\theta_i^*)$ is the acceptance for the given θ_i^* bin. In the above expression, mass was integrated




FIG. 10: Fits for two different momentum settings summed over θ^* bins for missing mass (top) and invariant mass (bottom). Fig. 8 contains the legend explanation.

over the ω line shape (Eqn. 8). The cross section was extracted at $Q^2 \approx 0.5 \text{ GeV}^2$ for 74 bins in θ^* , mostly for backward directions in the γ^*p CM system. Here, the Hand [15] convention was adopted in evaluating the virtual photon flux Γ_T . Identifying the ω meson production using only the $e'p$ final states introduced a statistical error of less than 25%. Systematic uncertainties associated with the background subtraction are less than 12%. Fixed electron kinematics and limited out-of-plane acceptance reduced the range of accepted ϕ^* angles to $\pm 30^\circ$ about 0° for the outermost angular setting ($\theta_{\gamma p} = 17.3^\circ$). This ϕ cut was also applied to the data of all other settings.

IV. RESULTS

With the use of the procedures described above, angular distributions of the differential cross sections for electroproduction of the ω meson were extracted for two different average values of the invariant mass W . The data were divided into two sets according to the average W which, for each data point, was determined using the




FIG. 11: Fits for two different momentum settings summed over θ^* bins for four-momentum transfer t (top) and four-momentum transfer Q^2 (bottom). Fig. 8 contains the legend explanation.

results of the fit. These two sets form the angular distributions that correspond to mean invariant masses $\langle W \rangle$ of 1.750 GeV and 1.790 GeV. The results are presented in Tables II and III.

These two sets of the data, however, do not constitute two independent angular distributions. There are large overlaps in the W ranges for both distributions that can readily be seen in the bottom of Figs. 8 and 10. Therefore, the cross sections of both angular distributions were scaled to a reference W of 1.785 GeV. This was done by rescaling their corresponding kinematic parts, *i.e.* phase space factors normalized to the incoming particle flux. Due to a significant variation with mass, the scaling factor was determined on an event-by-event basis and then averaged. The scaling can quantitatively be described by

$$\left(\frac{d\sigma}{d\Omega^*} \right)_{\text{scaled}} = \frac{\Gamma(W_{\text{ref}})}{\Gamma(W)} \frac{d\sigma}{d\Omega^*}, \quad (14)$$

where $\Gamma(W) = p^*(W)/q^*(W)W^2$ is a normalized phase space factor (compare with Eqn. (9)) and p^* and q^* are, respectively, the 3-momenta in the CM frame of the ω




FIG. 12: The results of signal-background separation, or, equivalently, ω yield extraction, for two different momentum settings of the hadron arm but the same, $\theta_{\gamma p} = 4.33^\circ$, angular setting, Top: central momentum $p_0 = 1.077$ GeV. Bottom: central momentum $p_0 = 0.929$ GeV.

and the virtual photon which, for fixed W , are determined only by the masses of the interacting particles. The result of this procedure is shown in Fig. 13. Correcting for the phase space, opening up above the threshold, removes practically all of the observed W dependence. It also shows that the shape of the distribution is not trivially induced by W variations of the phase space factors.

The enhancement of the backward-angle cross section over t -channel unnatural parity exchange (Fraas model, dashed line) is evident. This was suggested by the earlier electroproduction [2] and photoproduction [4, 5] data. Such a departure from the smooth fall-off of the t -channel processes, either in the angular distribution or t -dependence, has been attributed, theoretically, to s - and u -channel resonance contributions. Even though the energy dependence may not be sensitive to the details of the model, since it is integrated over full angular range, the inclusion of resonance formation was also necessary to reproduce the near threshold strength of the photoproduction cross section (see [16, 17]). Recent examples of such calculations [16, 17, 18, 19, 20, 21] mainly address

TABLE II: Differential cross sections for the lower average W ($\langle W \rangle = 1.75$ GeV). The bin width is 5^0 , centered on the quoted value, except for the first bin whose width is 10^0 . The cross sections were extracted for $|\phi^*| < 30^\circ$.

| θ^* (deg) | $d\sigma/d\Omega^*$ ($\mu\text{b}/\text{sr}$) | Uncertainty | | $\langle W \rangle$ (GeV) | $\langle Q^2 \rangle$ (GeV^2) |
|---------------------|--|-------------|-------|------------------------------|---|
| | | Stat. | Syst. | | |
| 45 | 0.257 | 0.057 | 0.015 | 1.753 | 0.501 |
| 75 | 0.116 | 0.026 | 0.011 | 1.745 | 0.512 |
| 80 | 0.170 | 0.026 | 0.006 | 1.747 | 0.511 |
| 85 | 0.112 | 0.024 | 0.006 | 1.747 | 0.510 |
| 90 | 0.131 | 0.024 | 0.006 | 1.747 | 0.510 |
| 95 | 0.163 | 0.024 | 0.008 | 1.749 | 0.510 |
| 100 | 0.176 | 0.023 | 0.010 | 1.752 | 0.509 |
| 101 | 0.170 | 0.028 | 0.012 | 1.752 | 0.509 |
| 105 | 0.260 | 0.023 | 0.012 | 1.755 | 0.505 |
| 106 | 0.267 | 0.028 | 0.012 | 1.756 | 0.508 |
| 110 | 0.292 | 0.024 | 0.012 | 1.758 | 0.504 |
| 111 | 0.311 | 0.025 | 0.013 | 1.761 | 0.505 |
| 115 | 0.440 | 0.026 | 0.014 | 1.763 | 0.501 |
| 120 | 0.466 | 0.025 | 0.013 | 1.766 | 0.499 |
| 125 | 0.425 | 0.025 | 0.013 | 1.766 | 0.498 |
| 130 | 0.399 | 0.026 | 0.012 | 1.762 | 0.498 |
| 135 | 0.412 | 0.031 | 0.012 | 1.759 | 0.500 |
| 138 | 0.400 | 0.031 | 0.012 | 1.749 | 0.497 |
| 140 | 0.458 | 0.044 | 0.012 | 1.755 | 0.501 |
| 143 | 0.466 | 0.033 | 0.012 | 1.751 | 0.498 |
| 148 | 0.367 | 0.028 | 0.010 | 1.751 | 0.497 |
| 153 | 0.352 | 0.030 | 0.010 | 1.750 | 0.501 |
| 158 | 0.308 | 0.031 | 0.010 | 1.748 | 0.501 |
| 163 | 0.353 | 0.039 | 0.009 | 1.747 | 0.501 |
| 168 | 0.288 | 0.040 | 0.010 | 1.745 | 0.504 |
| 173 | 0.199 | 0.050 | 0.012 | 1.742 | 0.510 |

SAPHIR data [5]. Some of these works [20, 21] showed that the dominant contributions could come from the missing resonances, $N_{3/2}^+(1910)$, and the $N_{3/2}^-(1960)$ (the latter is labeled $D_{13}(2080)$ by the Particle Data Group). Other calculations, however, differ in predicting which nucleonic excitations could contribute in the s -channel. It was found that the contribution from two resonances, $P_{13}(1720)$ and $F_{15}(1680)$, dominated and their inclusion was necessary to reproduce the available photoproduction data near threshold [16, 17, 18, 19].

From the point of view of the present work, the most interesting result of these theoretical models is that the nucleon resonances are the favored mechanism for producing backward-angle enhancements in the differential cross section. The solid line in Fig. 13 shows the comparison of the data with an unpublished, as of this writing, electroproduction calculation [22] complementary to the photoproduction model [17]. In this model, the diffractive nature of ω production is described by Pomeron exchange based on Regge phenomenology and SU(3) flavor symmetry. This contribution dominates the cross section above the resonance region. Neutral π exchange

TABLE III: Differential cross sections for the higher average W ($\langle W \rangle = 1.790$ GeV). The bin width is 5° , centered on the quoted value, except for the first two bins whose width is 10° . The cross sections were extracted for $|\phi^*| < 30^\circ$.

| θ^* (deg) | $d\sigma/d\Omega^*$ ($\mu\text{b}/\text{sr}$) | Uncertainty | | $\langle W \rangle$ (GeV) | $\langle Q^2 \rangle$ (GeV^2) |
|---------------------|--|-------------|-------|------------------------------|---|
| | | Stat. | Syst. | | |
| 25 | 0.501 | 0.058 | 0.015 | 1.778 | 0.505 |
| 35 | 0.360 | 0.053 | 0.015 | 1.765 | 0.502 |
| 62 | 0.229 | 0.034 | 0.015 | 1.808 | 0.493 |
| 67 | 0.263 | 0.030 | 0.014 | 1.808 | 0.489 |
| 72 | 0.186 | 0.027 | 0.014 | 1.811 | 0.488 |
| 73 | 0.171 | 0.033 | 0.014 | 1.771 | 0.504 |
| 77 | 0.193 | 0.025 | 0.014 | 1.814 | 0.484 |
| 78 | 0.168 | 0.032 | 0.014 | 1.773 | 0.503 |
| 82 | 0.141 | 0.023 | 0.013 | 1.819 | 0.479 |
| 83 | 0.175 | 0.030 | 0.013 | 1.775 | 0.502 |
| 84 | 0.173 | 0.030 | 0.012 | 1.780 | 0.512 |
| 88 | 0.225 | 0.029 | 0.012 | 1.779 | 0.500 |
| 87 | 0.226 | 0.024 | 0.012 | 1.821 | 0.477 |
| 89 | 0.256 | 0.024 | 0.012 | 1.782 | 0.502 |
| 92 | 0.251 | 0.027 | 0.012 | 1.825 | 0.475 |
| 93 | 0.249 | 0.028 | 0.012 | 1.784 | 0.496 |
| 94 | 0.237 | 0.020 | 0.012 | 1.789 | 0.498 |
| 97 | 0.282 | 0.031 | 0.012 | 1.827 | 0.473 |
| 98 | 0.329 | 0.027 | 0.012 | 1.791 | 0.493 |
| 99 | 0.321 | 0.019 | 0.012 | 1.800 | 0.491 |
| 102 | 0.309 | 0.047 | 0.012 | 1.827 | 0.470 |
| 103 | 0.395 | 0.028 | 0.011 | 1.798 | 0.489 |
| 104 | 0.352 | 0.018 | 0.012 | 1.808 | 0.485 |
| 108 | 0.365 | 0.030 | 0.012 | 1.796 | 0.488 |
| 109 | 0.391 | 0.020 | 0.011 | 1.811 | 0.484 |
| 113 | 0.318 | 0.037 | 0.012 | 1.791 | 0.488 |
| 114 | 0.450 | 0.023 | 0.011 | 1.814 | 0.481 |
| 116 | 0.396 | 0.025 | 0.011 | 1.772 | 0.500 |
| 119 | 0.515 | 0.029 | 0.011 | 1.816 | 0.481 |
| 121 | 0.486 | 0.024 | 0.010 | 1.785 | 0.493 |
| 124 | 0.524 | 0.044 | 0.010 | 1.816 | 0.482 |
| 126 | 0.503 | 0.023 | 0.010 | 1.795 | 0.488 |
| 131 | 0.506 | 0.024 | 0.010 | 1.796 | 0.488 |
| 136 | 0.518 | 0.027 | 0.010 | 1.792 | 0.490 |
| 140 | 0.530 | 0.024 | 0.010 | 1.772 | 0.489 |
| 141 | 0.495 | 0.031 | 0.010 | 1.788 | 0.492 |
| 145 | 0.538 | 0.020 | 0.010 | 1.781 | 0.488 |
| 146 | 0.536 | 0.046 | 0.010 | 1.779 | 0.501 |
| 150 | 0.492 | 0.019 | 0.010 | 1.783 | 0.489 |
| 151 | 0.471 | 0.102 | 0.014 | 1.765 | 0.516 |
| 155 | 0.431 | 0.019 | 0.010 | 1.780 | 0.492 |
| 160 | 0.425 | 0.021 | 0.010 | 1.777 | 0.494 |
| 165 | 0.439 | 0.026 | 0.010 | 1.773 | 0.498 |
| 166 | 0.652 | 0.093 | 0.014 | 1.785 | 0.470 |
| 170 | 0.442 | 0.036 | 0.011 | 1.764 | 0.504 |
| 171 | 0.419 | 0.059 | 0.012 | 1.780 | 0.482 |
| 175 | 0.485 | 0.070 | 0.012 | 1.755 | 0.513 |
| 176 | 0.392 | 0.077 | 0.014 | 1.778 | 0.489 |

in the t -channel is included to account for the peaking of the cross section in the forward direction, especially near threshold. Resonance formation processes in the s - and u -channel that dominate intermediate and backward scattering angles, where the other contributions are small, were modeled in an $SU(6) \times O(3)$ quark model symmetry limit. All contributions, summed coherently, give a strongly ϕ^* -dependent cross section (Eqn. (6)). To correctly compare this theoretical calculation with the data, the model was integrated over a range of the azimuthal angle ϕ^* corresponding to the cut used in the data analysis. The model was also averaged over the appropriate W and Q^2 ranges.

V. CONCLUSIONS

Cross sections for the ω meson electroproduction were obtained from the $^1\text{H}(e, e'p)\omega$ reaction at $E_e = 3.245$ GeV. The angular distribution of the differential cross section in the threshold regime has unprecedented granularity and much smaller statistical uncertainties than in previous work. The angular distribution exhibits a substantial backward-angle enhancement of the cross section over the pure t -channel expectation, similar to that found in the DESY [2], photoproduction [4], and SAPHIR data [5].

In comparing the result of this work to the Zhao model [22], the similarity of the angular distributions is evident. In the view of these results, this analysis pro-




FIG. 13: The angular distributions for different average W and for $|\phi^*| < 30^\circ$. Error bars are statistical. The DESY data correspond to the four-momentum transfer $Q^2 = 0.77$ GeV^2 , $W = 1.82$ GeV and full ϕ^* range. The Fraas model shown here was used in the cross section extraction. Both distributions were scaled to $W = 1.785$ GeV. The scaling procedure is described in the text.

vides significant evidence for resonance formation, possibly s -channel, in the $\gamma^*p \rightarrow \omega p$ reaction. It is worth noting that, although elastic πN scattering constitutes the main source of information on the nucleon excitation spectrum, it alone cannot distinguish among existing theoretical models [23], many of which predict a much richer baryonic, hence nucleonic, spectrum than currently observed [24, 25, 26, 27, 28, 29, 30, 31, 32]. If they exist, these states are either being masked by neighboring resonances with stronger couplings or they are altogether decoupled from the πN channel. There are decay modes, other than πN , however, that have sizeable resonance coupling constants [26, 33]. A calculation, based on the symmetric quark model [34], indeed predicts that vector meson decay channels, $N\rho$ and $N\omega$, have appreciable resonance couplings. Electroproduction of ω mesons, enhanced by its isospin selectivity, may therefore provide additional evidence in the search for resonances unobserved in πN scattering.

Acknowledgments

The authors would like to express sincere thanks to Dr. Qiang Zhao for sharing his electroproduction calculation

and for fruitful discussions on the underlying theory. The authors would like to acknowledge the support of the staff of the Accelerator division of Jefferson Lab. This work was supported in part by the U.S. Department of Energy under contract W-31-109-Eng-38 for Argonne National Laboratory, by contract DE-AC05-84ER40150, under which the Southeastern Universities Research Association (SURA) operates the Thomas Jefferson National Accelerator, and the National Science Foundation (grant no. NPS-PHY-9319984). It was also in part supported by Temple University, Philadelphia PA.

Pawel Ambrozewicz would like to thank Dr. Kees de Jager, the leader of Hall A at Jefferson Lab, for the support that allowed him to finish the analysis presented in this paper.

-
- [1] P. Joos et al., Nucl. Phys. **B113**, 53 (1976).
 - [2] P. Joos et al., Nucl. Phys. **B122**, 365 (1977).
 - [3] H. Fraas, Nuclear Physics **B36**, 191 (1971).
 - [4] ABBHHM Collaboration, Phys. Rev. **175**, 1669 (1968).
 - [5] F. Klein, Ph.D. thesis, Bonn University (1996).
 - [6] Jefferson Laboratory experiment E91-016 (Ben Zeidman spokesperson).
 - [7] P. Ambrozewicz, Ph.D. thesis, Temple University (2001).
 - [8] J. Dunne, *Cryo and dummy target information* (1997), JLab Hall C Internal Report (unpublished).
 - [9] D. Meekins, Ph.D. thesis, College of William & Mary (1998).
 - [10] H. Fraas and D. Schildknecht, Nuclear Physics **B14**, 543 (1969).
 - [11] M. Ross and L. Stodolsky, Phys. Rev. **149**, 1172 (1966).
 - [12] C. Caso et al., Eur. Phys. J. **C3** (1998).
 - [13] R. Ent et al., Phys. Rev. **64**, 054610 (2001).
 - [14] R. Barlow, Comp. Phys. Commun. **77**, 219 (1993).
 - [15] L. Hand, Phys. Rev. **129**, 1834 (1963).
 - [16] Q. Zhao, Nucl. Phys. **A675**, 217 (2000).
 - [17] Q. Zhao, Phys. Rev. C **63**, 025203 (2001).
 - [18] Q. Zhao, Z. Li, and C. Bennhold, Phys. Lett. **B436**, 42 (1998).
 - [19] Q. Zhao, Z. Li, and C. Bennhold, Phys. Rev. **C58**, 2393 (1998).
 - [20] Y. Oh, A. Titov, and T.-S. Lee (2000), arXiv:nucl-th/0004055 (preprint).
 - [21] Y. Oh, A. Titov, and T.-S. Lee, Phys. Rev. C **63**, 025203 (2001).
 - [22] Q. Zhao, *Private communication* (2003).
 - [23] N. Isgur and R. Koniuk, Phys. Rev. **D21**, 1868 (1980).
 - [24] R. Capps, Phys. Rev. Lett. **33**, 1637 (1974).
 - [25] S. Capstick and N. Isgur, Phys. Rev. **D34**, 2809 (1986).
 - [26] S. Capstick and W. Roberts, Phys. Rev. **D49**, 4570 (1994).
 - [27] R. Cutkosky and R. Hendrick, Phys. Rev. D **16**, 2902 (1977).
 - [28] C. Forsythe and R. Cutkosky, Z. Phys. C **18**, 219 (1983).
 - [29] N. Isgur and G. Karl, Phys. Lett. **72B**, 109 (1977).
 - [30] N. Isgur and G. Karl, Phys. Lett. **74B**, 353 (1978).
 - [31] N. Isgur and G. Karl, Phys. Rev. **D18**, 4187 (1978).
 - [32] N. Isgur and G. Karl, Phys. Rev. **D19**, 2653 (1979).
 - [33] S. Capstick, Phys. Rev. **D46**, 2864 (1992).
 - [34] S. Capstick and W. Roberts, Prog. Part. Nucl. Phys. **45**, S241 (2000).