# Event-by-Event Fluctuations at 40, 80, and 158 AGeV/c in Pb+Au Collisions

## Hiroyuki Sako† and Harald Appelshäuser for the CERES/NA45 collaboration

Gesellschaft für Schwerionenforschung (GSI), 64291 Darmstadt, Germany

E-mail: † h.sako@gsi.de

Abstract. Event-by-event fluctuations of mean  $p_T$  and net charge in 40, 80, and 158 AGeV/c Pb+Au collisions are presented. The observed dynamical mean  $p_T$ fluctuations in central events of about 1 % are very similar to results from RHIC. The centrality dependence of mean  $p_T$  fluctuations at 158 AGeV/c shows a peculiar nonmonotonic pattern with respect to the extrapolation from the fluctuations measured in p+p collisions. Dynamical net charge fluctuations smaller than the charge conservation limit are observed. After correction for charge conservation, the measured fluctuation signal in central events is comparable to RHIC results.

#### 1. Introduction

Event-by-event fluctuations of mean  $p_T$  have been proposed as a possible probe to search for the phase transition, especially the QCD critical point via their nonmonotonic variation with control parameters such as beam energy and centrality [1]. On the other hand, the fluctuations can be used to evaluate the extent of thermal equilibrium by comparison with the extrapolated fluctuations from nucleon-nucleon collisions. Suppressed net charge fluctuations have been proposed as a possible signal for the formation of a QGP, due to the smaller charge units of deconfined (anti-) quarks compared to those of confined hadrons [2, 3]. Comparisons with cascade models may clarify whether the observed fluctuations are described by resonance gas models.

A detailed description of CERES experiment at the CERN-SPS can be found elsewhere [4, 5]. For the present analysis, momenta of charged particle tracks produced in Pb+Au collisions at 40, 80, and 158 AGeV/c have been measured with a cylindrical Time Projection Chamber. The data set comprises a wide range of collision centrality. The acceptance of the TPC is around mid-rapidity ( $2.0 < \eta < 2.9$ ) and covers  $2\pi$  in azimuth (at 40 AGeV/c only  $1.3\pi$ , see also [5]).

#### 2. Mean $p_T$ Fluctuations

Two measures are used to evaluate mean  $p_T$  fluctuations. The  $\Sigma_{p_T}$  measures the dynamical fluctuation normalized with mean  $p_T$  [5]. This measure is proportional to



Figure 1:  $\Sigma_{p_T}$  (left) and  $F_{p_T}$  (right) as a function of Figure 2:  $\Sigma_{p_T}$  as a function of  $\sqrt{s_{NN}}$  in  $\Delta \eta$  at  $0.1 < p_T < 1.5 \text{ GeV/c}$  in 6.5 % central events.

central events.

mean covariance of all charged particle pairs per event [6]. Another measure  $F_{p_T}$  [8] is defined as a deviation from 1 of the ratio of the r.m.s. of the event-by-event mean  $p_T$  distribution in real events to that in mixed events. The  $F_{p_T}$  is approximately proportional to  $\langle N \rangle \Sigma_{p_T}^2$ , where  $\langle N \rangle$  is the mean charged particle multiplicity in the acceptance. In Fig. 1,  $\Sigma_{p_T}$  has a saturation tendency at the pseudo-rapidity interval  $(\Delta \eta)$  more than 0.4, whereas  $F_{p_T}$  steadily increases. We therefore use  $\Sigma_{p_T}$  to compare CERES results ( $\Delta \eta = 0.5$ ) with RHIC results ( $\Delta \eta = 0.7 \sim 2$ ).

Fig. 2 shows  $\Sigma_{p_T}$  as a function of nucleon-nucleon center of mass energy  $(\sqrt{s_{NN}})$ at 2.2 <  $\eta$  < 2.7 and 0.1 <  $p_T$  < 2 GeV/c in 6.5 % central events, compared to RHIC data at  $\sqrt{s_{NN}} = 130 \text{ GeV} [9, 10]$ , and 200 GeV [8]. For comparison to RHIC data, the CERES data are not corrected for short range momentum correlations. The observed fluctuations at SPS and at RHIC are similarly about 1 %. No indication for non-monotonic variation or enhanced fluctuations is observed. Note that a predicted fluctuation at the critical point is  $\sim 2\%$  [1], far above the measured values.

Fig. 3 shows fluctuations corrected for short range momentum correlations at 158 AGeV/c as a function of the number of participants  $(N_{part})$ . The fluctuations between the minimum-bias data set and the central data set are consistent. As a baseline (dashed lines) reflecting the superposition of individual nucleon-nucleon collisions, we employ the formula  $\Sigma_{p_T}^{AA} = 12\% \cdot (\langle N_{part} \rangle/2)^{-1/2}$ , where the 12 % is a measured fluctuation in p+p collisions at ISR [7], which corresponds to  $F_{p_T} \simeq 1.4$  % in the CERES acceptance. Significant deviations from the baseline are observed in semi-central events, whereas in peripheral events (centrality  $\geq 66\%$ ) and in central events (centrality  $\leq 20\%$ ), the fluctuations are consistent with the baseline. The maximum of  $F_{p_T}$  is about 2.8 % at  $N_{part}$  of 50 ~ 100, corresponding to 25 ~ 50 % central events. The observed centrality dependence as well as the increase of  $F_{p_T}$  with the upper  $p_T$  cut is similar to the observations by NA49 [11], PHENIX [8], and STAR [10].

In Fig. 4,  $F_{p_T}$  of RQMD and UrQMD without rescattering is consistent with the





Figure 3:  $\Sigma_{p_T}$  (left) and  $F_{p_T}$  (right) as a function Figure 4:  $F_{p_T}$  (left) and  $\overline{p_T}$  (right) as a function of  $N_{part}$  at 158 AGeV/c.

of  $N_{part}$  compared to RQMD and UrQMD at 158 AGeV/c.

baseline as expected. With rescattering, RQMD reproduces qualitatively both the nonmonotonic dependence of fluctuations and the increasing tendency of mean  $p_T$ , while UrQMD fails to describe both observables.

### 3. Net charge fluctuations

As a measure of net charge fluctuations, we use  $\nu_{dyn}$ , representing dynamical fluctuations of the difference between normalized multiplicity of positive particles and that of negative particles [12]. The measure is advantageous since the correction for the global charge conservation is additive, and no efficiency correction is necessary.

Fig. 5 shows  $\nu_{dyn}$  multiplied with mean charged particle multiplicity within the acceptance  $(\langle N \rangle)$  as a function of  $N_{part}$  at the three beam energies. The dashed lines show the charge conservation limit, defined as  $-4\langle N\rangle/\langle N\rangle_{4\pi}$ , where  $\langle N\rangle_{4\pi}$  is the total charged particle multiplicity which is assumed to scale with  $\langle N \rangle$ . The  $\langle N \rangle_{4\pi}$  is estimated based on the NA49 measurements [13]. The observed fluctuations are lower than the charge conservation limit. Since the fluctuations are much above the QGP model predictions of  $\sim -3.5$  [2, 3], no indication for the formation of a deconfined phase is found. Decreasing tendencies of  $\langle N \rangle \nu_{dyn}$  with  $N_{part}$  are observed at the three beam energies. Whether this is due to a deviation of the charge conservation limit from the multiplicity scaling, or due to a real deviation of the fluctuations from the multiplicity scaling, is under investigation.

Fig. 6 shows, as a function of  $\sqrt{s_{NN}}$ , a compilation of CERES, STAR [14, 15], and PHENIX results [16] for  $\nu_{dyn}$  corrected for charge conservation  $(\tilde{\nu}_{dyn})$ . A decrease of  $\tilde{\nu}_{dyn}$  is observed at SPS energies, while a decrease is tiny from SPS top energy to RHIC energies. Both RQMD and UrQMD are in reasonable agreement with the CERES data.



Figure 5:  $\langle N \rangle \nu_{dyn}$  as a function of  $N_{part}$  at 2.05 <  $\eta < 2.85$  and  $0.1 < p_T < 2.5$  GeV/c.



Figure 6:  $\tilde{\nu}_{dyn}$  as a function of  $\sqrt{s_{NN}}$  in central events.

#### 4. Conclusions

Dynamical mean  $p_T$  fluctuations of ~ 1 % in central events are observed, which are similar to the fluctuations measured at RHIC. The  $F_{p_T}$  shows non-monotonic centrality dependence with a maximum of ~ 2.8 % in semi-central events, which is about a factor of 2 enhanced over the p+p extrapolation. A similar tendency is observed also by NA49, PHENIX, and STAR. RQMD qualitatively reproduces both enhancement of fluctuations in semi-central events and increase of mean  $p_T$  as a function of centrality.

The observed dynamical net charge fluctuations are smaller than the charge conservation limit. The  $\langle N \rangle \nu_{dyn}$  slightly decreases from peripheral to central events. The  $\tilde{\nu}_{dyn}$  decreases at SPS energies but varies little from the SPS top energy to RHIC energies. UrQMD and RQMD reproduce the observed fluctuations at SPS.

#### References

- [1] Stephanov M A, Rajagopal K and Shuryak E V 1999 Phys. Rev. D 60 114028
- [2] Asakawa M, Heinz U and Mueller B 2000 Phys. Rev. Lett. 85 2072
- [3] Jeon S and Koch V 2000 Phys. Rev. Lett. 85 2076
  - Bleicher M, Jeon S and Koch V 2000 Phys. Rev. C 62 061902
- [4] CERES Collaboration, Adamová D et al 2003 Nucl. Phys. A 714 124
- [5] CERES Collaboration, Adamová D et al 2003 Nucl. Phys. A 727 97
- [6] Voloshin S, Koch V, and Ritter H 1999 Phys. Rev. C 60 024901
- [7] Braune K et al 1983 Phys. Lett. B 123 467
- [8] PHENIX Collaboration, Adler S et al 2003 nucl-ex/0310005
- [9] STAR Collaboration, Voloshin S et al 2001 nucl-ex/0109006
- [10] STAR Collaboration, Adams J et al 2003 nucl-ex/0308033
- [11] NA49 Collaboration, Anticic T et al 2003 hep-ph/0311009
- [12] Pruneau C et al 2002 Phys. Rev. C 66 044904
- [13] NA49 Collaboration, 2002 Phys. Rev. C 66 054902
- [14] STAR Collaboration, Adams J et al 2003 Phys. Rev. C 68 044905
- [15] Pruneau C et al 2003 nucl-ex/0304021
- [16] PHENIX Collaboration, Adcox K et al 2002 Phys. Rev. Lett. 89 082301