Measurement of the tensor analyzing power T_{20} in the dd \rightarrow ³Hen and dd \rightarrow ³Hp at intermediate energies and at zero degree.

V.P. Ladygin^{h,†}, T. Uesaka^a, T. Saito^b, M. Hatano^b, A.Yu. Isupov^h, H. Kato^b,

N.B. Ladygina^h, Y. Maeda^a, A.I. Malakhov^h, J. Nishikawa^d, T. Ohnishi^c, H. Okamura^e,

S.G. Reznikov^h, H. Sakai^{a,b}, N. Sakamoto^c, S. Sakoda^b, Y. Satou^f, K. Sekiguchi^c,

K. Suda^a, A. Tamii^f, N. Uchigashima^b and K. Yako^b

^a Center for Nuclear Study, University of Tokyo, Bunkyo, Tokyo 113-0033, Japan

^b Department of Physics, University of Tokyo, Bunkyo, Tokyo 113-0033, Japan

^c RIKEN, Wako, Saitama 351-0198, Japan

^d Department of Physics, Saitama University, Urawa 338-8570, Japan

^e CYRIC, Tohoku University, Sendai, Miyagi 980-8578, Japan

^f Research Center for Nuclear Physics, Osaka University, Ibaraki 567-0047, Japan

^g Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan

^h LHE-JINR, 141980, Dubna, Moscow region, Russia

Abstract

The data on the tensor analyzing power T_{20} in the $dd \rightarrow {}^{3}\text{He}n$ and $dd \rightarrow {}^{3}\text{H}p$ reactions at 140, 200 and 270 MeV of the deuteron kinetic energy and at zero degree obtained at RIKEN Accelerator Research Facility are presented. The observed positive sign of T_{20} clearly demonstrates the sensitivity to the D/S wave ratios in the ${}^{3}\text{He}$ and ${}^{3}\text{H}$ in the energy domain of the measurements. The T_{20} data for the ${}^{3}\text{He}-n$ and ${}^{3}\text{H}-p$ channels are in agreement within experimental accuracy.

PACS: 24.70.+s; 21.45.+v

Keywords: tensor analyzing power; spin structure; deuteron; triton; ³He

[†] - Corresponding author:
Dr. V.P. Ladygin,
LHE-JINR,
Joliot Curie 6,
141980, Dubna, Moscow region, Russia
phone: 7-09621-63929
FAX : 7-09621-65180
E-mail address: ladygin@sunhe.jinr.ru

The spin structure of the light nuclei has been extensively investigated during the last decades using both electromagnetic and hadronic probes. The main purposes of these studies at intermediate and high energies are to obtain the information on the high-momentum components of light nuclei affected by the relativistic effects and to search the manifestation of non-nucleonic degrees of freedom. Especially, three nucleon bound states are of interest, because even a fundamental constant as the binding energy in the system cannot be reproduced by calculations with modern pairwise nucleon-nucleon potentials [1]. Since the binding energy is known to have a strong correlation with strength of spin-dependent forces such as tensor force and/or three-nucleon force, experimental study of the spin structure of three-nucleon bound system is expected to provide a clue to understand the source of the missing energy.

The non-relativistic Faddeev calculations [2] for three-nucleon bound state predict that the dominant components of the ³He ground state are a symmetric S-state, where the ³He spin due to the neutron and two protons are in a spin singlet state and a D-state, where all three nucleon spins are oriented opposite to the ³He spin. The S-state is found to dominate at small momenta while D-state dominates at large momenta. The relative sign of D- and S- wave in the momentum space is positive at small and moderate nucleon momenta [3].

The sensitivity to the different components of ${}^{3}\text{He}$ can be observed in polarization observables in both hadronic and electromagnetic processes.

Polarized electron scattering on polarized ³He target, ³He(\vec{e}, e')X, can be used to study the different components of the ³He wave function [2]. However, it is necessary to take into account final state interaction (FSI) and meson exchange currents (MEC) in addition to the plane wave impulse approximation (PWIA) to describe the experimental results [4] obtained at different relative orientations of electron and ³He spins. Recent CEBAF data on the transverse asymmetry $A_{T'}$ [5] at Q^2 values of 0.1 and 0.2 GeV/c² have been described using full Faddeev calculation with the MEC effects.

The ${}^{3}\text{He}(\vec{p}, 2p)$ and ${}^{3}\text{He}(\vec{p}, pn)$ breakup reactions were studied at TRIUMF in quasielastic kinematics at 200 [6] and 290 MeV [7] of incident proton energy. In the last experiment, spin observables A_{on} , A_{on} and A_{nn} were measured up to $q \sim 190$ and ~ 80 MeV/c for ${}^{3}\vec{\text{He}}(\vec{p}, 2p)$ and ${}^{3}\vec{\text{He}}(\vec{p}, pn)$ reactions, respectively. The results indicate that analyzing powers A_{no} , A_{on} and A_{nn} are close to the PWIA calculations for the ${}^{3}\vec{\text{He}}(\vec{p}, 2p)$ reaction, while for the ${}^{3}\vec{\text{He}}(\vec{p}, pn)$ there is a strong disagreement with these predictions. The same observables were recently measured at 197 MeV at IUCF Cooler Ring [8] up to $q \sim 400$ MeV/c. It was observed that the polarization of the neutron and proton at zero nucleon momentum in ${}^{3}\text{He}$ are $P_n \sim 0.98$ and $P_p \sim -0.16$, respectively, that is in good agreement with the Faddeev calculations [2]. However, at higher momenta there is the discrepancy, which can be due to the uncertainty of the theoretical calculations, as well as to large rescattering effects.

One Nucleon Exchange (ONE) reactions, like dp \rightarrow pd, d³He \rightarrow p⁴He or d³He \rightarrow ³Hed, are the simplest processes with large momentum transfer and, therefore, can be used as an effective tool to investigate the structure of the deuteron and ³He at short distances. In the framework of the ONE approximation [9] the polarization observables of the above reactions are expressed in terms of the D/S- wave ratios of these nuclei. For instance, tensor analyzing power T_{20} for the dp \rightarrow pd reaction in the collinear geometry is expressed in terms of D and S wave of the deuteron in the momentum space

$$T_{20} = \frac{2\sqrt{2}r - r^2}{\sqrt{2}(1+r^2)},\tag{1}$$

where r is the D/S- wave ratio in the deuteron at the corresponding internal momentum [9].

A significant amount of the data devoted to the investigation of the deuteron and ${}^{3}\text{He}({}^{3}\text{H})$ spin structure at short distances have been accumulated in the last years. Recently, the tensor analyzing power T_{20} and polarization transfer coefficients in backward elastic scattering, dp \rightarrow pd, have been measured at Saclay, Dubna and RIKEN [10, 11, 12, 13]. Another binary reaction, d³He \rightarrow p⁴He, has been investigated at RIKEN using both polarized deuteron and ³He up to 270 MeV [14, 15, 16]. All the data show the sensitivity to the deuteron spin structure at short distances. For instance, T_{20} for both dp \rightarrow pd [10, 12, 13] and d³He \rightarrow p⁴He [15, 16] reactions at intermediate energies has a large negative value reflecting the negative sign of the D/S- wave ratio in the deuteron in the momentum space.

As concerns ³He spin structure, the tensor analyzing power T_{20} in the d^3 He backward elastic scattering has been measured at 140, 200 and 270 MeV [17]. The sign of T_{20} is found to be positive in accordance with the positive sign of D/S- wave ratio in the ³He [3].

The data sensitive to the three-nucleon bound state spin structure are scarce, and, new polarization data, especially, at short distances is of great importance.

The $dd \rightarrow {}^{3}\text{H}p({}^{3}\text{H}en)$ process is the simplest ONE reaction where three nucleon structure is relevant. The theoretical analysis of the polarization phenomena for this reaction in the collinear geometry has been performed [18, 19]. It has been shown that the tensor analyzing power T_{20} due to polarization of the incident deuteron can be expressed in the terms of the D/S- wave ratio in the ${}^{3}\text{H}({}^{3}\text{He})$, when three-nucleon bound state is emitted in the forward direction in the cms. A new experiment has been proposed [20] to measure the energy and angular dependence of the tensor analyzing powers in the $dd \rightarrow {}^{3}\text{H}p({}^{3}\text{He}n)$ process at RIKEN.

In this paper the data on the tensor analyzing power T_{20} due to the incident deuteron polarization in the $dd \rightarrow {}^{3}\text{He}(0^{\circ})n$ and $dd \rightarrow {}^{3}\text{H}(0^{\circ})p$ reactions at 140, 200 and 270 MeV of the deuteron kinetic energy are presented.

The experiment has been performed at RIKEN Accelerator Research Facility (RARF). A polarized deuteron beam was produced by the high-intensity polarized ion source (PIS) [21] and accelerated by AVF and Ring cyclotrons up to the required energy. The direction of the beam polarization axis was controlled with a Wien filter located at the exit of the PIS. The polarization of the beam has been measured with the beam-line polarimeters based on the asymmetry measurements in dp- elastic scattering reaction, which has large values of tensor and vector analyzing powers [22, 12]. New values of the analyzing powers for dp- elastic scattering at 140 and 270 MeV obtained during absolute calibration of the deuteron beam polarization via the ${}^{12}C(d, \alpha){}^{10}B^*[2^+]$ reaction [23] have been used.

SMART (Swinger and Magnetic Analyzer with a Rotator and a Twister) spectrograph [24] has been used for the measurements. In this system, the incident beam can be swung by rotating the swinger magnet, while the magnetic analyzer is fixed. The measurement of the particle's momentum and separation from the primary beam was achieved by the

magnetic system of SMART spectrograph consisting of two dipole and three quadrupole magnets (Q-Q-D-Q-D).

The beam intensity during the experiment was about 1-2 nA, the solid angle was 10^{-2} sr. The live time of DAQ system [25] was more than 80% at the trigger rate of few thousands per second. A CD₂ thin sheet [26] has been used as a deuterium target. The measurements on CD₂ and carbon targets were made for each setup setting to obtain the contribution from deuterium via the CD₂-C subtraction. The thicknesses of the CD₂ and carbon targets used were 54 mg/cm² and 34 mg/cm², respectively.

Three plastic scintillators viewed by photomultipliers from the both sides were used in coincidence for the trigger and to provide the information about time-of-flight and energy losses of the particles. The distance between the target and detection point is about 17 m, which is enough to separate tritons, deuterons and protons with the same momentum. The time difference between the appearence of the trigger signal and radio-frequency signal of cyclotron was used as the time-of-flight information (TOF). The information about the charge of the particles was used at the trigger level. Protons and deuterons were partly suppressed in the cases of ³He and ³H detection by rasing threshold levels of the discriminators.

The event was used in the following analysis only in the case when information on the pulse height in all tree scintillation counters was consistent with the energy losses for particle of interest.

The set of multiwire drift chambers has been used to obtain the information on the hit position in the focal plane and, therefore, on the momentum of the particle and emission angle from the target. The energy resolution provided by the tracking system was ~ 300 keV. The typical track reconstruction efficiency was better than 99%.

In the experiment the data for ³He-*n* channels have been obtained at 140, 200 and 270 MeV, while for the ³H-*p* channel at 140 and 200 MeV only, because the momentum of the ³H at 270 MeV is higher than the maximal rigidity of SMART [24].

The quality of the CD₂-C subtraction procedure for the $dd \rightarrow {}^{3}\text{He}(0^{\circ})n$ reaction at 270, 200 and 140 MeV is demonstrated in Fig.1 a), b) and c), respectively. The spectra are plotted versus excitation energy E_X defined as following

$$E_X = \sqrt{((E_0 - E_{3N})^2 - (\mathbf{P}_0 - \mathbf{P}_{3N})^2 - M_N,}$$
(2)

where \mathbf{P}_0 is the incident momentum; $E_0 = 2M_d + T_d$ is the total initial energy; E_{3N} and \mathbf{P}_{3N} are energy and momentum of the three-nucleon system, respectively; M_N is the nucleon mass. Peaks at $E_X = 0$ MeV correspond to ³He from the $dd \rightarrow {}^{3}$ Hen reaction, while the shadowed areas show the carbon contribution.

For the ³H detection case the yield from carbon under peak at $E_X = 0$ MeV is negligibly small. The peak for the binary reaction on deuterium, $dd \rightarrow {}^{3}\text{H}p$, is separated from the peaks for the reaction $d^{12}C \rightarrow {}^{3}\text{H}X$ by ~ 5 and ~ 10 MeV at 200 and 140 MeV, respectively.

The events considered for the analysis were selected within polar angle acceptance $\leq 1.4^{\circ}$.

Only the events for unpolarized spin mode and two spin modes with the ideal values of the tensor polarization $p_{zz} = -2$ and $p_{zz} = +1$ [21] were used for the analysis. Actual values of the beam polarization were $\sim 75\%$, $\sim 50\%$ and $\sim 15\%$ of the ideal values at 270 MeV, 200 MeV and 140 MeV, respectively.

Tensor analyzing power T_{20} for each polarized spin mode was calculated from the following expression

$$T_{20} = -\frac{2\sqrt{2}}{p_{ii}} \left(\frac{\sigma_{pol}}{\sigma_0} - 1\right),$$
(3)

where p_{ii} is the corresponding tensor polarization of the beam, σ_{pol} and σ_0 are yields in polarized and unpolarized spin modes obtained via CD₂-C subtraction after corrections of dead-time, detection efficiency and beam intensity. The analyzing power T_{20} was taken as weighted averaged for two spin modes.

The results on the tensor analyzing power T_{20} for the $dd \rightarrow {}^{3}\text{He}(0^{\circ})n$ and $dd \rightarrow {}^{3}\text{H}(0^{\circ})p$ reactions are given in the Table 1. The systematic error due to uncertainty in the beam polarization and statistical error are added in quadrature.

These data are also shown in Fig.2. The open triangles and full circles correspond to the ³H-p and ³He-n channels, respectively. The T_{20} values obtained for the both charge symmetrical ³H-p and ³He-n channels at 140 and 200 MeV in this experiment are in good agreement within achieved experimental accuracy. No evidence of possible charge symmetry breaking is found in these processes.

The positive sign of T_{20} values is in a striking contrast to negative T_{20} for dp \rightarrow pd or other reactions where deuteron structure is relevant. The energy dependence of T_{20} demonstrates the increasing of T_{20} magnitude with the energy. This behaviour can be understood in terms of D/S wave ratio in the ³He(³H) with the help of ONE.

Within ONE approximation tensor analyzing power T_{20} for the $dd \rightarrow {}^{3}\text{H}p({}^{3}\text{He}n)$ reaction in the collinear geometry is expressed in terms of D and S wave ratio r of the ${}^{3}\text{H}({}^{3}\text{He})$ [18, 19, 20] (see expression (1)). The positive sign of T_{20} in the explored energy domain reflects the positive sign of the D/S wave ratio in the ${}^{3}\text{He}({}^{3}\text{H})$ in the momentum space [2, 3]. In this respect one can conclude that our data are sensitive to the D-state in the ${}^{3}\text{He}({}^{3}\text{H})$.

The solid, dashed and dotted curves in Fig. 2 are the results of non-relativistic ONE calculations [20] using Urbana [27], Paris [28] and RSC [29] (with the parametrization from [30]) ³He wave functions. Paris deuteron wave function [31] was used in the calculations. The data are in a qualitative agreement with the ONE calculations [20].

The behaviour of our data is consistent with the behaviour of T_{20} for other reactions sensitive to the ³He spin structure. In Fig.3 the data on T_{20} in the $dd \rightarrow {}^{3}\text{He}(0^{\circ})n$ are plotted versus internal momentum k along with the data obtained for the $d{}^{3}\text{He} \rightarrow {}^{3}\text{He}d$ reaction [17]. The data for the both processes demonstrate the universality in the kbehaviour. The solid curve is the result of ONE calculation using Urbana ³He wave function [27] according to Eq.(1). The relativistic effects are taken into account by the minimal relativization scheme [32].

The discrepancy between the data and the calculations can be due both to the contribution from the reaction mechanism other than ONE and to the nonadequate description of the short-range ³He spin structure. Concerning the reaction mechanism, the virtual excitation to the other channels, for example, excitation to Δ -isobar, is considered phenomenologically in Ref. [17] to reproduce the energy dependence of T_{20} for the $d^{3}\text{He} \rightarrow {}^{3}\text{He}d$ process. The microscopic calculation by Laget et al.[28] shows that the Δ -isobar contribution to the $dd \rightarrow {}^{3}\text{He}n$ process is less than 10% at energies lower than 300 MeV. At higher energies, in GeV region, it is shown that the coherent sum of ONE and the Δ -isobar excitation reproduces the cross section data reasonably. Thus it is expected that the uncertainty in the reaction mechanism is too small to explain the descrepancy between the data and the ONE calculation. Further theoretical investigations of the short-range ³He spin structure as well as the reaction mechanism of the $dd \rightarrow {}^{3}\text{He}n$ process are needed to understand the T_{20} data presented here.

The extension of the T_{20} measurements to the higher energies, namely to larger internal momenta, is of great interest. In particular, the measurement of T_{20} in the vicinity of $k \sim 0.4 \text{ GeV/c}$, where the changing of the T_{20} sign is expected, could distinguish different models of the short range ³He spin structure description.

The results can be summarized as following.

The data on the tensor analyzing power T_{20} in the $dd \rightarrow {}^{3}\text{H}p$ and $dd \rightarrow {}^{3}\text{H}en$ reactions at intermediate energies and in collinear geometry are obtained. The sign of T_{20} is positive being in the agreement with the results on T_{20} in the $d^{3}\text{He} \rightarrow {}^{3}\text{He}d$ reaction [17], on the one hand, and with the ONE calculations using standard ${}^{3}\text{He}$ wave functions, on the other hand.

In general, ONE reproduces the global feature of the T_{20} energy dependence. However, the deviation of the experimental data from the ONE calculations can be due to not only the nonadequate description of the short range ³He spin structure, but also the influence of the mechanisms additional to ONE. To improve the description of the obtained data further theoretical calculations are required.

The extension of T_{20} measurements to a GeV energy range is important to explore the short range ³He spin structure.

The authors express their thanks to the staff of RARF for providing of excellent conditions for the R308n experiment. They are grateful to H.Kumasaka, R.Suzuki and R.Taki for the help during experiment. Russian part of collaboration thanks RIKEN Directorate for kind hospitality during the experiment. The work has been supported in part by the JINR-Bulgaria grants for 2001 and 2002 years and by the Russian Foundation for Fundamental Research (grant N^o 04-02-17107).

References

- [1] W. Glöckle, H. Witala, D. Hüber, H. Kamada, J. Golak, Phys.Rep. 274 (1996) 107.
- B. Blankleider, R.M. Woloshyn, Phys.Rev. C 29 (1984) 538; J.L. Friar, B.F. Gibson, G.L. Payne, A.M. Bernstein, T.E. Chupp, Phys.Rev. C 42 (1990) 2310; R.-W. Schulze, P.U. Sauer, Phys.Rev. C 48 (1993) 38.
- [3] A.M. Eiro, F.D. Santos, J.Phys.G: Nucl.Phys. 16 (1990) 1139.
- [4] C.E. Woodward, et al., Phys.Rev.Lett.65 (1990) 698; C.E. Jones-Woodward, et al., Phys.Rev. C 44 (1991) R571; A.K. Thompson, et al., Phys.Rev.Lett. 68 (1992) 2901; M. Meyerhoff, et.al., Phys.Lett. B 327 (1994) 201.
- [5] W. Xu, et al., Phys. Rev. Lett. 85 (2000) 2900.
- [6] E.J. Brash, et al., Phys.Rev. C 47 (1993) 2064.

- [7] A. Rahav, et al, Phys.Lett. B 275 (1992) 259;
 A. Rahav, et al., Phys.Rev. C 46 (1992) 1167.
- [8] M.A. Miller, et al, Phys.Rev.Lett. 74 (1995) 502.
- [9] S.S. Vasan, Phys.Rev. D8 (1973) 4092; V.A. Karmanov, Yad.Fiz.34 (1981) 1020.
- [10] V. Punjabi, et al., Phys.Lett. B 350 (1995) 178.
- [11] L.S. Azhgirey, et al., Phys.Lett. B 391 (1997) 22; Yad.Fiz. 61 (1998) 494.
- [12] K. Sekiguchi, et al., Phys.Rev. C 65 (2002) 034003.
- [13] H. Sakai, et al., Phys.Rev.Lett. 84 (2000) 5288.
- [14] T. Uesaka, et al., Phys.Lett. B 467 (1999) 199.
- [15] T. Uesaka, et al., Few-Body Systems Suppl. 12 (2000) 497.
- [16] T. Uesaka, et al., Phys.Lett. B 533 (2002) 1.
- [17] M. Tanifuji, et al., Phys.Rev. C61 (2000) 024602.
- [18] V.P. Ladygin, N.B. Ladygina, Phys.Atom.Nucl. 59 (1996) 789.
- [19] V.P. Ladygin, N.B. Ladygina, Nuovo Cim. A 112 (1999) 855.
- [20] V.P. Ladygin, N.B. Ladygina, H. Sakai, T. Uesaka, Part.and Nucl.Lett. 3[100]-2000 (2000) 74.
- [21] H. Okamura, et al., AIP Conf. Proc. 293 (1993) 84.
- [22] N. Sakamoto, et al., Phys.Lett. B 367 (1996) 60.
- [23] K. Suda, et al., AIP Conf. Proc. 570 (2001) 806;
 RIKEN Accel.Prog.Rep. 35 (2002) 174.
- [24] T. Ichihara, et al., Nucl. Phys. A569 (1994) 287c.
- [25] H. Okamura, Nucl.Instr. and Meth. in Phys.Res. A 443 (2000) 194.
- [26] Y. Maeda, H. Sakai, K. Hatanaka, A. Tamii, Nucl.Instr. and Meth. in Phys.Res. A 490 (2002) 518.
- [27] R. Schiavilla, V.R. Pandharipande, R.B. Wiringa, Nucl. Phys. A 449 (1986) 219.
- [28] J.-M. Laget, J.F. Lecolley, F. Lefebvres, Nucl. Phys. A 370 (1981) 479.
- [29] F.D. Santos, A.M. Eiro, A. Barosso, Phys.Rev. C 19 (1979) 238.
- [30] Yu.N. Uzikov, EChAYa 29 (1998) 1010.
- [31] M. Lacombe, B. Loiseau, R. Vinh Mau, J. Cote, P. Pires, R. de Tourreil, Phys.Lett. B 101 (1981) 139.
- [32] P.A.M. Dirac, Rev.Mod.Phys. 21 (1949) 392; S. Weinberg, Phys.Rev. 150 (1966) 1313; L.L. Frankfurt, M.I. Strikman, Phys.Rep. 76 (1981) 215.

Energy, MeV	Reaction	T_{20}	ΔT_{20}
140	$dd \rightarrow {}^{3}Hen$	0.112	0.019
140	$\mathrm{dd} \to {}^{3}\mathrm{Hp}$	0.082	0.018
200	$dd \rightarrow {}^{3}Hen$	0.172	0.020
200	$\mathrm{dd} \to {}^{3}\mathrm{Hp}$	0.165	0.018
270	$dd \rightarrow {}^{3}Hen$	0.300	0.013

Table 1. Tensor analyzing power T_{20} in the dd \rightarrow ³Hen and dd \rightarrow ³Hp reactions.

Figures caption

Fig.1. $CD_2 - C$ subtraction for the $dd \rightarrow^3 He(0^\circ)n$ channel: a) at 270 MeV, b) at 200 MeV and c) at 140 MeV. The open and shadowed histograms correspond to the yields from CD_2 and carbon targets, respectively.

Fig.2. Tensor analyzing power T_{20} in the dd \rightarrow ³Hen(solid circles) and dd \rightarrow ³Hp (open triangles) reactions in collinear geometry versus incident momentum of deuteron. The solid, dashed and dotted curves are the results of the non-relativistic ONE calculations [20] for the forward emission of the ³He(³H) in the cms, using Urbana [27], Paris [28] and RSC [29] ³He wave functions, respectively. Paris deuteron wave function [31] was used for the deuteron structure description.

Fig.3. Tensor analyzing power T_{20} in the dd \rightarrow ³Hen(solid circles) and d³He \rightarrow ³Hed (open squares) [17] reactions versus internal momentum k. The solid curve is the result of the ONE calculations using Urbana ³He wave function [27].



Figure 1:



Figure 2:



Figure 3: