# The <sup>18</sup>F(p,α)<sup>15</sup>O reaction rate for application to nova γ-ray emission N. de Séréville<sup>a\*</sup>, E. Berthoumieux<sup>b</sup> and A. Coc<sup>a</sup> <sup>a</sup>CSNSM, CNRS/IN2P3/UPS, Bât. 104, 91405 Orsay Campus, France <sup>b</sup>CEA, DAPNIA/SPhN, F-91191 Gif/Yvette Cedex, France

The <sup>18</sup>F(p, $\alpha$ )<sup>15</sup>O reaction is recognized as one of the most important reaction for nova gamma-ray astronomy as it governs the early  $\leq 511$  keV emission. However, its rate remains largely uncertain at nova temperatures due to unknown low-energy resonance strengths. We report here on our last results concerning the study of the D(<sup>18</sup>F,p $\alpha$ )<sup>15</sup>N reaction, as well as on the determination of the <sup>18</sup>F(p, $\alpha$ )<sup>15</sup>O reaction rate using the Rmatrix theory. Remaining uncertainties are discussed.

# 1. INTRODUCTION

Gamma-ray emission from classical novae is dominated, during the first hours, by positron annihilation following the beta decay of <sup>18</sup>F[1, 2, 3]. However, even though it has been the object of many recent experiments[4, 5, 6] the rate of its main mode of destruction, through the <sup>18</sup>F(p, $\alpha$ )<sup>15</sup>O reaction, remains highly uncertain. This was mainly due to the unknown proton widths of the first three <sup>19</sup>Ne levels above the proton emission threshold ( $E_x$ ,  $J^{\pi} = 6.419$  MeV,  $3/2^+$ ; 6.437 MeV,  $1/2^-$  and 6.449 MeV,  $3/2^+$ ). The tails of the corresponding resonances (at respectively,  $E_R = 8$ , 26 and 38 keV) can dominate the astrophysical S-factor in the relevant energy range[3]. As a consequence of these nuclear uncertainties, the <sup>18</sup>F production in nova and the early gamma-ray emission was uncertain by a factor of  $\approx 300[3]$ . Since a direct measurement of the relevant resonance strengths is impossible due to the very low Coulomb barrier penetrability, we used an indirect method aiming at determining the one-nucleon spectroscopic factors in the analog levels of the mirror nucleus (<sup>19</sup>F) by the neutron transfer reaction D(<sup>18</sup>F,p)<sup>19</sup>F[7]. Recently, the same reaction has been studied at higher energy[8]. We present here additional information on the extraction of spectroscopic factors as well as new <sup>18</sup>F(p, $\alpha$ )<sup>15</sup>O reaction rates.

## 2. EXPERIMENT AND ANALYSIS

We refer to de Séréville et al. [7] for experimental details and present in Figure 1 the reconstructed excitation energy spectrum. The well populated <sup>19</sup>F levels at  $E_X =$ 2.780 and 5.106 MeV observed in single events as well as the  $E_X =$  7.262 + 7.368 MeV group observed in coincidence events were used to make an internal calibration. This procedure resulted in an uncertainty on the excitation energy calibration of  $\approx 3$  keV in

<sup>\*</sup>Presently at IEEC, Edifici Nexus 201, C/ Gran Capita 2-4, 08034 Barcelona, Spain



Figure 1. <sup>19</sup>F excitation energy calibration spectrum with a global fit (see text). The  $E_X = 6.527$  MeV level is the most populated.

the region of interest. During this analysis a careful study of the systematic errors has been done. To extract the relative contribution of the two  $3/2^+$  levels, a simultaneous fit of the 6.5 MeV group ( $E_X = 6.497 + 6.528$  MeV), the 6.25 MeV group ( $E_X = 6.255$ + 6.282 + 6.330 MeV) and the 6.9 MeV group ( $E_X = 6.787 + 6.838 + 6.891$  MeV) was performed. The background is described by a lorentzian of width  $\Gamma = 280 \text{ keV}$ corresponding to the  $1/2^- E_X = 6.429$  MeV. The result favors the dominant contribution of one single level ( $E_X = 6.527$  MeV) while another more recent study [8] favors the other one. In any case, the nuclear structure of these two  $3/2^+$  levels seem to be very different according to the results of an inelastic electron scattering measurement on <sup>19</sup>F [ 9]. From the angular information of the 6.5 MeV peak, we have obtained the angular distribution that we have analyzed performing a finite range DWBA analysis (FRESCO) 10] code), including a compound nucleus component. The extracted spectroscopic factor is  $C^2S = 0.17$  (0.21 when neglecting the compound nucleus contribution 7) and is weakly dependent of the optical potential parameters. It is to be noted that our experiment has been recently repeated at higher energy [8] and that a slightly lower spectroscopic factor is found  $(C^2 S = 0.12)$ .

# 3. REMAINING UNCERTAINTIES AND REACTION RATE

Before deriving a new  ${}^{18}F(p,\alpha){}^{15}O$  reaction rate and because the previous analysis showed that the  $3/2^+$  levels of interest *cannot* be neglected, it seems important to focus

on the remaining uncertainties. First, the spectroscopic factors obtained correspond to <sup>19</sup>F levels and should be transposed to the analog <sup>19</sup>Ne levels in order to deduced the proton widths used to calculate the reaction rate. Such a common practice in nuclear astrophysics leads to an uncertainty of a factor of about two on the proton widths (based on a statistical study of analog states in the same mass region). Moreover, due to the small separation energy of this doublet (only 30 keV), the assignation of analog levels is not very clear and in the following we will always deal with two cases: no inversion where the low energy level in  ${}^{19}$ F is the analog of the low energy one in  ${}^{19}$ Ne and *inversion* where the low energy level in  $^{19}$ F is the analog of the high energy one in  $^{19}$ Ne. Second, the  $\alpha$ -widths of the <sup>19</sup>Ne levels are unknown and calculated from the reduced widths of the corresponding analog levels in <sup>19</sup>F. Unlike the one-nucleon case (see above), the associated uncertainty could be as large as a factor of 10 [11]. Third, the two  $3/2^+$ levels of astrophysical interest can interfere with another  $3/2^+$  level at  $E_X = 7.076$  MeV and no information about the sign of interferences is available at present day. In case of destructive interference, the astrophysical S-factor decreases drastically in the Gamow region at nova temperature.

Figure 2 is an example of the influence on the astrophysical S-factor of the  $\alpha$ -width variation when constructive and destructive interferences are considered between the two levels  $E_X = 6.449$  and 7.076 MeV (no inversion). Experimental data from Bardayan et al.[6] are also displayed. These data points are used to constrain the astrophysical S-factor with the help of R-matrix fits (ANARKI [12] code). The free parameters are the  $\alpha$ -width of the  $E_r = 8$  or 38 keV resonances, the proton width of the  $E_r = 330$  keV, the proton and  $\alpha$ -width and position of the  $E_r = 665$  keV resonance and the sign of the interferences. The best fit is obtained for constructive interferences for the  $E_X = 6.449$  MeV level and is then used for determining the new  ${}^{18}\text{F}(p,\alpha){}^{15}\text{O}$  nominal rate. The upper reaction rate is given for a constructive interference for the  $E_X = 6.419$  MeV (inversion) and an  $\alpha$ -width such as the astrophysical S-factor correspond to the upper limit of the error bar of the low-energy data point from [6] ( $E_{c.m.} = 375$  keV). In the same way, the lower rate is given for a destructive interference for the  $E_X = 6.449$  MeV (no inversion) and an  $\alpha$ -width such as the astrophysical S-factor correspond to the upper limit of the error bar of the low-energy data point from [6] ( $E_{c.m.} = 375$  keV). In the same way, the lower rate is given for a destructive interference for the  $E_X = 6.449$  MeV (no inversion) and an  $\alpha$ -width such as the astrophysical S-factor correspond to the lower limit of the error bar at  $E_{c.m.} = 375$  keV.

Since we consider that the inversion of the analog levels is a possibility, the conclusions are the following. The new nominal rate is within a factor of two of the former one [3]. Furthermore the upper rate is reduced and the global uncertainty is reduced but remains large.

# 4. ACKNOWLEDGMENTS

This work is based on experimental results obtained in collaboration with C. Angulo, M. Assunção, D. Beaumel, B. Bouzid, S. Cherubini, M. Couder, P. Demaret, F. de Oliveira Santos, P. Figuera, S. Fortier, M. Gaelens, F. Hammache, J. Kiener, D. Labar, A. Lefebvre–Schuhl, P. Leleux, M. Loiselet, A. Ninane, S. Ouichaoui, G. Ryckewaert, N. Smirnova, V. Tatischeff and J.-P. Thibaud.

This work has been supported by the European Community-Access to Research Infrastructure action of the Improving Human Potential Programme, contract N<sup>o</sup> HPRI-CT-



Figure 2. Astrophysical S-factor for constructive and destructive interferences between the two <sup>19</sup>Ne levels  $E_X = 6.449$  and 7.076 MeV (dashed lines). Cases for different  $\alpha$ width are reported. (The 6.449 MeV spectroscopic factor is from our experiment while data points are from Bardayan et al.[6].). Solid lines represent the astrophysical S-factor used for the lower and upper <sup>18</sup>F(p, $\alpha$ )<sup>15</sup>O reaction rate (see text). Gamow windows for temperatures relevant to nova are shown.

1999-00110.

### REFERENCES

- 1. J. Gómez–Gomar, et al., MNRAS 296, 913 (1998).
- 2. M. Hernanz, et al., Astrophys. J. 526, L97 (1999).
- 3. A. Coc, et al., Astron. Astrophys. 357, 561 (2000).
- 4. J.-S. Graulich, et al., *Phys. Rev.* C63, 011302(R) (2000).
- 5. D.W. Bardayan, et al., *Phys. Rev.* C63, 065802 (2001).
- 6. D.W. Bardayan, et al., *Phys. Rev. Lett.* **89**, 262501 (2002).
- 7. N. de Séréville, et al., *Phys. Rev.* C67, 052801 (R) (2003).
- 8. R. L. Kozub et al., see contribution in these proceedings.
- 9. B. A. Brown et al., *Phys. Rev.* C32, 1127 (1985).
- 10. I. J. Thompson Computer Physics Report 7, 167 (1988).
- 11. F. de Oliveira et al., *Phys. Rev.* C55, 3149 (1997).

12. E. Berthoumieux et al., Nucl. Inst. and Meth. **B136**, 55 (1998).