

YAG:Nd crystals as possible detector to search for 2β and α decay of neodymium

F.A. Danevich¹, V.V. Kobychiev, S.S. Nagorny, V.I. Tretyak

Institute for Nuclear Research, MSP 03680 Kiev, Ukraine

Abstract

Energy resolution, α/β ratio, pulse-shape discrimination for γ rays and α particles, radioactive contamination were studied with neodymium doped yttrium-aluminum garnet (YAG:Nd). Applicability of YAG:Nd scintillators to search for 2β decay and α activity of natural neodymium isotopes are discussed.

PACS numbers: 29.40.Mc; 23.40.-s; 23.60.+e; 95.35.+d

Keywords: Scintillation detector, Double beta decay, Alpha decay, Dark matter, Neodymium doped yttrium-aluminum garnet

1 INTRODUCTION

The great interest to the double beta (2β) decay search is related, in particular, with the recent evidence of neutrino oscillations which strongly suggests that neutrinos have nonzero mass [1, 2, 3, 4]. While oscillation experiments are sensitive to the neutrinos mass difference, only the measured neutrinoless (0ν) double beta decay rate could set the Majorana nature of the neutrinos and give the absolute scale of the effective neutrino mass.

^{150}Nd is one of the most promising candidate for 2β decay study due to its large transition energy ($Q_{2\beta}$). As a result, the calculated value of the phase space integral $G_{mm}^{0\nu}$ of the $0\nu 2\beta$ decay of ^{150}Nd is the largest one among 35 possible $2\beta^-$ decay candidates [5, 6]. The theoretical predictions of the product $T_{1/2}^{0\nu} \cdot \langle m_\nu \rangle^2$ are in the interval $3.4 \times 10^{22} - 3.4 \times 10^{24}$ yr (half-lives for the neutrino mass mechanism with $\langle m_\nu \rangle = 1$ eV) [7]. Moreover, the larger the $Q_{2\beta}$ energy, the simpler, from an experimental point of view, is to overcome background problems. Note that the background from natural radioactivity drops sharply above 2615 keV, which is the energy of the γ from ^{208}Tl decay (^{232}Th family). In addition, a contribution of cosmogenic activation, which is important problem of the next generation 2β decay experiments [8], decreases at higher energies.

There exist no appropriate detector containing neodymium, which could serve as both source and detector simultaneously (so called "active" source experimental method [2]). It should be noted that this experimental approach will assure high registration efficiency which is especially important if very expensive enriched ^{150}Nd isotope would be used in an experiment.

The purpose of our work was investigation of scintillation properties and preliminary check of radioactive contamination of neodymium doped yttrium-aluminum garnet (YAG:Nd) as a possible detector for double beta decay experiment with ^{150}Nd . The active source method allows

¹Corresponding author. *E-mail address:* danevich@kinr.kiev.ua

Table 1: Properties of YAG:Nd crystal scintillators

Density (g/cm ³)	4.56
Melting point	1970°C
Crystal structure	Cubic Garnet
Hardness (Mohs)	8.5
Refractive index	1.82
Average decay time*	4 μ s
Photoelectron yield relatively to NaI(Tl)*	8%

*For γ rays, at room temperature.

also to investigate other rare processes in Nd nuclei, such as α decay of naturally occurring neodymium isotopes.

2 MEASUREMENTS AND RESULTS

2.1 Energy resolution and α/β ratio

Main properties of YAG:Nd (chemical formula $Y_3Al_5O_{12}:Nd$) are presented in Table 1. These crystals are well known and their production is well developed due to their wide application in laser technique. The scintillation properties of cerium doped yttrium-aluminum garnet (YAG:Ce) have been studied in [9], properties of YAG:Yb have been reported in [10], whereas YAG:Nd, to our knowledge, was never studied as a scintillator.

All the measurements were carried out for the $\varnothing 17 \times 6$ mm YAG:Nd crystal with ≈ 2 mol% of Nd. The mass of the crystal is 7.16 g. Photoelectron yield was estimated with the Philips XP2412 bialkali photomultiplier (PMT) as 8% of NaI(Tl). The energy resolution FWHM=13.6% was measured for 662 keV γ line of ¹³⁷Cs with the YAG:Nd crystal wrapped by PTFE reflector tape and optically coupled to the PMT XP2412. A substantial improvement of the light output ($\approx 22\%$) and energy resolution was achieved by placing the crystal in liquid (silicone oil with index of refraction ≈ 1.5). The crystal was fixed in center of the teflon container $\varnothing 70 \times 90$ mm and viewed by two PMTs XP2412. Fig. 1 demonstrates the energy spectra of ¹³⁷Cs and ²⁰⁷Bi obtained in such a way with the YAG:Nd crystal. The energy resolution of 9.3% was obtained with the YAG:Nd crystal scintillator for 662 keV γ line of ¹³⁷Cs.

We estimate the energy resolution of the YAG:Nd scintillation detector at the energy $Q_{2\beta}$ of ¹⁵⁰Nd using the results of measurements with ¹³⁷Cs and ²⁰⁷Bi γ sources. These data were fitted by function $FWHM(\text{keV}) = a + \sqrt{b \times E_\gamma}$ with values $a = 2$ keV and $b = 5.2$ keV, where energy of γ quanta E_γ is in keV. The energy resolution $FWHM \approx 4\%$ could be achieved at the $Q_{2\beta}$ energy of ¹⁵⁰Nd.

The α/β ratio was measured with the help of the collimated α particles of a ²⁴¹Am source. As it was checked by surface-barrier detector, the energy of α particles was reduced to about 5.25 MeV by ≈ 1 mm of air, due to passing through the collimator. Fig. 2 shows the energy spectrum of the α particles measured by the YAG:Nd scintillator. The α/β ratio is 0.33. The YAG:Nd crystal was irradiated in three perpendicular directions with aim to check a possible dependence of the α signal on direction of irradiation. While earlier such a dependence has




Figure 1: Energy spectra of ^{137}Cs (a) and ^{207}Bi (b) γ rays measured with the YAG:Nd scintillation crystal. The crystal was located in liquid and viewed by two distant PMTs (see text).

been found for CdWO_4 [11] and ZnWO_4 [12] scintillators, we did not observe it for YAG:Nd crystal.

2.2 Pulse-shape discrimination for γ quanta and α particles

The shapes of scintillation light pulses in YAG:Nd crystals were studied for 5.25 MeV α particles and $\approx 1.5 - 1.8$ MeV γ quanta with the help of a 12 bit 20 MHz transient digitizer as described in [11, 13]. The pulse shape can be fitted by sum of exponential functions:

$$f(t) = \sum A_i / (\tau_i - \tau_0) \cdot (e^{-t/\tau_i} - e^{-t/\tau_0}), \quad t > 0,$$

where A_i are intensities (in %), and τ_i are decay constants for different light emission components, τ_0 is integration constant of electronics ($\approx 0.2 \mu\text{s}$). The values of A_i and τ_i obtained by fitting the average of ≈ 3 thousand individual α and γ pulses in the time interval $0 - 60 \mu\text{s}$ (see Fig. 3) are presented in Table 2. Observed difference in light pulse shapes allows to discriminate $\gamma(\beta)$ events from α particles. We applied for this purpose the optimal filter method proposed in [14] and developed in [13]. To obtain the numerical characteristic of YAG:Nd signal, so called shape indicator (SI), the following formula was applied for each pulse:

$$SI = \sum f(t_k) \cdot P(t_k) / \sum f(t_k),$$




Figure 2: The energy spectrum of α particles from collimated ^{241}Am source.

where the sum is over time channels k , starting from the origin of pulse and up to $50 \mu\text{s}$, $f(t_k)$ is the digitized amplitude (at the time t_k) of a given signal. The weight function $P(t)$ is defined as: $P(t) = \{f_\alpha(t) - f_\gamma(t)\} / \{f_\alpha(t) + f_\gamma(t)\}$, where $f_\alpha(t)$ and $f_\gamma(t)$ are the reference pulse shapes for α particles and γ quanta. Reasonable discrimination between α particles and γ rays was achieved using this approach as one can see in Fig. 4 where the shape indicator distributions measured by the YAG:Nd scintillation crystal with α particles ($E_\alpha \approx 5.3 \text{ MeV}$) and γ quanta ($\approx 1.5 - 1.8 \text{ MeV}$) are shown.

2.2.1 Radioactive contamination of the YAG:Nd crystal

To determine activity of α active nuclides from U/Th contamination in the YAG:Nd crystal, the pulse shape analysis was applied to data of 14.8 h low background measurement carried out

Table 2: Decay time of YAG:Nd scintillator for γ quanta and α particles. The decay constants and intensities (in percentage of the total intensity) are denoted as τ_i and A_i , respectively.

Type of irradiation	Decay constants, μs		
	$\tau_1 (A_1)$	$\tau_2 (A_2)$	$\tau_3 (A_3)$
γ rays	1.1 (10%)	4.1 (87%)	20 (3%)
α particles	1.0 (15%)	3.9 (83%)	18 (2%)




Figure 3: Decay of scintillation in YAG:Nd for γ rays and α particles (≈ 3000 forms of γ and α events were added) and their fit by sum of three exponential components with decay constants $\tau_i \approx 1, \approx 4$, and $\approx 20 \mu\text{s}$.

at the earth surface laboratory in Kiev. The YAG:Nd crystal was viewed by the low-radioactive PMT (EMI D724KFLB) through plastic scintillator light-guide 10 cm in diameter and 4 cm long. The active light-guide reduces effect of γ radiation from the PMT and provides suppression of cosmic rays induced background due to the pulse-shape discrimination of scintillation signals. Passive shield was consisted of steel (2 cm) and lead (10 cm). The energy resolution of the detector was 11% for 662 keV γ rays.

To estimate contamination of the crystal by α active nuclides of U/Th families, the α events were selected from the data accumulated with the YAG:Nd crystal with the help of the pulse-shape discrimination technique. In the α spectrum there is no peculiarities which can be surely attributed to α active U/Th daughters. Analyzing these data we set the limit on total α activity of ^{235}U , ^{238}U , and ^{232}Th daughters in the YAG:Nd crystal: ≤ 20 mBq/kg.

3 DISCUSSION

3.1 2β decay of neodymium

Among neodymium isotopes there are three possible 2β candidates: ^{146}Nd , ^{148}Nd , and ^{150}Nd . They are listed in Table 3 where the $Q_{2\beta}$ energies and the isotopic abundances are given. As it was mentioned in the Introduction, the most interesting of them is ^{150}Nd due to the




Figure 4: The shape indicator (see text) distributions measured by YAG:Nd scintillation crystal with α particles ($E_\alpha = 5.25$ MeV) and γ quanta ($\approx 1.5 - 1.8$ MeV).

large value of the transition energy. Two neutrino 2β decay of ^{150}Nd with the half-life $T_{1/2} = 1.9_{-0.4}^{+0.7} \times 10^{19}$ yr was observed in experiment [17] by using a time projection chamber and samples of enriched ^{150}Nd and natural neodymium. In [18] the double beta decay of ^{150}Nd was studied with similar technique (only enriched ^{150}Nd sample was used). The two-neutrino half-life $T_{1/2} = (6.8 \pm 0.8) \times 10^{18}$ yr was measured. Recently, the observation of ^{150}Nd $2\nu 2\beta$ decay to the second excited level of ^{150}Sm (0_1^+ at 741 keV) with $T_{1/2} = 1.4_{-0.4}^{+0.5} \times 10^{20}$ yr was reported in [19], while only the limit $T_{1/2} > 1.5 \times 10^{20}$ yr was determined previously at 90% C.L. [20]. The most stringent limit on the neutrinoless mode $T_{1/2} \geq 1.2 \times 10^{21}$ yr was set in the experiment [18].

YAG:Nd scintillators provide a possibility to search for $0\nu 2\beta$ decay of ^{150}Nd by "active" source experimental method. The studied pulse-shape discrimination ability is an important

Table 3: 2β unstable neodymium isotopes

Transition	Isotopic abundance (%) [15]	$Q_{2\beta}$ (keV) [16]
$^{146}\text{Nd} \rightarrow ^{146}\text{Sm}$	17.2(0.3)	70.2(2.9)
$^{148}\text{Nd} \rightarrow ^{148}\text{Sm}$	5.7(0.1)	1928.8(1.9)
$^{150}\text{Nd} \rightarrow ^{150}\text{Sm}$	5.6(0.2)	3367.5(2.2)

Table 4: Theoretical calculations of half-lives for α decay of natural Nd isotopes. Uncertainties given for calculations with cluster model of Ref. [25] and semiempirical formula of Ref. [26] are related only with uncertainty in the Q_α values. For ^{143}Nd , we take into account that change in a parity and spin additionally suppresses the ^{143}Nd decay rate.

Isotope	Abund. [15]	Q_α , MeV [16]	Calculated $T_{1/2}$, yr			Exp. $T_{1/2}$, yr
			based on [25]	based on [26]	[27]	
^{143}Nd	12.2%	0.520(7)	$> 1.1_{-1.0}^{+7.8} \times 10^{80}$	$5.2_{-4.7}^{+51} \times 10^{93}$	–	–
^{144}Nd	23.8%	1.905(2)	$1.9_{-0.1}^{+0.2} \times 10^{15}$	$4.3_{-0.3}^{+0.3} \times 10^{15}$	–	$2.29 \pm 0.16 \times 10^{15}$
^{145}Nd	8.3%	1.578(2)	$1.7_{-0.1}^{+0.2} \times 10^{22}$	$3.9_{-0.4}^{+0.3} \times 10^{23}$	3.7×10^{22}	$> 1 \times 10^{17}$
^{146}Nd	17.2%	1.182(2)	$2.0_{-0.4}^{+0.4} \times 10^{34}$	$3.9_{-0.7}^{+0.8} \times 10^{34}$	–	–
^{148}Nd	5.7%	0.599(3)	$6.1_{-3.1}^{+6.7} \times 10^{70}$	$1.1_{-0.6}^{+1.2} \times 10^{71}$	–	–

advantage of YAG:Nd scintillators as a low counting rate detector. Background of scintillation detector caused by radioactive contamination by U/Th daughters can be effectively rejected due to pulse-shape analysis as it was demonstrated in experiments with CdWO_4 [11, 21] and CaWO_4 [8, 22] crystal scintillators. Obvious disadvantage of YAG:Nd detector is rather small concentration of Nd. In the studied crystal the mass concentration is only $\approx 0.03\%$ of ^{150}Nd . It should be noted, however, that YAG:Nd crystals can be produced with neodymium concentration up to 8 mol%, which corresponds to ≈ 0.1 mass% of ^{150}Nd . In this connection we would like to refer the project CANDLES [23] which intends to use a few tons of non-enriched CaF_2 crystals to search for $0\nu 2\beta$ decay of ^{48}Ca . The concentration of target ^{48}Ca nuclei in CaF_2 detector is at the same level: ≈ 0.1 mass% of ^{48}Ca .

The value of energy resolution $\text{FWHM}=4\%$ could allow to discover [24] the $0\nu 2\beta$ decay of ^{150}Nd with the half-life $T_{1/2}^{0\nu 2\beta} \sim 10^{25}$ yr, which corresponds to the effective Majorana neutrino mass $m_\nu \sim 0.06-0.6$ eV (taking into account all existing calculations of nuclear matrix elements for $0\nu 2\beta$ decay of ^{150}Nd). As for the sensitivity [24], an experiment involving ≈ 20 tons of non-enriched YAG:Nd crystals with ≈ 8 mol% of Nd could reach the half-life sensitivity $\approx 3 \times 10^{26}$ yr (supposing zero background during ten years of measurements), which corresponds to the Majorana neutrino mass $0.01 - 0.1$ eV.

3.2 α decay of neodymium

Alpha decay is allowed energetically for five naturally occurring isotopes of neodymium. Their energies of α decay (Q_α) and natural abundances are listed in Table 4. Calculation of half-life values based on models [25, 26], and the value of the half-life of ^{145}Nd calculated in [27] are also presented. Alpha decay was observed only for ^{144}Nd . Average half-life $T_{1/2} = 2.29 \pm 0.16 \times 10^{15}$ yr [28] was derived on the basis of four measurements: $T_{1/2} = 2.2 \times 10^{15}$ yr [29], $T_{1/2} = 1.9 \times 10^{15}$ yr [30], $T_{1/2} = 2.4 \pm 0.3 \times 10^{15}$ yr [31] and $T_{1/2} = 2.65 \pm 0.37 \times 10^{15}$ yr [32]. It is obvious that, due to low Q_α values and long expected $T_{1/2}$'s, there is no perspective to observe the α decay of ^{143}Nd , ^{146}Nd and ^{148}Nd . As for ^{145}Nd , only two $T_{1/2}$ limits are known: $T_{1/2} > 6 \times 10^{16}$ yr [30] and $T_{1/2} > 1 \times 10^{17}$ yr [33].

The indication on the α decay of natural ^{180}W isotope with half-life $T_{1/2} = 1.2_{-0.4}^{+0.8}$ (stat) ± 0.3 (syst) $\times 10^{18}$ yr has been recently observed in the experiment [11] with the help of the low background $^{116}\text{CdWO}_4$ crystal scintillators. This result was confirmed in the measurements with CaWO_4 crystal as scintillator [22] and bolometer [34]. It should be also referred the excellent result on the detection of alpha decay of ^{209}Bi with the half-life $T_{1/2} = (1.9 \pm 0.2) \times 10^{19}$ y [35]. In both the experiments [34] and [35] the cryogenic technique, which uses simultaneous registration of heat and light signals, have been applied. This method provides perfect selection of α events on the background caused by γ rays (electrons). It should be interesting to check YAG:Nd crystal as the cryogenic detector to search for alpha decay of ^{145}Nd and to measure more accurately the ^{144}Nd half-life.

It should be also mentioned a possibility to search for spin-dependent inelastic scattering of weakly interacting massive particles (WIMP) with excitation of low energy nuclear levels of ^{145}Nd (the lowest one: $3/2^-$ 67 keV, E2 transition, and the second level: $5/2^-$ 72 keV, M1 transition) due to nonzero spin of this nucleus ($7/2^-$). Identification of such "mixed" (nuclear recoil plus γ quanta) events could be possible due to the simultaneous registration of heat and light signals (such a techniques have been used in [36, 37]). A heat/light ratio for such events would differ from "pure" nuclear recoils or γ (electron) events.

4 CONCLUSIONS

The scintillation properties of YAG:Nd crystals were studied. The energy resolution 9.3% (662 keV ^{137}Cs γ line) was obtained with the YAG:Nd crystal scintillator placed in liquid and viewed by two PMTs. Shapes of scintillation signals were investigated, and reasonable pulse-shape discrimination for γ rays and α particles was achieved. The α/β ratio was measured with the YAG:Nd scintillator to be equal 0.33 at the energy of alpha particles 5.25 MeV. Radioactive contamination of the YAG:Nd crystal by α active nuclides from U/Th chains was estimated as ≤ 20 mBq/kg.

Three potentially 2β active neodymium isotopes can be studied with the help of YAG:Nd crystal. Due to good scintillation characteristics and pulse-shape discrimination ability the YAG:Nd scintillators seems to be encouraging material for the 2β experiment with ^{150}Nd .

The YAG:Nd crystals could be used to search for α decay of natural neodymium isotopes, in particular, ^{145}Nd . A strong signature of α decay can be assured via simultaneous registration of heat and light signals using cryogenic technique. However, ability of YAG:Nd crystal as cryogenic detector has to be elaborated in an experiment.

5 ACKNOWLEDGEMENT

The authors would like to thank Dr. Yu.D. Glinka from the Institute of Surface Chemistry of the National Academy of Sciences of Ukraine for lending the YAG:Nd crystal used in the present study.

References

- [1] J.D. Vergados, Phys. Rep. 361 (2002) 1.
- [2] Yu.G. Zdesenko, Rev. Mod. Phys. 74 (2002) 663.
- [3] S.R. Elliot and P. Vogel, Ann. Rev. Nucl. Part. Sci. 52 (2002) 115.
- [4] V.I. Tretyak and Yu.G. Zdesenko, At. Data Nucl. Data Tables 61 (1995) 43; 80 (2002) 83.
- [5] M. Doi et al., Prog. Theor. Phys. Suppl. 83 (1985) 1.
- [6] J. Suhonen and O. Civitarese, Phys. Rep. 300 (1998) 123.
- [7] M. Doi et al., Prog. Th. Phys. 69 (1983) 602;
A. Staudt et al., Europhys. Lett. 13 (1990) 31;
T. Tomoda, Rep. Prog. Phys. 54 (1991) 53;
M. Hirsh et al., Z. Phys. A 347 (1994) 151;
J.G. Hirsh et al., Nucl. Phys. A 582 (1995) 124;
F. Simkovic et al., Phys. Rev. C 60 (1999) 055502.
- [8] see, for instance: Yu.G. Zdesenko et al., "CARVEL: an experiment with $^{48}\text{CaWO}_4$ crystal scintillators for the double β decay study of ^{48}Ca ", submitted to Astropart. Phys.
- [9] M. Moszynski et al., Nucl. Instr. Meth. A 345 (1994) 461;
T. Ludziejewski et al., Nucl. Instr. Meth. A 398 (1997) 287;
T. Bhattacharjee et al., Nucl. Instr. Meth. A 484 (2002) 364.
- [10] P. Antonini et al., Nucl. Instr. Meth. A 460 (2001) 469.
- [11] F.A. Danevich et al., Phys. Rev. C 67 (2003) 014310.
- [12] F.A. Danevich, nucl-ex/0409014; submitted to Nucl. Instr. Meth. A.
- [13] T. Fazzini et al., Nucl. Instr. Meth. A 410 (1998) 213.
- [14] E. Gatti, F. De Martini, Nuclear Electronics 2, IAEA, Vienna, 1962, p. 265.
- [15] K.J.R. Rosman, P.D.P. Taylor, Pure and Appl. Chem. 70 (1998) 217.
- [16] G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A 729 (2003) 337.
- [17] V. Artemiev et al., Phys. Lett. B 345 (1995) 564.
- [18] A. De Silva et al., Phys. Rev. C 56 (1997) 2451.
- [19] A.S. Barabash et al., JETP Letters 79 (2004) 12.
- [20] A.A. Klimenko et al., Part. Nucl. Lett. 5 (2001) 80.
- [21] F.A. Danevich et al., Phys. Rev. C 62 (2000) 045501; 68 (2003) 035501.
- [22] Yu.G. Zdesenko et al., Nucl. Instr. Meth. A, in press.

- [23] I. Ogawa et al., Proc. 5-th Int. Workshop on Neutrino Oscillation and their Origin, NOON 2004, 10-15 Feb. 2004, Tokyo, Japan, to be published in Proc.
- [24] see discussion about discovery potential and sensitivity of 2β decay experiments in: Yu.G. Zdesenko et al., J. Phys. G.: Nucl. Part. Phys. 30 (2004) 971.
- [25] B. Buck, A.C. Merchant, S.M. Perez, J. Phys. G 17 (1991) 1223.
- [26] D.N. Poenaru, M. Ivascu, J. Physique 44 (1983) 791.
- [27] C. Xu, Z. Ren, Phys. Rev. C 69 (2004) 024614.
- [28] A.A. Sonzogni, Nucl. Data Sheets 93 (2001) 599.
- [29] W. Porschen, W. Riezler, Z. Naturforsch. 11a (1956) 143.
- [30] A. Isola, M. Nurmia, Z. Naturforsch. 20a (1965) 541.
- [31] R.D. Macfarlane, T.P. Kohman, Phys. Rev. 121 (1961) 1758.
- [32] B. Al-Bataina, J. Janecke, Radiochim. Acta 42 (1987) 159.
- [33] G. Kauw, Forsch. Laudes Nord.-Westfalen N 1640 (1966).
- [34] C. Cozzini et al., nucl-ex/0408006.
- [35] P. de Marcillac et al., Nature 422 (2003) 876.
- [36] M. Bravin et al., Astropart. Phys. 12 (1999) 107.
- [37] S. Cebrian et al., Phys. Lett. B 563 (2003) 48.