## arXiv:nucl-ex/0411006v1 4 Nov 2004

## Parallel momentum distribution of the <sup>28</sup>Si fragments from <sup>29</sup>P\*

WEI Yi-Bin,<sup>1,2</sup> MA Yu-Gang,<sup>1,†</sup> CAI Xiang-Zhou,<sup>1,‡</sup> ZHONG Chen,<sup>1</sup> CHEN Jin-Gen,<sup>1</sup> ZHANG Hu-Yong,<sup>1</sup>

FANG De-Qing,<sup>1</sup> WANG Kun,<sup>1</sup> MA Guo-Liang,<sup>1</sup> GUO Wei,<sup>1</sup> TIAN Wen-Dong,<sup>1</sup> SHEN Wen-Qing,<sup>1</sup> ZHAN Wen-Long,<sup>3</sup> XIAO Guo-Qing,<sup>3</sup> XU Hu-Shan,<sup>3</sup> SUN Zhi-Yu,<sup>3</sup> LI Jia-Xing,<sup>3</sup> GUO Zhong-Yan,<sup>3</sup> WANG Meng,<sup>3</sup> CHEN Zhi-Qiang,<sup>3</sup> HU Zheng-Guo,<sup>3</sup> CHEN Li-Xin,<sup>3</sup> LI Chen,<sup>3</sup> MAO Rui-Shi,<sup>3</sup> and BAI Jie<sup>3</sup>

<sup>1</sup>Shanghai Institute of Applied Physics, Chinese Academy of Sciences, P.O. Box 800-204, Shanghai 201800 <sup>2</sup>Graduate School of the Chinese Academy of Sciences

<sup>3</sup>Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000

Distribution of the parallel momentum of <sup>28</sup>Si fragments from the breakup of 30.7 MeV/nucleon  $^{29}$ P has been measured on C targets. The distribution has the FWHM with the value of 110.5  $\pm$ 23.5 MeV/c which is consistent quantitatively with Galuber model calculation assuming by a valence proton in <sup>29</sup>P. The density distribution is also predicted by Skyrme-Hartree-Fock calculation. Results show that there might exist the proton-skin structure in <sup>29</sup>P.

PACS numbers: 25.60.Gc, 23.20.En, 27.30.+t

Studies have been performed for many years for exotic nuclei with the increasing availability of radioactive nuclear beam around the world. The exotic structures of nuclei refer to the weakly bound systems that display the very diffuse surface of nearly pure nucleonic matter at densities far below that of normal nuclear matter. The nucleus <sup>11</sup>Li is the first observed case as a neutron halo nucleus [1], and other nuclei that are considered to have neutron halo are  ${}^{6}He$  [2],  ${}^{14}Be$  [3, 4] and etc. Such kind of halo nucleus provide good venue to investigate the weakly bound quantum system, which was not easily accessible before. Nuclear interactions in a low-density nuclear matter and loosely bound three-body interactions are among the current themes of interest related to these nuclei. In addition to the neutron halo, several experiments suggest the existence of proton halo nuclei. However, it is not yet fully established. More attentions are needed on the proton halo structure. Recently, the nuclei including  ${}^{8}B$  [1],  ${}^{12}N$  [5] and  ${}^{23}Al$  [6], which have the very low one-proton separated energy, are considered to have single proton halo structure. Compared with the neutron halo, proton halo is difficult to be formed because of the Coulomb repulsion interaction, but it is still an experimental observable phenomenon.

The methods which are used to study the halo structure have been explored and extended in recent years. For the size of the nuclei, the methods of the total reaction cross section and the one nucleon knocked-out

<sup>†</sup>Email: ygma@sinr.ac.cn

<sup>‡</sup>Email: caixz@sinr.ac.cn

cross section are widely used [7, 8, 9, 10]. From which, the sensitive probes to the abnormal structure of the nuclei, namely the radius and the density distribution can be deduced. However, to some nuclei, which have the larger cross section compared with the neighbored isotopes, may be not the candidates of the halo ones. This attributes to the extended density distribution of the core of the nucleus, which would induce the larger cross section [11, 12]. Therefore, it is necessary to find a suitable approach to explore the internal structure of the nuclei. The momentum distribution of the nucleus fragment could accord with such requirement. In a highly simplified picture, the longitudinal momentum distribution of fragment from the breakup of a loosely bound projectile directly reflects the internal momentum distribution of the valence nucleon and hence the square of the Fourier transform of its wave function. In the previous works, it was shown that the width of the momentum distribution can be understood in terms of the Fermi motion or a temperature corresponding to the nuclear binding energy [13, 14], and the momentum distribution of the valence-nucleon in a beam nucleus can be deduced from the observed momentum distribution of the fragment by taken the evaporation correction [15, 16]. Thus, halo formation in such loosely bound nucleus can be investigated by measuring the momentum distribution of the fragment from a breakup reaction. The wide spatial dispersion of a halo nucleon translates into a narrow momentum distribution. Many practical works have been done in the measurements of the momentum distribution widely both theoretically and experimentally [1, 2, 17]. It has been reliable while searching for the exotic structure of the nuclei. Very recently, the momentum correlation function among the nucleons of the neutron-rich nuclei has also been investigated and it seems to be sensitive to the binding energy per nucleon and single neutron separation energy [18, 19].

In the previous works, the study of the proton-halo

<sup>\*</sup>This work was supported partially by the Major State Basic Research Development Program under Contract No G200077400, the Chinese Academy of Sciences Grant for the Distinguished Young Scholars of National Natural Science Foundation of China (NNSFC) under Grant No. 19725521, NNSFC under Grant Nos 10135030 and 10328509 and the Shanghai Phosphor Program Under Contract Number 03 QA 14066.



FIG. 1: The experiment set-up for the measurement of parallel momentum distribution in RIBLL. For details see texts.

structure mainly focuses on the light proton-rich nuclei, whose mass numbers are not more than 25. It is interesting to investigate the characters of the nuclei with the mass number around 30. Navin et al. performed an experiment to study the structure of the  ${}^{26-28}$ P [20]. Some investigations of the density distribution of these P isotopes have also been done[21, 22]. Judging from the small single proton separation  $(S_p)$  energy (0.14 MeV, 0.9 MeV, 2.065 MeV, respectively) for <sup>26,27,28</sup>P [23], it is easy to detach the valence-proton from the core. This favors to form the proton halo structure, which has been approved through their narrow fragment momentum distributions. Considering the small  $S_p$  of <sup>29</sup>P (2.748 MeV), it should be interesting to measure the momentum distribution of the one-proton removal fragment of  $^{29}$ P. In this Letter, we report the experimental measurement and model analysis for the parallel momentum distribution of  $^{28}Si$  from the break-up of  $^{29}P$  on C target.

The experiment was performed at the Radioactive Ion Beam Line in Lanzhou (RIBLL). <sup>29</sup>P secondary beam was produced by the fragmentation of a 69 MeV/nucleon <sup>36</sup>Ar primary beam on Be target which was delivered by the Heavy Ion Research Facility in Lanzhou (HIRFL). A maximum intensity of 60 enA for <sup>36</sup>Ar was used to produce the secondary nuclear fragments. As shown in Fig. 1, the production target (Be) was mounted on the target box (T0). Before the target there is one Si detector which could give the identification of the nuclei. The angular acceptance was 6.5 msr, and the fragments emitted between  $-2.61^{\circ}$  and  $2.61^{\circ}$  could be transmitted through the spectrometer. The good collection is obtained in a cone around these angles because of the strongly focus of the projectile fragments. After T0, RIBLL can be used as a doubly achromatic magnetic spectrometer where nuclei produced in the reaction are separated and selected by means of magnetic rigidity  $(B\rho)$  and energy degrader ( $\Delta E$ ). The momentum acceptance  $\Delta p/p$  of the magnetic spectrometer was set at about 10%

The variable slits and energy degraders, which were made by Al foils and mounted on curved frames, were placed at the dispersive focal plane C1 and C2. The



FIG. 2: The momentum distribution data from one-proton removal in  $^{29}$ P on *Be* target at 30.7 MeV/nucleon. The dashed line is the Glauber calculation result. The error bars are only the statistical.

time of flight (TOF) of the fragments was measured by two scintillator detectors installed at the first (T1) and second (T2) achromatic focal planes with a flight distance of 16.8m. The measurement of momentum distribution was done using the TOF. The *Si* detector before *C* target is 300  $\mu$ m. A telescope was installed at T2, which consisted of three transmission *Si* surface barrier detectors followed by a CsI (T1) crystal and gave the energy losses ( $\Delta E$ 's) and total energy of the reaction products, where the identification of nuclei could be completed. The thickness of three *Si* detectors were 150, 150 and 1000  $\mu$ m, respectively, and the energy resolutions of the Si detectors were not greater than 0.8%. It is possible to identify the reaction products and get the momentum distribution combining TOF and  $\Delta E$  [24, 25, 26].

With the  $B\rho$ - $\Delta E$  selection of RIBLL, the secondary beam of  $^{29}P$  can be identified before C target. The PPACs will ensure the position of  $^{29}$ P to focus on the center of the target. After the target, the reaction products will be selected by the  $B\rho$ - $\Delta E$  again. After the flight during the TOF2, the significative events will be accepted by the final three Si detectors. With the TOF2, the fragments of the reaction will be achieved. Thus, the calibration of the TOF and Si detectors are very important. The good linear relationship between the channel number and the flight time (ns) have been obtained. From the flight time of the nuclei, we could get the longitudinal (parallel) momentum distribution. Also, in this experiment, the angular distribution can be deduced from the position signals obtained through three PPACs before and after the C target.

Fig. 2 shows the experimental result of the parallel momentum distribution and Fig. 3 displays the angular distribution of the <sup>28</sup>Si fragment in <sup>29</sup>P. The width (FWHM) of momentum distribution was found to be 110.5  $\pm$  23.5 MeV/c and that of scattering angular distribution is 9.2°



FIG. 3: The angular distribution data from one-proton removal in  $^{29}$ P on *Be* target at 30.7 MeV/nucleon. The solid line is the fit with single Gaussian distribution.

which is obtained from the one-Gaussian fit.

As we know, the lower nucleon separation energy favors to form the exotic structure. For the light proton halo nuclei which have been discovered, the one proton separation energy is smaller than that of the neighboring nuclei among its isotopes. The proton most far away from the core due to its small separation energy will appear in the more extended region which is defined halo. For the candidates of proton halo nuclei, <sup>8</sup>B, <sup>14</sup>N, <sup>17</sup>Ne and <sup>23</sup>Al all have the smaller one proton separation energies. For proton-rich nuclei, it is possible to form the proton halo structure just similar with the neutron halo existing in the neutron-rich nuclei. For P-isotope, the number of proton is 15.  $^{26-29}P$  have the more number of protons than that of neutrons. Considering the small one proton separation energy, one could imagine the possibility of the proton halo structure existing in the proton-rich nuclei of P-isotope. For  ${}^{26,27}P$  nuclei, the experiment results have achieved to indicate their halo structure [20, 25, 26, 27]. But for <sup>28</sup>P, it is not sure for the existence of the exotic structure [26]. Compared with the separation energy of <sup>28</sup>P, it is more difficult to form the halo structure for <sup>29</sup>P. But it is still interesting to investigate its structure for the normal or proton-skin character.

We attempt to obtain the theoretical results using the Glauber model, which can give the reasonable results of the fragment momentum distribution. The Glauber model is a microscopic reaction theory based on the eikonal approximation and on the bare nucleon-nucleon interaction. It is now a standard tool to calculate the reaction of a weakly bound nucleus. The observed interaction cross sections can be related to the wave functions of these nuclei through this model and also one can obtain information on the structure of these exotic isotopes. The accurate nuclear wave functions play the very important role in the reaction simulation. Varga et al developed the Monte Carlo integration in Glauber model and some good results have been obtained [28]. Abu-Ibrahim et al. developed this model and some reasonable results have been achieved compared with the experiment [29]. We took his code in our calculation. More elaborated treatment beyond the optical-limit approximation is worked out for the valence-nucleon part by using the Monte Carlo quadrature with the Metropolis algorithm. This is suitable for reactions involving those nuclei which have spatially extended and low density distribution as in a halo nucleus.

In the model, the density of the projectile is treated as two parts: the core and the valence nucleon density distributions. The core density is assumed to be given by a combination of Gaussians:

$$\rho(r) = \Sigma c_i e^{-a_i r^2} \tag{1}$$

Also, the density  $\rho$  is normalized to the mass number of a nucleus,  $\int d\mathbf{r} \rho(\mathbf{r}) = \mathbf{A}$ . The valence-nucleon density distribution will be generated through the model with the typical parameters including the separated energy, the angular momentum value and so on. The density of the target is treated similar with that of the core of the projectile. Thus, the valence-nucleon character will be included in the Glauber calculation and the results show that such consideration is reliable. More details about Glauber model could be found in Ref. [29].

Using such a model, we calculated the reaction of the 30.7MeV/nucleon <sup>29</sup>P on <sup>12</sup>C target. The parallel momentum distribution of the <sup>28</sup>Si fragment after oneproton removal from <sup>29</sup>P has been obtained. The calculated FWHM is 91 MeV/c and the result is the dashed line shown in Fig. 2. In our calculation, the valence proton density distribution of <sup>29</sup>P is decided by the separation energy, the angular momentum value and the node of the density distribution of the valence proton. The core of the projectile and the target density distributions are fitted with five gauss function according to the calculated results with Skyrme-Hartree-Fock(SHF) method. Thus, the parallel momentum distribution of <sup>29</sup>P fragmentation is self-consistent in experiment and theory.

In Ref. [30], the fragment angular distribution will give the almost same character with the momentum distribution. The angular distribution indicates the scattering character of the projectile. Compared with the neutronhalo nuclei [31], the angular distribution could also describe the structure in the proton-rich nuclei. Fig. 3 shows our experimental results. The peak of the angular distribution of <sup>28</sup>Si fragment shows the scattering angular during the reaction. From the value of the angular distribution, it is hard to say the existence of the exotic halo structure in <sup>29</sup>P. But the proton-skin structure might exist.

In order to understand the structure of  $^{29}$ P further, it is more clear to describe  $^{29}$ P from the sight of nucleon orbits. For  $^{29}P$ , the last proton orbit is 1s1/2, and the other 14 protons and 14 neutrons form the structure that similar with that of  $^{28}$ Si. The single-particle energies of



FIG. 4: The calculated density distribution of protons and neutrons for  ${}^{28}Si$  (a) and  ${}^{29}P$  (b) with SHF model. In (c), the valenceproton density distribution is from Glauber model while the density distribution of  ${}^{28}Si$  and  ${}^{29}P$  are from SHF model; the total one is the sum of the valence-proton density and the density of  ${}^{28}Si$ .

protons in level 1d5/2 and 2s1/2 for <sup>29</sup>P are, respectively, 11.585MeV and 2.748MeV. These indicate that <sup>29</sup>P can be approximately considered as a <sup>28</sup>Si core plus one proton because the valence-proton is weakly bound. One could use some models to study such nucleus through the density distribution. The SHF is one of the suitable models which can give the reasonable nucleus structure

[21, 22]. We calculated the density distributions of  $^{29}$ P and  $^{28}$ Si shown in Fig. 4. The dashed lines in (a) and (b) are the proton density distributions. From the figure one find that the density distribution of neutrons in  $^{28}$ Si and  $^{29}$ P are very similar, but the density distribution of protons in  $^{29}$ P extends farther than that of  $^{28}$ Si. It displays that the radii of proton and matter of  $^{29}$ P are a little larger than those of  $^{28}$ Si. This displays the weakly unbound status of the last proton of  $^{29}$ P, namely the proton-skin behavior.

In the Glauber model the valence-nucleon's wave function can be obtained with the three input parameters which has been mentioned above. It is easy to get the density distribution from the wave function after normalization. The solid line shown in Fig.4(c) is the valenceproton of <sup>29</sup>P density distribution from the Glauber model. With the density distribution of the core, <sup>28</sup>Si, we get <sup>29</sup>P density distribution, which is the total one shown in Fig.4(c). Compared with the result from SHF model, the density distribution of <sup>29</sup>P from Glauber model is suitable to embody the nuclear character and it would be reliable to get the momentum distribution.

In summary, the experimental measurement of the parallel momentum distribution of <sup>28</sup>Si fragment from the breakup of 30.7 MeV/nucleon <sup>29</sup>P has been reported for the first time. The parallel momentum width is 110.5± 23.5 MeV/c FWHM which is consistent with the calculation of the Glauber model assuming the core plus one proton. The Skyrme-Hartree-Fock calculation also gives a proton-skin density distribution in comparison with <sup>28</sup>Si. Taken together with the theoretical analysis of the density distribution and the nucleon orbit arrangement, it is obviously hard to form the halo structure in <sup>29</sup>P. But the possibility of the proton-skin structure of <sup>29</sup>P exists, and this nucleus could be understood as the transition one between the exotic and stable isotopes.

We appreciate helpful discussions with Prof. Gen-Ming Jin of Institute of Modern Physics.

- [1] Tanihata I et al. 1988 Phys. Rev. Lett. 55 2676.
- [2] Kobayashi T et al., Phys. Rev. Lett. 60 (1988) 2599.
- [3] Tanihata I 1988 Nucl. Phys. A 488 113.
- [4] Suzuki T et al. 1999 Nucl. Phys. A 658 313.
- [5] Warner R et al. 1998 Nucl. Phys. A 635 292.
- [6] Cai X Z et al. 2002 Phys. Rev. C 65 024610.
- [7] Shen W Q et al. 1989 Nucl. Phys. A 491 130.
- [8] Ma Y G et al. 1993 Chin. Phys. Lett. 10 464.
- [9] Ma Y G et al. 1993 Phys. Lett. B **302** 386.
- [10] Ma Y G et al. 1993 Phys. Rev. C 48 850.
- [11] Saint-Laurent M G et al. 1989 Z. Phys. A 332 457.
- [12] Bazin D et al. 1998 Phys. Rev. C 57 2156.
- [13] Goldhaber A 1974 Phys. Lett. B 53 306.
- [14] Fujita T and H $\ddot{u}$ fner H 1980 Nucl. Phys. A **343** 493.
- [15] Hanelt R et al. 1993 Z. Phys. A **346** 43.
- [16] Ma Y G et al. 2002 Phys. Rev. C 65 051602(R).

- [17] Guimarães V et al. 2000 Phys. Rev. C $\mathbf{61}$ 064609
- [18] Wei Y B, Ma Y G et al. 2004 Chin. Phys. Lett. 21 629;
- [19] Wei Y B, Ma Y G et al. 2004 Phys. Lett. B 586 225.
- [20] Navin A et al. 1998 Phys. Rev. Lett. **81** 5089.
- [21] Ren Z Z et al. 1996 Phys. Rev. C 53 R572.
- [22] Ren Z Z et al. 1999 Nucl. Phys. A 652 250.
- [23] Wapstra A H and Audi G 1985 Nucl. Phys. A 432 1.
- [24] Cai X Z et al. 2001 Chin. Phys. Lett. 18 1189.
- [25] Fang D Q et al. 2001 Chin. Phys. Lett. 18 1033.
- [26] Fang D Q et al. 2001 Eur. Phys. J. A **12** 335.
- [27] Liu Z H et al. 2004 Phys. Rev. C 69 034326.
- [28] Varga K et al. 2002 Phys. Rev. C 66 034611.
- [29] Abu-Ibrahim B et al. 2003 Comp. Phys. Comm 151 369.
- [30] Zahar M et al. 1993 Phys. Rev. C 48 R1484.
- [31] Marques F M et al. 1996 Phys. Lett. B 381 407.