

## High $p_T$ $\gamma$ -hadron and $\pi^0$ -hadron correlations in $\sqrt{s_{NN}} = 200$ GeV Au + Au collisions in PHENIX

Jiamin Jin<sup>1</sup> (for the PHENIX Collaboration)

<sup>1</sup> Columbia University, New York, NY 10025, USA

*Received 1 January 2005*

**Abstract.** This poster presents the PHENIX results on high- $p_T$   $\gamma$ -hadron and  $\pi^0$ -hadron correlations and shows the modification of away side jet shape by the medium. By comparing their jet shapes and yields, we see some evidence of the direct  $\gamma$  contribution to correlation functions.

*Keywords:* Correlation Function, Jets, Direct Photon

*PACS:* 25.75.-q

### 1. Introduction

PHENIX has measured direct photon production in Au + Au at  $\sqrt{s_{NN}} = 200$  GeV [1]. Due to the suppression of high- $p_T$  pion production, a large photon excess over meson decay background is seen in central Au + Au collisions at  $p_T > 6$  GeV/c. The much higher statistics 2004 RHIC Au + Au run has been used to further explore the possibility of constructing direct photon-hadron correlations. Since direct photons, once produced, interact with the medium much more weakly than hadrons, they provide a better measurement of the energy and direction of the away side jets, thus we gain direct information on the initial state effect and better control on the modification of the away side jets by the strongly interacting medium. The high  $p_T$  photon sample PHENIX measures is a mixture of direct photons and fragmentation (mostly  $\pi^0$  decay) photons. Therefore, it is important to study the high- $p_T$   $\gamma$ -hadron and  $\pi^0$ -hadron correlations and compare them systematically.

### 2. High $p_T$ $\gamma$ -hadron Correlations and $\gamma$ Per Trigger Yield


The  $\gamma$ -hadron correlation analysis presented in QM2005 poster represents 1 billion minimum bias events. We use photons with  $p_T > 5$  GeV/c as leading particles and charged hadrons with  $1\text{GeV}/c < p_T < 5$  GeV/c as associated particles. The

correlation function  $C(\Delta\phi)$  is constructed by pairing leading and associated particles.  $C(\Delta\phi) = \frac{N_{real}(\Delta\phi)}{N_{mix}(\Delta\phi)}$ , where  $\Delta\phi$  is the difference of the azimuthal angles of the pair. The  $N_{real}(\Delta\phi)$  is built from pair members belonging to the same event and  $N_{mix}(\Delta\phi)$  is built from pair members belonging to different events. This event mixing technique removes the detector acceptance effects. The correlation function can be decomposed into jets and flow components [ 2]:

$$C(\Delta\phi) = J(\Delta\phi) + \xi(1 + 2v_2^l v_2^a \cos(2\Delta\phi)) \quad (1)$$

where  $J(\Delta\phi)$  is the jet contribution,  $v_2^l(v_2^a)$  is the elliptic flow for the leading(associated) particles and  $\xi$  is the background level. The  $v_2$  values are measured using reaction plane method, whereas  $\xi$  is fixed using ZYAM(Zero jet Yield At Minimum) assumption [ 2, 3]. After removing the  $v_2$  modulated background, we correct the remaining  $J(\Delta\phi)$  by single particle efficiency and PHENIX  $\Delta\eta$  acceptance, normalize it by the number of triggers, thus obtain the photon per trigger yield.

Figure 1 shows the typical  $\gamma$ -hadron correlation function and per trigger yield for central Au + Au collisions. We see a strongly modified(non-gaussian) away side shape.




**Fig. 1.** a)  $\gamma$ -hadron correlation function, lines indicated the level of flow modulated background and its systematic error band. b) Corresponding photon per trigger yield.

### 3. Results


A simultaneous fit is applied to the correlation functions to determine the jet and the flow contributions. The jet widths as well as the near side jet yield are extracted directly from the fit, whereas the away side yield is obtained by integrating the correlation function over  $\Delta\phi \in [\pi/2, 3\pi/2]$ , then subtracting the background in the same range. This region is chosen so that the overall flow contribution cancels out, thus making the yield not sensitive to  $v_2$ .

PHENIX has measured  $\pi^0$ -hadron correlations in a similar way with trigger  $\pi^0$   $p_T > 5$  GeV/c [ 4]. Figure 2 shows the comparison of near side jet width and yield between  $\gamma$ -hadron and  $\pi^0$ -hadron correlations. They have comparable widths,

whereas  $\gamma$  jet yields are consistently lower than  $\pi^0$  jet yields. Figure 3 plots the ratio of  $\gamma$  near side jet yield over  $\pi^0$  near side jet yield as a function of associated particle  $p_T$  for different centrality bins. The ratios are all below one. Direct photons contribute to this yield reduction since they are not from jets, thus having zero yield on the near side. The ratios also become smaller for more central bins. This trend is consistent with the results in [ 1 ], which established that there is larger direct photon signal in central collisions.




**Fig. 2.** a) comparison of near side width between  $\gamma$  jet (left) and  $\pi^0$  jet (right), b) same for near side yield



**Fig. 3.** near side yield ratio between  $\gamma$  jet and  $\pi^0$  jet

Figure 4 shows  $\gamma$ -hadron and  $\pi^0$ -hadron correlation functions on top of each other for 3 centrality bins and 2 associated particle  $p_T$  bins. It allows us to do a

direct comparison and have some feeling about the subtle jet shape change between the two. In addition to the near side jet yield reduction noted above, the upper panels show that the away side peaks in  $\gamma$ -hadron correlations seem to have a stronger distortion at low  $p_T$ , whereas the lower panels show that once the  $p_T$  of the associated hadrons reach 3 GeV/c, the two correlations appear similar on the away side.



**Fig. 4.** Upper row:  $\gamma$ -hadron (filled points) and  $\pi^0$ -hadron (open circles) correlations for 3 different centrality bins and 1–2 GeV/c associated  $p_T$  bin. Lower row: same for 3–5 GeV/c associated  $p_T$  bin.

## 4. Conclusions

High  $p_T$   $\gamma$ -hadron correlation functions are measured in different centrality bins and associated particle  $p_T$  bins. A distortion on the away side jet due to the coupling of the hadrons and the strong interacting medium is observed. Comparison between  $\gamma$ -hadron and  $\pi^0$ -hadron correlations shows a decrease of the near side  $\gamma$  jet yield and a stronger modification on the away side  $\gamma$  jet, both effects can be attributed to the presence of direct photons. Further investigation on the role direct photon plays in correlation studies is undergoing.

## References

1. S.S. Adler et al., *Phys. Rev. Lett.* **94** 232301 (2005).
2. J. Jia, *arXiv:nucl-ex/0510019*.
3. S.S. Adler et al. [PHENIX Collaboration], *arXiv:nucl-ex/0507004*
4. P. Hua [PHENIX Collaboration], quark matter proceedings.