

## MEASUREMENTS OF $C_Z$ , $C_X$ FOR $K^+\Lambda$ AND $K^+\Sigma^0$ PHOTOPRODUCTION

ROBERT BRADFORD

*Department of Physics and Astronomy  
University of Rochester  
500 Wilson Boulevard  
Rochester, NY 14627-0171, USA  
E-mail: bradford@pas.rochester.edu*

REINHARD SCHUMACHER

*Dept of Physics  
Carnegie Mellon University  
5000 Forbes Avenue  
Pittsburgh, PA 15213  
E-mail: schumacher@cmu.edu*

FOR THE CLAS COLLABORATION.

The CLAS collaboration has recently completed first measurements of the double polarization observables  $C_x$  and  $C_z$  for the reactions  $\gamma p \rightarrow K^+\Lambda$  and  $\gamma p \rightarrow K^+\Sigma^0$ .  $C_x$  and  $C_z$  are the beam-recoil polarization asymmetries measuring the polarization transfer from incoming circularly polarized photons to outgoing hyperons along two directions in the reaction's production plane. The  $\Lambda$  is found to nearly maximally polarized along the direction of incident photon's polarization for forward-going kaons. Polarization transfer to the  $\Sigma^0$  is different from the  $\Lambda$  case.

### 1. Introduction

Measurement of polarization observables have long been recognized as key to unraveling baryon production mechanisms. This work represents the first measurement of  $C_x$  and  $C_z$  for  $\gamma p \rightarrow K^+\Lambda$  and  $\gamma p \rightarrow K^+\Sigma^0$ .

### 2. $C_x$ and $C_z$

$C_x$  and  $C_z$  measure the polarization transfer from a circularly polarized incident photon beam to the recoiling  $\Lambda$  or  $\Sigma^0$  baryons along two orthogonal directions in the production plane of the  $K^+$ -hyperon system. The  $\hat{x}$  and  $\hat{z}$

directions are defined in the CM frame, with  $\hat{z}$  lying along the directions of the incident photon beam's polarization. In this analysis, the polarization transfer was measured through the beam helicity asymmetry according to

$$A_{x/z}(\cos\theta_p) = \frac{N^+(\cos\theta_p) - N^-(\cos\theta_p)}{N^+(\cos\theta_p) + N^-(\cos\theta_p)} = \alpha_{eff}\eta C_{x/z} \cos\theta_p \quad (1)$$

where  $\cos\theta_p$  is the direction of the proton from the hyperon's decay measured in the hyperon rest frame with respect to the  $\hat{x}$  or  $\hat{z}$  axis.  $\eta$  is the polarization of the incident photon beam,  $N^+(\cos\theta_p)$  and  $N^-(\cos\theta_p)$  are the beam helicity dependent hyperon yields in a given  $\cos\theta_p$  bin.  $\alpha_{eff}$  is the effective weak decay asymmetry parameter, and has a value of 0.642 for  $K^+\Lambda$  and -0.165 for  $K^+\Sigma^0$ . The value of  $\alpha_{eff}$  for the  $\Sigma^0$  decay arises from our technique of measuring the proton distribution in the rest frame of the  $\Sigma^0$ , not the  $\Lambda$ ; this dilutes its value to less than the nominal -0.642/3.


### 3. Experimental Setup and Analysis

The data were taken using the CLAS spectrometer in Hall B at Jefferson Lab. The experiment used a circularly polarized photon beam incident on a liquid hydrogen target. Data were taken with endpoint photon energies of 2.4 and 2.9 GeV. From this dataset, we also measured differential cross sections, which are currently available in preprint<sup>1</sup>.


All analyzed events were required to have explicit detection of the  $K^+$  and proton. The  $\Lambda$  or  $\Sigma^0$  hyperons were identified in the  $p(\gamma, K^+)Y$  missing mass. The data were binned in beam helicity, the cosine of the kaon angle in the CM frame ( $\cos(\theta_{KCM})$ ), the cosine of the proton angle ( $\cos(\theta_p)$ ) and photon energy ( $E_\gamma$ ). Within kinematic each bin, yields were extracted by fitting a Gaussian peak to each hyperon in the missing mass spectrum. Backgrounds were modeled with a polynomial. The beam helicity asymmetry was plotted against  $\cos(\theta_p)$  and the slope of this distribution was extracted with a linear fit. Complete analysis details are available elsewhere<sup>2</sup>.

### 4. Results and Discussion

Sample results are presented in Figures 1 and 2. The data are plotted with predictions from the Kaon-MAID<sup>4</sup> (solid line) and Janssen<sup>3</sup> (dashed line) isobar models. The data plotted are for only a few representative bins in  $\cos(\theta_{KCM})$ . The full results include nine bins in  $\cos(\theta_{KCM})$  for  $K^+\Lambda$  and six bins for  $K^+\Sigma^0$ .



(a)



(b)

Figure 1.  $C_x$  and  $C_z$  for  $K^+\Lambda$  in two different kaon-angle bins. Top:  $\cos(\theta_K) = -0.75$ , bottom:  $\cos(\theta_K) = 0.25$ . The data are a subset of the 2005 CLAS results. The curves are predictions from the Kaon-MAID (solid line,<sup>4</sup>) and Janssen (dashed line,<sup>3</sup>) isobar models.




Figure 2.  $C_x$  and  $C_z$  for  $K^+\Sigma^0$  for  $\cos(\theta_K) = 0.5$ . The data are a subset the 2005 CLAS results. The curves are predictions from the Kaon-MAID (solid line, <sup>4</sup>) and Janssen (dashed line, <sup>3</sup>) isobar models.

The  $\Lambda$  results show some  $W$ -dependent structure at backward kaon angles and then stabilize at more forward-going kaon angles. For this hyperon,  $C_x$  is near one over most of the forward hemisphere of the kaon angle while  $C_z$  is near zero for the same range.

The  $K^+\Sigma^0$  results show no preferred direction for the polarization over the kaon angle range. The precision of these results here appears worse due to the small value of  $\alpha_{eff}$ .

Of the models shown, the Janssen does a good job of following the data. The MAID curve does not fair as well. This model has the oddity of predicting that  $C_z$  saturates at -1 in  $K^+\Lambda$  for forward-going kaons.

## References

1. R. Bradford, *et al.* Preprint available at: nucl-ex/0509033. Submitted to *Phys. Rev. C*.
2. R. Bradford, *Measurement of differential cross sections and  $C_x$  and  $C_z$  for  $\gamma p \rightarrow K^+\Lambda$  and  $\gamma p \rightarrow K^+\Sigma^0$  using CLAS at Jefferson Lab.* Ph.D. thesis, Carnegie Mellon University, 2005, to be published. Available on-line at: <http://www.jlab.org/Hall-B/general/thesis/bradford/index.html>
3. S. Janssen, *Strangeness production on the nucleon.* Ph.D. thesis, University of Gent, 2002.
4. F.X. Lee, *et al.*, *Nucl. Phys. A* **695**, 237 (2001).