$\Lambda(1520)$ production in d+Au collisions at RHIC

Christina Markert for the STAR collaboration

[†] Physics Department, Kent State University, Kent, OH 44242, USA

Abstract. Recent results of $\Lambda(1520)$ resonance production in d+Au collisions at $\sqrt{s_{\rm NN}} = 200$ GeV are presented and discussed in terms of the evolution and freezeout conditions of a hot and dense fireball medium. Yields and spectra are compared to results from p+p and Au+Au collisions. The $\Lambda(1520)/\Lambda$ ratio in d+Au collisions ratio is consistent with the ratio in p+p collisions. This suggests a short time for elastic interactions between chemical and thermal freeze-out. One can conclude that the interaction volume in d+Au collisions is small.

1. Introduction

In heavy ion collisions an extended hot and dense medium is created. The properties (mass, width, momentum distribution, yield) of the produced resonances depend on the fireball conditions of temperature, pressure and lifetime. During the fireball expansion the short lived resonances and their hadronic decay daughters may interact with the medium. Two freeze-out surfaces can be defined, chemical and thermal, representing the conditions when inelastic and elastic interactions cease respectively. In a dynamically evolving system produced resonances decay and may by regenerated. Hadronic decay daughters of resonances which decay inside the medium may also scatter with other particles from the medium. This results in a signal loss, because the reconstructed invariant mass of the decay daughters no longer matches that of the parent. The strength of the rescattering and regeneration via the inverse decay processes for resonances and their decay particles which are dominant after chemical but before the kinetic freezeout depend on individual cross sections. These interactions can lead to changes of the reconstructed resonance yields, momentum spectra, widths and masses [1]. Rescattering will decrease the measured resonance yields while regeneration will increase them. For small system sizes such as the one produced in p+p and d+Au collisions we would expect a short or even no interaction time of hadrons since the interaction volume is very small. In order to try to understand the medium effect in a d+Au system during the evolution and expansion of the fireball, we compare resonance yields and spectra to elementary p+p collisions, and system size and time extended Au+Au collisions.

2. Resonance Reconstruction

The $\Lambda(1520)$ resonance is determined by the invariant mass reconstruction using the p and K⁻ decay daughters. The decay candidates are identified by their energy loss (dE/dx) in the STAR *Time Projection Camber* (TPC). The resonance signal is obtained by the invariant mass reconstruction of each daughter combination and the subsequent subtraction of the combinatorial background calculated via a mixed event technique. Fig 1 shows the invariant mass spectrum before (insert) and after background subtraction. The trigger selection is set to a minimum bias trigger.



Figure 1. Invariant mass distributions in d+Au collisions at $\sqrt{s_{\text{NN}}} = 200 \text{ GeV}$ before (inset) and after mixed-event background subtraction for $\Lambda^* \to p + K^-$.

3. Resonance Spectra and Yields

The $\Lambda(1520)$ transverse mass spectra at mid-rapidity are shown in Fig 2 (left) for minimum bias p+p and d+Au collisions. The yield is obtained by the extrapolation and integration of the momentum distribution. For Au+Au collisions an inverse slope parameter has to be assumed (T = 350 MeV) due to the small signal. Yield and $\langle p_T \rangle$ comparisons are shown in Table 1. In order to compare different collision systems we normalize the yield to the yield of the corresponding measured ground state particle. Fig.2 (right) shows the $\Lambda(1520)/\Lambda$ ratios versus number of participants for different collision systems. This ratio remains constant in p+p and d+Au collisions and decreases in Au+Au collisions with increasing centrality [2, 3, 4]. The comparison to microscopic (UrQMD [1]) and thermal ([5, 6]) model predictions for Au+Au collisions indicates a hadronic interaction phase were re-scattering and regeneration may occur, which can not be described by the thermal model. Elastic interactions after chemical freeze-out happen only in an extended reaction volume. In d+Au collisions we would not expect an

	T [MeV]	$\langle p_T \rangle ~[MeV/c]$	$(dn/dy) _{y=0}$
$(\bar{\Lambda}^* + \Lambda^*)/2_{\rm p+p}$	338 ± 42	1.08 ± 0.09	0.0034 ± 0.0006
$(\bar{\Lambda}^* + \Lambda^*)/2_{d+Au}$	382 ± 75	1.17 ± 0.15	0.0149 ± 0.0014
$\Lambda^*_{\rm Au+Au}$	fixed $T=350$		0.58 ± 0.28

Table 1. Temperature , yield, and $\langle p_T \rangle$ from an exponential fit to the p_T spectra.



Figure 2. Left: The transverse mass spectra for $\Lambda(1520)$ in p+p, d+Au collisions at $\sqrt{s_{\rm NN}} = 200$ GeV. Right: $\Lambda(1520)/\Lambda$ ratio for p+p, d+Au and Au+Au collisions at $\sqrt{s_{\rm NN}} = 200$ GeV at mid-rapidity. Statistical and systematic errors are included. Thermal model and microscopic model predictions (UrQMD) for $\Lambda(1520)/\Lambda$ in central Au+Au systems are shown. Statistical and systematic errors are included.

extended reaction volume and therefore the $\Lambda(1520)/\Lambda$ ratio is expected to be similar to p+p collisions, which is in agreement with the data. Based on the data a lower limit of $\Delta \tau > 4$ fm/c can be placed on the interaction time in central Au+Au collisions [7, 8, 9]. The interaction time here is defined as the time between chemical and kinetic freeze-out. As the $\Lambda(1520)/\Lambda$ in d+Au is in agreement with the thermal model and the value for p+p, where one expects a very short time ($\Delta \tau \approx 0$ fm/c) for a hadronic phase, we can also conclude a $\Delta \tau \approx 0$ fm/c for the elastic interaction hadronic phase in a d+Au collision.

4. Conclusion

The $\Lambda(1520)/\Lambda$ ratio in d+Au collisions at $\sqrt{s_{\rm NN}} = 200$ GeV is in agreement with the ratio in p+p collisions. This indicates a zero life time between the chemical and thermal freeze-out where elastic interactions occur. The slopes of the momentum spectrum of the $\Lambda(1520)$ measured in p+p and d+Au is equal, which also leads to the conclusion of a no hadronic phase after chemical freeze-out.

REFERENCES

- [1] M. Bleicher and J. Aichelin, Phys. Lett. B530 (2002) 81, hep-ph/0201123.
- [2] C. Markert at al., J.Phys. **G30** (2004) 1313.
- [3] C. Markert et al., Proc. 19th Winter Workshop on Nuclear Dynamics (2003) 71.
- [4] L. Gaudichet J.Phys. **G30** (2004) 549.
- [5] P. Braun-Munzinger, D.Magestro, K. Redlich and J.Stachel, Phys. Lett. B518 (2001) 41, private communication.
- [6] W. Florkowski, W. Broniowski and P. Bozek, J.Phys. G30 (2004) 1321.
- [7] G. Torrieri and J. Rafelski, Phys. Lett. **B509** (2001) 239-245.
- [8] C. Markert, G. Torrieri, J. Rafelski, Campos do Jordao 2002, New states of matter in hadronic
- [9] G. Torrieri and J. Rafelski, J.Phys. G28 (2002) 1911-1920. interactions 533-552, hep-ph/0206260.