Target Structure Independent ⁷Li Elastic Scattering at Low Momentum Transfers

O.A. Momotyuk^{a,1}, N. Keeley^{b,*}, K.W. Kemper^a,

B.T. Roeder^a, A.M. Crisp^a, W. Cluff^a, B.G. Schmidt^a,

M. Wiedeking^a, F. Maréchal^c, K. Rusek^d, S.Yu. Mezhevych^d,

J. Liendo^e

^aDepartment of Physics, Florida State University, Tallahassee, Florida 32306-4350, USA

^bCEA-Saclay, DSM/DAPNIA/SPhN, F-91191 Gif-sur-Yvette, France

^cInstitut de Recherches Subatomiques, CNRS/IN₂P₃-ULP, F-67037 Strasbourg Cedex 2, France

^dDepartment of Nuclear Reactions, The Andrzej Soltan Institute for Nuclear Studies, Hoża 69, PL-00-681 Warsaw, Poland

^ePhysics Department, Simon Bolivar University, Caracas, Venezuela

Abstract

Analyzing powers and cross sections for the elastic scattering of polarized ⁷Li by targets of ⁶Li, ⁷Li and ¹²C are shown to depend only on the properties of the projectile for momentum transfers of less than 1.0 fm⁻¹. The result of a detailed analysis of the experimental data within the framework of the coupled channels model with ground state reorientation and transitions to the excited states of the projectile and targets included in the coupling schemes are presented. This work suggests that nuclear properties of weakly-bound nuclei can be tested by elastic scattering experiments, independent of the target used, if data are acquired for momentum transfers less than ~ 1.0 fm⁻¹.

Key words: ${}^{12}C({}^{7}Li, {}^{7}Li), {}^{7}Li({}^{7}Li, {}^{7}Li), {}^{6}Li({}^{7}Li, {}^{7}Li)$, coupled channels calculations PACS: 25.70.Bc, 24.70.+s, 24.10.Eq

It has been known for some time [1,2,3,4,5,6] that the structure of a heavyion projectile determines the characteristics of its elastic scattering. For ¹⁰B,

* Corresponding author

Email address: keeley@nucmar.physics.fsu.edu (N. Keeley).

¹ Permanent address: Institute for Nuclear Research, Kyiv, Ukraine.

a nucleus with a large ground state quadrupole moment, the addition of a quadrupole reorientation term to the optical potential decreases the depth of the minima in the elastic scattering cross section angular distribution at large scattering angles when compared with ¹²C scattering from the same target [2]. The clearest signature of the influence of the projectile's internal structure on the elastic scattering was provided by comparing ⁶Li and ⁷Li vector and tensor analyzing powers (APs) for energies close to the Coulomb barrier [5,6]. The difference in sign of the vector analyzing powers between ⁶Li and ⁷Li scattering was shown to arise from strong coupling between the ground, bound and resonant excited states of these loosely bound nuclei. This difference in analyzing powers was also shown to be present for nuclear dominated scattering [7,8]. The strong influence of the projectile structure as observed through the analyzing powers is made possible by the weak spin-orbit force in the projectile-target interaction [9].

The surprising result to date is that no clear signature for the structure of the target has been observed in Li scattering APs. This could be a result of the limited number of data sets, or the fact that none of the targets are loosely bound. For this reason, we have undertaken a study of ⁷Li scattering from the loosely bound targets ⁶Li and ⁷Li. The ⁶Li nucleus is bound by 1.47 MeV and it possesses a triplet of states $(3^+, 2^+, 1^+)$ that are strongly excited in nuclear scattering. While ⁷Li is more tightly bound (2.45 MeV), it has a large ground state quadrupole moment that results in a strong reorientation during scattering as well as having strongly coupled excited states. These two targets have different properties that significantly affect their elastic scattering and might yield knowledge about the importance of target choice when extracting the properties of loosely bound nuclei like ⁶He and ¹¹Li from elastic scattering.

In this work, we report new cross section and analyzing power angular distributions for the scattering of polarized ⁷Li beams from targets of ⁶Li and ⁷Li. A polarized ⁷Li beam was produced by the Florida State University (FSU) optically pumped lithium ion source [10] and accelerated to 42 MeV by the FSU Tandem/Linac accelerator. This energy was chosen because of well known analyzing powers that allow the beam polarization to be determined [7,11]. The beam polarizations were: $t_{10} = 0.76 \pm 0.07$ and $t_{20} = 0.37 \pm 0.07$. The targets were enriched in ⁶Li to 95% and transferred under vacuum to the main scattering chamber. Absolute cross section normalisations were obtained using the target thicknesses determined by comparison of proton elastic scattering measurements with previously accurately determined absolute cross sections [12]. The 5% content of ⁷Li in the targets was enough to allow both sets of measurements to be carried out simultaneously, thus allowing a direct comparison between these two data sets. The left panel in Fig. 1 shows the data plotted in the traditional way as a function of angle. Also included are previously reported data for 34 MeV ⁷Li scattering from ¹²C [8]. It is quite difficult to discern any pattern in the displayed data, primarily because the oscillations



Fig. 1. Experimental data for the differential cross section, vector analyzing power ${}^{T}T_{10}$ and second rank tensor analyzing power ${}^{T}T_{20}$ for ${}^{7}\vec{\text{Li}}$ elastic scattering from ${}^{6}\text{Li}$, ${}^{7}\text{Li}$ and ${}^{12}\text{C}$ [8] plotted versus angle (centre of mass) and the momentum transfer. Error bars are omitted for clarity and the lines are drawn to guide the eye.

in the data are determined by the size of the target nuclei and the energy of the beam relative to the Coulomb barrier, and occur at very different angles.

It has been found useful in searching for patterns when comparing elastic scattering between the same target and beam but at multiple bombarding energies to display the data as a function of momentum transfer rather than angle [13], and this was done for the present data sets. The momentum transfer was calculated from $q_t = 2k \cdot \sin(\theta_{\text{c.m.}}/2)$ where k is the wave number of the projectile.

Fig. 1 shows that the structures of the cross sections and vector analyzing powers are target independent for momentum transfers up to 1.0 fm⁻¹, while the second rank tensor analyzing powers $^{T}T_{20}$ are very similar out to momentum transfer of 1.8 fm⁻¹. With this observation, it should be possible through detailed calculations to separate the projectile dependent and projectile-target

	V	r_V	a_V	W	r_W	a_W	$V_{\rm so}$	$r_{\rm so}$	$a_{\rm so}$
⁶ Li	107.8	0.750	0.853	37.9	1.000	0.757	2.5	0.950	0.45
$^{7}\mathrm{Li}$	107.8	0.750	0.855	37.9	0.910	0.757	2.5	0.950	0.45
$^{12}\mathrm{C}$	107.8	0.750	0.800	47.9	1.000	0.757	1.75	0.926	0.45

Table 1

Optical model parameters for ⁷Li elastic scattering from ⁶Li and ⁷Li at 42 MeV and from ¹²C at 34 MeV. For all radii the convention $R_i = r_i(A_t^{1/3} + A_p^{1/3})$ is used.

dependent contributions to the scattering process, and then extract projectile structure information.

Such calculations were performed within the framework of the coupled channels (CC) method. Standard optical model (OM) potentials of Woods-Saxon form with volume absorption were used and the parameters are presented in Table 1. The potential parameters were taken from ref. [14], and were slightly modified in order to obtain a better description of the experimental data. The CC calculations were carried out using the code FRESCO [15], version FRXY.1i. The rôle of different processes in the ⁷Li interaction was investigated using various coupling schemes. The results of the calculations are shown in Figs. 2-4.

The first calculation investigated the rôle of the ground state reorientation coupling in the projectile. The standard rotational model was employed with coupling strengths taken from ref. [3]. The observed $^{T}T_{20}$ analyzing powers are reproduced by this process in the forward angular region (dashed curves in Fig. 2) up to $\theta = 60^{\circ}$ (or $q_t = 1.5 \text{ fm}^{-1}$ momentum transfer) for scattering from ⁶Li and ¹²C, and up to $\theta \sim 40^{\circ}$ ($q_t \sim 1.3 \text{ fm}^{-1}$) for scattering from ⁷Li (in the region of the first two maxima, in agreement with previous work on ⁷Li scattering on ⁴He [7], ¹²C [8], and ²⁶Mg [16]). The projectile ground state reorientation process also partly reproduces the $^{T}T_{10}$ analyzing powers, providing a rather good description at small momentum transfers, but the amplitudes of the oscillations at the larger momentum transfers are small compared to the data.

To understand whether the spin-orbit potential gives a significant contribution to the scattering process, further calculations were performed with a real spin-orbit interaction included and the results are also shown in Fig. 2 (solid curves). As can be seen, while the inclusion of the spin-orbit potential has essentially no effect on $^{T}T_{20}$, it significantly increases the magnitude of the oscillations in $^{T}T_{10}$, particularly at larger momentum transfers, except for the 12 C target where it has little effect. This result is an obvious indication that the spin-orbit interaction, while weak, can still make an important contribution to the scattering process. The lack of effect for the 12 C target is at least partly due to the weaker potential employed for this target – the use of a



Fig. 2. Cross section, vector analyzing power ${}^{T}T_{10}$ and second rank tensor analyzing power ${}^{T}T_{20}$ momentum transfer distributions for ⁷Li elastic scattering from ⁶Li and ⁷Li targets at 42 MeV and from a ¹²C target at 34 MeV. The dashed curves represent the results of CC calculations with coupling to the ⁷Li ground state reorientation only. The solid curves denote the results of similar calculations including a spin-orbit potential.

stronger potential destroyed the good agreement with $^{T}T_{10}$ at low momentum transfer generated by the ground state reorientation coupling.

To investigate the rôle of coupling to the excited states of ⁷Li (bound $1/2^{-}$ and $5/2^{-}$ resonances) as well as target-dependent effects, CC calculations with corresponding coupling schemes were performed and their results are shown in Figs. 3-4. Initially, couplings to the excited states of ⁷Li only were included. The rotational model was again employed, and all allowed quadrupole couplings between these states were included. The results are shown in Fig. 3 as the solid and dashed curves for calculations with and without the real spin-orbit potential, respectively. The inclusion of these couplings necessitated a retuning of the optical potential parameters for ⁷Li + ¹²C in order to recover reasonable agreement with the cross section; the new parameters were: V = 107.8 MeV, $r_V = 0.846$ fm, $a_V = 0.80$ fm, W = 31.65 MeV, $r_W = 1.00$ fm, $a_W = 0.757$ fm (the spin-orbit potential parameters were



Fig. 3. Data as in Fig. 2. The solid and dashed curves denote the result of the same calculations as in Fig. 2 with added transition to the 3/2-, 1/2- and 7/2- excited states of ⁷Li with and without the real spin-orbit potential, respectively.

unchanged).

Adding couplings to the excited states of ⁷Li gives a better description of ^T T_{20} for ⁷Li + ⁶Li and ⁷Li + ⁷Li scattering in the region of the second and third maxima while tending to damp out their magnitudes for ⁷Li + ¹²C, giving a worse description of this observable. However, they do not play any significant rôle in producing the second rank analyzing powers in the momentum transfer region up to 1.5 fm⁻¹. Including these processes in the coupling scheme tends to worsen the agreement between the calculated and measured vector analyzing powers, even when the OM potential with the real spin-orbit potential is used (dashed and solid curves in Fig. 3). We again see that the inclusion of a real spin-orbit potential has little or no influence on ^T T_{20} , its main effect being on ^T T_{10} for momentum transfers of greater than about 1.0 fm⁻¹. Overall, we may conclude that the effect of couplings to the excited states of ⁷Li on the first and second rank analyzing powers is relatively unimportant compared to that due to ground state reorientation, with the exception of ^T T_{20} at momentum transfers greater than about 1.5 fm⁻¹ for ⁷Li + ⁷Li scattering.

Couplings to the first excited states of the targets do not play an important

rôle at small momentum transfers in the ⁷Li-target interactions studied here and their inclusion only slightly modifies the analyzing powers in the range $0 \le q_t \le 1.0 \text{ fm}^{-1}$, as shown in Fig. 4 by the solid curves. Adding couplings to the excited states of the targets again gives a somewhat worse description of the experimental data in comparison to previous calculations. This could be due to the increase in the number of exact values which are needed to perform the theoretical calculations. However, the description obtained is good enough to define the region in the momentum transfer distribution where the dominant rôle is played by projectile structure. Couplings for ⁷Li and ¹²C used the rotational model, the coupling strengths for ¹²C being taken from ref. [17], while the couplings for ⁶Li followed the method in ref. [18]. The "target" excitation coupling for the $^{7}Li + ^{7}Li$ system also included ground state reorientation. The ${}^{7}\text{Li} + {}^{12}\text{C}$ potentials were again adjusted in order to recover a reasonable description of the cross section data, the new parameters being: V = 81.92 MeV, $r_V = 0.850$ fm, $a_V = 0.80$ fm, W = 14.59 MeV, $r_W = 1.00$ fm, $a_W = 1.10$ fm (the spin-orbit potential parameters were again unchanged). The large imaginary diffuseness a_W is required if a reasonable description of the data is to be obtained, χ^2 minimizations confirming that this large diffuseness cannot be compensated for by an increase in the imaginary radius r_W when coupling to the ¹²C 2_1^+ state is included.

Including these target excitations in the coupling schemes tends to damp out the oscillations in both the cross sections and analyzing powers for all ⁷Li scattering reactions analyzed here (cf. the dashed and solid curves in Fig. 4). In general, one may conclude that target excitation effects are only significantly manifest for momentum transfers greater than about 1.0 fm⁻¹. It should be manifest that while the ground state reorientation coupling for ⁷Li as a *projectile* is very important, particularly for the generation of the second rank tensor analysing power $^TT_{20}$, when this coupling is included for the ⁷Li *target* it has little or no effect. This remarkable result suggests that, at least for small momentum transfers, the target is very much a "spectator" as far as the polarisation observables are concerned.

In conclusion, ⁷Li elastic scattering cross section and analyzing power data when graphed as a function of momentum transfer rather than angle, show similar structure for momentum transfers less than 1.0 fm⁻¹ in the nuclear scattering energy regime studied here. While the internal structures of ⁶Li and ⁷Li have been shown to strongly influence elastic scattering when they are used as beams, their influence on the scattering is not readily apparent when they are used as targets. Although the analyzing powers induced by polarized beams of Li have provided a rich environment for investigating the effects of projectile excitation and reorientation, a detailed picture of the scattering has not emerged for larger momentum transfers at energies in the nuclear scattering regime. The calculations carried out to date do not show any clear dependence on the excitation of the target for any system studied. This result



Fig. 4. Data as in Fig. 2. The curves denote the result of the CC calculations with transition to the 3/2-, 1/2- and 7/2- excited states of ⁷Li couplings only (dashed line) as well as added coupling to excited states of targets (solid line).

suggests that the properties of loosely bound projectiles such as ⁶He, ⁸B and ¹¹Li could be probed by elastic scattering no matter what target is used if the momentum transfers are not too large.

This work was supported in part by the State of Florida, the U.S. National Science Foundation and NATO.

References

- [1] J.S. Blair, Phys. Rev. 115 (1959) 928.
- [2] L.A. Parks, K.W. Kemper, A.H. Lumpkin, R.I. Cutler, L.H. Harwood, D. Stanley, P. Nagel, and F. Petrovich, Phys. Lett. 70B (1977) 27.
- [3] V. Hnizdo, K.W. Kemper, J. Szymakowski, Phys. Rev. Lett. 46 (1981) 590.
- [4] G. Tungate, R. Böttger, P. Egelhof, K.-H. Möbius, Z. Moroz, E. Steffens, W. Dreves, I. Koenig, and D. Fick, Phys. Lett. 98B (1981) 347.

- [5] K. Rusek, Z. Moroz, R. Čaplar, P. Egelhof, K.-H. Möbius, E. Steffens, I. Koenig, A. Weller, and D. Fick, Nucl. Phys. A407 (1983) 208.
- [6] Z. Moroz, P. Zupranski, R. Böttger, P. Egelhof, K.-H. Möbius, G. Tungate, E. Steffens, W. Dreves, I. Koenig, and D. Fick, Nucl. Phys. A381 (1982) 294.
- [7] P.D. Cathers, E.E. Bartosz, M.W. Cooper, N. Curtis, N. Keeley, K.W. Kemper, F. Maréchal, E.G. Myers, B.G. Schmidt, K. Rusek, and V. Hnizdo, Phys. Rev. C 63 (2001) 064601.
- [8] E.E. Bartosz, N. Keeley, P.D. Cathers, M.W. Cooper, K.W. Kemper, F. Maréchal, and K. Rusek, Phys. Rev. C 64 (2001) 014606.
- [9] H. Amakawa and K.I. Kubo, Nucl. Phys. A266 (1976) 521.
- [10] A.J. Mendez, E.G. Myers, K.W. Kemper, P.L. Kerr, E.L. Reber, and B.G. Schmidt, Nucl. Instrum. Methods Phys. Res. A 329 (1993) 37.
- [11] G. Tungate, W. Ott, and H. Leucker, Z. Phys. A 321 (1985) 103.
- [12] H.G. Bingham, A.R. Zander, K.W. Kemper, and N.R. Fletcher, Nucl. Phys. A173 (1971) 265.
- [13] Dao T. Khoa, W. von Oertzen, H.G. Bohlen, and F. Nuoffer, Nucl. Phys. A672 (2000) 387.
- [14] J. Jänecke, T. Annakkage, G.P.A. Berg, B.A. Brown, J.A. Brown, G. Crawley, S. Danczyk, M. Fujiwara, D.J. Mercer, K. Pham, D.A. Roberts, J. Stasko, J.S. Winfield, and G.H. Yoo, Phys. Rev. C 54 (1996) 1070.
- [15] I.J. Thompson, Comp. Phys. Rep. 7 (1988) 167.
- [16] W. Ott, R. Butsch, H.J. Jänsch, K.-H. Möbius, P. Paul, G. Tungate, E. Steffens, K. Rusek, Z. Moroz, I.M. Turkiewicz, K. Becker, K. Blatt, H. Leucker and D. Fick, Nucl. Phys. A489 (1988) 329.
- [17] M.F. Vineyard, J. Cook, and K.W. Kemper, Phys. Rev. C 31, (1985) 879.
- [18] E.L. Reber, K.W. Kemper, P.V. Green, P.L. Kerr, A.J. Mendez, E.G. Myers, B.G. Schmidt, and V. Hnizdo, Phys. Rev. C 50 (1994) 2917.