

# Prior to “Quark Matter 2006” predictions within retarded jet absorption scenario at RHIC

V.S. Pantuev

University at Stony Brook, Stony Brook, NY 11794-3800

On leave from Institute for Nuclear Research Russian Academy of Sciences, Moscow

## Abstract

Predictions for some experimental physical observables in nucleus-nucleus collisions at RHIC energies are presented. I utilize the previous suggestion that the retarded, by time about 2-3 fm/c, jet absorption in opaque core is a natural explanation of many experimental data. This assumption is applied only to the particles with high transverse momentum above 4 GeV/c resulting from parton fragmentation. I calculate nuclear modification factor  $R_{AA}$ ,  $R_{AA}$  in- and out of the reaction plane, azimuthal anisotropy parameter  $v_2$ , jet suppression  $I_{AA}$  for the away side jet and its dependence versus the reaction plane orientation. The systems under consideration are Au+Au, Cu+Cu, at 200 GeV and 62 GeV. Most of numbers are predictions prior to the QM2006.

PACS numbers: 25.75.Dw

At QM2006 conference we expect that many new experimental data will be shown. I already get few requests from different groups to calculate some experimentally observed parameters. To be fair, I present all these numbers prior to the conference.

In previous papers [1, 2] I propose a simple model to explain experimental data on the angular dependence of the nuclear modification factor  $R_{AA}$  at high transverse momentum in the reaction plane. The model has one free parameter  $L \simeq 2$  fm to describe the thickness of the corona area with no absorption and was adjusted to fit the experimental data on AuAu collisions at centrality 50-60%. The model nicely describes the  $R_{AA}$  dependence for all centrality classes. I extract the second Fourier component amplitude,  $v_2$ , for high  $p_T$  particle azimuthal distribution and found  $v_2$  should be at the level of 11-12% purely from the geometry of the collision with particle absorption in the core. At that time I made a prediction for  $R_{AA}$  in Cu+Cu collisions at 200 GeV which, as later was found, is in very good agreement with experimental data. Physical interpretation of the parameter  $L$  could be that it is actually retarded jet absorption caused by the plasma formation time  $T = L/c \simeq 2$  fm/c, or at least non-trivial response of strongly interacting plasma to fast moving color charge.

This value of formation time is true for the dilute corona region where density is small. For more central interaction region it should be smaller, probably changing like mean distance between interaction points or  $1/\sqrt{\rho(x, y)}$ , where  $\rho(x, y)$  is two-dimensional (projected into  $x - y$  plane) density of  $N_{binary}$  or  $N_{part}$ . Extrapolation of  $T$  in a such way gives values of the order of 0.7-1 fm/c in the center of collision zone.

As usual, nuclear modification factor  $R_{AA}$  is defined as:

$$R_{AA}(p_T) = \frac{(1/N_{evt}) d^2 N^{A+A}/dp_T d\eta}{(\langle N_{binary} \rangle / \sigma_{inel}^{N+N}) d^2 \sigma^{N+N}/dp_T d\eta}, \quad (1)$$

where  $\langle N_{binary} \rangle$  is a number of binary nucleon-nucleon collisions at particular centrality class.

I use a Monte Carlo simulation of nucleus-nucleus collisions based on the Glauber approach with Woods-Saxon density distribution. Assume that jets are produced in accordance with the  $N_{coll}$  distribution in the  $x - y$  transverse plane. I neglect longitudinal expansion for the first 2-3 fm/c. Let all jets fly in all possible directions (some of them will move from the interaction zone determined by the envelop of Woods-Saxon radii, some will go into this zone) for some time  $T$ . If after that time jet will be still inside the interaction zone, it will be completely absorbed. If not, jet leaves the zone without interaction. We found from ex-

perimental data that this time should be about 2.3 fm/c at 200 GeV and 3.5 fm/c at 62 GeV center of mass beam energy. Absorption can not happen instantly (or in the infinitely thin layer), plus there should be statistical fluctuations. To take these into account I smooth the cut edge: an arbitrary weight function in the form of a Fermi distribution with diffuseness parameter  $a$  was applied:

$$weight(l) = \frac{1}{1 + exp((T - l)/a)}. \quad (2)$$

The value of this function changes from 0.1 to 0.9 at  $\Delta l=4.4*a$  around  $T$ . Parameter  $a$  varied from 0.01 to 0.5 fm/c. It was found that the results for  $R_{AA}$  and  $v_2$  do not change much for this parameter range. By default it was chosen to be  $a=0.2$  fm/c. The results of calculation for  $R_{AA}$  are presented in Fig. 1. All values for  $R_{AA}$  and  $v_2$  are presented in tables at the bottom of this preprint. It is worth to mention, values of  $R_{AA}$  and  $v_2$  for mesons and baryons at high  $p_T$  should be the same. Obviously,  $R_{AA}$  is flat versus  $p_T$  if no other physics process is “switched on” at high  $p_T$ , except retarded absorption in opaque core considered here.

For di-jets, another parameter,  $I_{AA}$ , was introduced in Ref. [5]. For away side jet  $I_{AA}$  is the ratio of particle yield per trigger in the particular momentum range of trigger and associate particles, to the similar yield in p+p collisions. To calculate this value I perform the same procedure like for  $R_{AA}$ , except that in addition the location of away side jet after time  $T$  should be taken into account. Away side jet may not go exactly in opposite relative to the trigger jet direction, but is distributed around 180 degree angle. It introduces another critical parameter for the calculation, which should be taken from the experimentally observed width of the away side jets in azimuthal angle. I use simple Gausse distribution around 180 degrees for the away side jet. Depending on the momentum range of the associate particle, width of the away side jet varies from 0.5 radians for the low momentum range to 0.2 radians at momentum above 5-6 GeV/c.

Results of calculation for  $I_{AA}$  in- and out of plane for Au+Au collisions at 200 GeV are presented in Fig. 2. One can see the main feature of the surface biased scenario: in the most central events there are more di-jets out of the reaction plane than di-jets in the plane. This is opposite to the  $R_{AA}$  reaction plane dependence or *to the scenario which assumes only “punch-through” di-jets*. In mid-central events the situation is mixed, and in peripheral collisions there are more in-plane di-jets. Wide centrality bin of 20 to 60%, commonly used

in experiments, corresponds to  $N_{part}=125$  if number of di-jets is weighted by the number of trigger particles. Results for Cu+Cu collisions are shown in Fig. 3.

I want to emphasize that the role of event by event fluctuations should be treated by a better way than I do it here by parameter  $a$ . It is known that nucleon position fluctuations in the colliding Cu+Cu nuclei can significantly increase the observed  $v_2$  in the most central collisions [6]. This is why static models, including our model presented here, underestimate  $v_2$  in the central Cu+Cu collisions. The same should be true when we consider di-jets in Au+Au or Cu+Cu: local density fluctuations induce positive correlation between surviving probability of the trigger jet and the away side jet. Again, such fluctuations could increase the observed value of  $I_{AA}$ , especially in the central collisions.

Few more comments: as I pointed out in the previous paper [2], if we see a significant suppression of non-photonic electrons [7], one can expect values of  $v_2$  for such electrons at high  $p_T$  to be equal to  $v_2$  for high  $p_T$  hadrons. Another point: suppression of the away side jets in direct gamma-hadron correlations,  $I_{AA}$ , should be equal to the  $R_{AA}$  of inclusive hadrons at about gamma energy. Here I ignore the difference in quark and gluon fragmentation functions. The reason, that this may change the result, is that hadrons in the away side jets for the direct gamma-hadron correlations are mostly from quark fragmentation, while inclusive hadrons below 10 GeV/c are dominantly from gluon fragmentation.

All numbers for Au+Au collisions are presented at 200 GeV in Table I, at 62 GeV in Table II. Numbers for Cu+Cu collisions at 200 GeV are in Table III, and at 62 GeV are in Table IV.

- 
- [1] V. S. Pantuev, hep-ph/0506095.
  - [2] V. S. Pantuev, hep-ph/0604268.
  - [3] S. S. Adler et al., [PHENIX Collaboration], *Phys. Rev.* **C69** (2004) 034910.
  - [4] M. Shumomura, [PHENIX collaboration], *Nucl. Phys.* **A774** (2006) 457, arXiv:nucl-ex/0510023.
  - [5] J. Adams et al., [STAR collaboration], *Phys. Rev. Lett.* **93** (2004) 252901.
  - [6] S. Manly, [PHOBOS collaboration], *Nucl. Phys.* **A774** (2006) 523, arXiv:nucl-ex/0510031.
  - [7] S. Butsyk, [PHENIX collaboration], *Nucl. Phys.* **A774** (2006) 669, arXiv:nucl-ex/0510010.

## Figures




FIG. 1: Calculated  $R_{AA}$  for Au+Au and Cu+Cu collisions at 200 GeV and 62 GeV versus the number of participant nucleons,  $N_{part}$ . Solid curves are for 200 GeV, dashed are for 62 GeV. The circles are experimental  $\pi^0$  data for 200 GeV Au+Au collisions integrated for  $p_T \geq 4$  GeV/c [3]. The triangles are data for 200 GeV Cu+Cu collisions of  $p_T \geq 7$  GeV/c [4]. Only statistical errors are shown.




FIG. 2: Calculated  $I_{AA}$  in- and out of plane for Au+Au collisions at 200 GeV versus the number of participant nucleons,  $N_{part}$ . Width of the away side jet is  $\sigma=0.22$  radians. Small triangles are estimation for in plane, squares- for out of plane. The circles with error bars are data from [5].




FIG. 3: Calculated  $I_{AA}$  for Cu+Cu collisions at 200 GeV versus the number of participant nucleons,  $N_{part}$ . Width of the away side jet is  $\sigma=0.22$  radians. Circles are estimation for in plane, triangles - for out of plane.

## Tables

TABLE I: Results of calculations for Au+Au at 200 GeV. In all calculations  $a=0.2$  fm/c and  $\sigma=0.22$  radians.

cent, %	$N_{part}$	R	$R_{in}$	$R_{out}$	$v_2, \%$	$\langle I_{AA} \rangle$	$I_{in}$	$I_{out}$
0-5	353	0.195	0.213	0.177	4.71	0.145	0.0882	0.202
5-10	300	0.235	0.281	0.211	7.06	0.198	0.135	0.262
10-15	255	0.29	0.348	0.246	8.64	0.259	0.204	0.314
15-20	215	0.335	0.417	0.282	9.68	0.34	0.317	0.364

20-25	182	0.39	0.483	0.316	10.47	0.448	0.488	0.407
25-30	152	0.45	0.563	0.362	10.87	0.575	0.687	0.463
30-35	126	0.505	0.632	0.40	11.23	0.659	0.809	0.508
35-40	103	0.565	0.705	0.447	11.19	0.727	0.897	0.557
40-45	84	0.635	0.779	0.512	10.37	0.783	0.947	0.62
45-50	67	0.715	0.848	0.589	9.03	0.833	0.974	0.69
50-55	52	0.79	0.902	0.681	7.0	0.877	0.99	0.763
55-60	40	0.84	0.958	0.847	5.53			
cent, %	$N_{part}$	R	$R_{in}$	$R_{out}$	$v_2, \%$	$\langle I_{AA} \rangle$	$I_{in}$	$I_{out}$

TABLE II: Results of calculations for Au+Au at 62 GeV.

0-5	345	0.333	0.364	0.302	4.65			
5-10	296	0.402	0.461	0.344	7.29			
10-15	250	0.468	0.553	0.383	9.06			
15-20	211	0.532	0.639	0.425	10.			
20-25	178	0.591	0.719	0.462	10.9			
25-30	149	0.656	0.801	0.512	11.			
30-35	123.8	0.707	0.861	0.553	10.9			
35-40	102	0.756	0.911	0.601	10.3			
40-45	82.9	0.805	0.948	0.663	8.83			
45-50	66.2	0.851	0.971	0.731	7.05			
50-55	51.9	0.895	0.984	0.805	5.01			
55-60	40	0.922	0.99	0.854	3.69			
cent, %	$N_{part}$	R	$R_{in}$	$R_{out}$	$v_2, \%$			

TABLE III: Results of calculations for Cu+Cu at 200 GeV.



cent, %	$N_{part}$	R	$R_{in}$	$R_{out}$	$v_2, \%$	$\langle I_{AA} \rangle$	$I_{in}$	$I_{out}$
0-5	110	0.37	0.40	0.34	3.4	0.194	0.136	0.251
5-10	95	0.435	0.48	0.39	5.1	0.263	0.203	0.324
10-15	82	0.51	0.57	0.45	6.0	0.346	0.31	0.382
15-20	70.5	0.585	0.66	0.51	6.3	0.482	0.497	0.467
20-25	60	0.655	0.74	0.57	6.4	0.581	0.637	0.526
25-30	51	0.7	0.79	0.61	6.4	0.645	0.723	0.568
30-35	43	0.775	0.86	0.69	5.4	0.733	0.817	0.648
35-40	36	0.825	0.9	0.75	4.3	0.784	0.865	0.704
40-45	29	0.87	0.93	0.81	3.3	0.833	0.903	0.762
45-50	24	0.91	0.95	0.87	2.1			
50-55	19	0.95	0.97	0.93	0.9			
55-60	15	0.98	0.98	0.98	0			

cent, %	$N_{part}$	R	$R_{in}$	$R_{out}$	$v_2, \%$	$\langle I_{AA} \rangle$	$I_{in}$	$I_{out}$
---------	------------	---	----------	-----------	-----------	--------------------------	----------	-----------

TABLE IV: Results of calculations for Cu+Cu at 62 GeV.

cent, %	$N_{part}$	R	$R_{in}$	$R_{out}$	$v_2, \%$
0-5	104	0.598	0.638	0.559	3.28
5-10	92	0.672	0.737	0.607	4.82
10-15	80	0.736	0.817	0.656	5.48
15-20	68	0.8	0.888	0.712	5.48
20-25	57	0.845	0.931	0.759	5.08
25-30	48.6	0.874	0.954	0.793	4.61
30-35	40.6	0.91	0.975	0.846	3.52
35-40	34	0.932	0.982	0.882	2.7
40-45	28	0.952	0.989	0.915	1.93

cent, %	$N_{part}$	R	$R_{in}$	$R_{out}$	$v_2, \%$
---------	------------	---	----------	-----------	-----------