The anisotropic flow coefficients v_2 and v_4 in Au+Au collisions at RHIC

Yuting Bai for the STAR[‡] Collaboration

NIKHEF, Kruislaan 409, 1098SJ Amsterdam, The Netherlands

E-mail: Yuting.Bai@nikhef.nl

Abstract. We present measurements by STAR of the anisotropic flow coefficients v_2 and v_4 as a function of particle-type, centrality, transverse momentum and pseudorapidity in Au+Au collisions at RHIC.

Anisotropic flow is an azimuthal correlation of the particle momenta with respect to the reaction plane. This flow is recognized as one of the main observables that provide information on the early stage of a heavy-ion collision [1].

In this study, we used 13×10^6 minimum-bias Au+Au events taken at a center-ofmass energy of 200 GeV and 6×10^6 events taken at 62.4 GeV. The particles were detected by the STAR main TPC [2] and by the forward TPCs [3] and cover a pseudorapidity of $|\eta| < 1.3$ and $2.5 < |\eta| < 4.0$, respectively. From these data, v_2 is obtained by the 4-particle cumulant method [4] and is denoted by v_2 {4}. This method is less sensitive to non-flow effects compared to measurements based on two particle correlations like v_2 {2} or v_2 { EP_2 }. The v_4 coefficient is obtained with respect to the second harmonic event plane and is denoted by v_4 { EP_2 }. The flow coefficients are studied for different particle species as function of transverse momentum (p_t), pseudorapidity (η) and centrality. Only statistical errors are shown unless specified.

Transverse momentum dependence. The left panel of Fig. 1 shows the charged particle v_2 as a function of p_t for mid-central (20–60%) Au+Au collisions at 200 and 62.4 GeV. It is seen that the measured v_2 increases with p_t , reaches its maximum around 3 GeV/c and then decreases again. At 200 GeV, v_2 is measured up to 10 GeV/c and is still sizable above 8 GeV/c. The behavior of v_2 at 62.4 GeV is similar to that observed at 200 GeV. It is argued in [5] that v_2 at large p_t might be related to the parton energy loss mechanism and may thus provide a constraint on the initial gluon density.

The p_t dependence of the charged particle v_4 is shown for both energies in the right panel of Fig. 1. It is seen that v_4 increases quadratically at low p_t and has, like v_2 , its maximum around 3 GeV/c. At 200 GeV, v_4 is measured up to 7 GeV/c and is still sizable above 6 GeV/c. Similar values are obtained at 62.4 GeV.

[‡] For the full list of STAR authors and acknowledgements, see appendix 'Collaborations' in this volume.



Figure 1. Charged particle $v_2\{4\}$ (left-hand plot) and $v_4\{EP_2\}$ (right-hand plot) versus p_t for 20–60% centrality at $|\eta| < 1.3$ in Au + Au collisions at 200 and 62.4 GeV.



Figure 2. Charged particle $v_2\{4\}$ and $v_4\{EP_2\}$ versus $\eta - y_{beam}$ for different centrality bins at 200 (full circles) and 62.4 GeV (open circles). The flow coefficients are shown for particles in the forward hemisphere only.

Rapidity dependence. It has been shown that particle production in the fragmentation region exhibits longitudinal scaling when plotted as a function of $\eta - y_{beam}$ [6]. It is also known that the integrated elliptic flow for fixed centrality at mid-rapidity is proportional to the particle yield dN/dy [7]. If this scaling with dN/dy holds at all rapidities, then v_2 is also expected to show a longitudinal scaling behavior. Figure 2 (left) shows v_2 as a function of $\eta - y_{beam}$ for different centralities at 200 and 62.4 GeV. The v_2 values measured at both energies fall on a universal curve, indicating that the longitudinal scaling approximately holds. This scaling is also observed for v_4 as can be seen in the right panel of Fig. 2.

Mass and particle type dependence. It has been shown by STAR [8] that v_2 for identified particles at low p_t exhibits a mass ordering. This ordering is well described by hydrodynamic calculations which indicates that all particles flow with a common



Figure 3. The scaled flow coefficients v_2/n_q versus p_t/n_q and $(m_T - m_0)/n_q$. Left: v_2/n_q for 3 different centrality classes at $|\eta| < 1.0$ measured at 200 GeV. Right: v_2/n_q versus $(m_T - m_0)/n_q$ for minbias data at 62.4 GeV for $|\eta| < 1.0$. The curve shows a polynominal fit to the data.

velocity. At intermediate p_t , v_2 scales with the number of constituent quarks n_q [9]. This scaling can be explained in the coalescence picture and is indicative of the partonic origin of flow [10]. The ratio v_2/n_q is shown in Fig. 3 (left) for three different centralities at 200 GeV as a function of p_t/n_q and as a function of the scaled transverse mass $(m_T - m_0)/n_q$. The scaled transverse mass, which takes into account relativistic effects, is sometimes considered to be a better scaling variable than p_t/n_q [11].

An indication that v_2 follows constituent quark scaling independent of centrality is given by the fact that the v_2/n_q falls onto a universal curve for each centrality bin. Figure 3 shows that the constituent quark scaling holds at both 200 GeV and 62.4 GeV.

The ratio v_4/v_2^2 In recent work [12], v_4/v_2^2 is proposed as a more sensitive probe of ideal hydrodynamic behavior. Furthermore, this ratio is directly related to the degree of thermalization of the medium, see also [13]. Under the assumption that $v_2\{4\}$ is a genuine measure of elliptic flow, the systematic error in the ratio v_4/v_2^2 is dominated by non-flow contributions to v_4 . It can be shown that the non-flow contribution to v_4 is proportional to the difference $v_2\{2\}^2 - v_2\{4\}^2$. Obtaining this difference from the data allows us to estimate the systematic error on v_4/v_2^2 . However, the difference can also originate from flow fluctuations [14]. In that case, the systematic error will be reduced. A detailed description of this systematic analysis is beyond the scope of these proceedings and will be described in a future publication.

Figure 4 shows the ratio v_4/v_2^2 measured at 200 GeV in the 20–30% centrality interval. The horizontal brackets in this figure show the lower limit of the systematic uncertainty as presently estimated. In this figure, the data are compared to two



Figure 4. The ratio of v_4/v_2^2 versus p_t for charged particles at $|\eta| < 1.3$ in Au+ Au collisions at 200 GeV. The horizontal brackets indicate the systematic uncertainty. The curves correspond to two hydrodynamic calculations [15, 12]. The filled area shows the AMPT model calculations [16].

hydrodynamic model calculations [12, 15] (curves) and to model predictions based on a microscopic partonic and hadronic description of the collision (AMPT model, filled area) [16]. It is seen that the data lie above the model predictions. However, the present systematic uncertainties do not allow us to either validate or exclude these two models.

References

- [1] STAR Collaboration, Adams J et al., 2005 Nucl. Phys. A 757 102, and references therein.
- [2] Anderson M et al., 2003 Nucl. Instr. Meth. A 499, 659.
- [3] Ackermann K H et al., 2003 Nucl. Instr. Meth. A 499, 713.
- [4] Borghini N, Dinh P M and Ollitrault J-Y, 2001 Phys. Rev. C 64 054901.
- Snellings R, Poskanzer A M and Voloshin S A, nucl-ex/9904003; Gyulassy M, Vitev I and Wang X N, 2001 Phys. Rev. Lett. 86 2537.
- [6] PHOBOS Collaboration, Back B B et al., 2006 Phys. Rev. C 74 021901.
- [7] STAR Collaboration, Adler C et al., 2002 Phys. Rev. C 66, 034904.
- [8] STAR Collaboration, Adler C et al., 2001 Phys. Rev. Lett. 87, 182301.
- [9] STAR Collaboration, Adams J et al., 2004 Phys. Rev. Lett. 92, 052302.
- [10] Molnar D, Voloshin S A, 2003 Phys. Rev. Lett. 91 092301.
- [11] PHENIX Collaboration, Adare A et al., nucl-ex/0608033.
- [12] Borghini N, Ollitrault J-Y, 2006 Phys.Lett. B 642, 227-231.
- [13] Ko C M, in these proceedings;
- [14] Miller M, Snellings R, nucl-ex/0312008; Sorensen P, in these proceedings; Tang A H, in these proceedings; Voloshin S A, in these proceedings.
- [15] Kolb P F, 2003 Phys. Rev. C 68, 031902.
- [16] Chen L, Ko C M and Lin Z, 2004 Phys. Rev. C 69, 031901.