

# Azimuthal and pseudo-rapidity correlations with strange particles at intermediate- $p_T$ at RHIC

**J Bielcikova for the STAR Collaboration**

Physics Department, Yale University, New Haven, CT-06520, USA

E-mail: jana.bielcikova@yale.edu

**Abstract.** We present results on two-particle azimuthal correlations with strange trigger particles ( $K_S^0$ ,  $\Lambda$ ,  $\Xi$ ,  $\Omega$ ) associated with unidentified charged particles in d+Au and Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV. We investigate, in detail, the near-side associated yield as a function of centrality,  $p_T$  and strangeness content in the trigger particle to look for possible flavor and baryon/meson differences. We compare our results to a fragmentation and recombination model, where the study of  $\Omega$ -triggered correlations is used as a critical test of the validity of the recombination picture.

PACS numbers: 25.75.-q, 25.75.Gz


## 1. Introduction

The suppression of inclusive  $p_T$  spectra of identified particles in central Au+Au collisions with respect to p+p and peripheral Au+Au collisions [1, 2] and the enhanced baryon/meson ratios [3, 4] show that baryons and mesons behave differently than in p+p collisions for  $p_T = 2-6$  GeV/ $c$ . This indicates that fragmentation is not dominant and parton recombination and coalescence models [5, 6, 7, 8] have been suggested as alternative mechanisms of particle production. Moreover, studies of di-hadron correlations in Au+Au revealed the presence of an additional long-range pseudo-rapidity correlation (*ridge*) on the near-side [9], which is absent in p+p and d+Au collisions.

In this paper, we discuss the properties of two-particle correlations using strange trigger particles ( $K_S^0$ ,  $\Lambda$ ,  $\Xi$ ,  $\Omega$ ) associated with unidentified charged particles in d+Au and Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV with STAR. We investigate the near-side associated yield as a function of centrality, system size,  $p_T$  and strangeness content in the trigger particle, to look for possible flavor and baryon/meson differences. We compare our results to a fragmentation and recombination model.

## 2. Data analysis

The correlation functions, normalized to the number of trigger particles, are corrected for the reconstruction efficiency of associated particles, elliptic flow ( $v_2$ ) and, unless mentioned otherwise, triangular acceptance in  $\Delta\eta$ . The near-side yield of associated



**Figure 1.** Centrality dependence of the ridge yield (a) and jet yield (b) of associated charged particles for various trigger species in d+Au and Au+Au collisions. The error bands indicate systematic errors on the ridge yield due to the  $v_2$  subtraction.


particles is calculated as the area under the Gaussian peak obtained from a fit. We study separately the jet and ridge contributions to the near-side yield by analyzing the correlations in two  $\Delta\eta$  windows:  $|\Delta\eta| < 0.7$  containing both jet and ridge correlations, and  $|\Delta\eta| > 0.7$  containing only the ridge contributions, assuming the jet contribution at large  $\Delta\eta$  is negligible. The jet yield is free of systematic uncertainties due to the  $v_2$  subtraction if a uniformity of  $v_2$  with  $\eta$  is assumed. For the ridge yield, these systematic errors are estimated by subtracting the  $v_2$  measured by the event plane method (the lower bound) and by the 4-particle cumulant method (the upper bound).

### 3. Correlations with $K_S^0$ , $\Lambda$ and charged trigger particles

Comparing the near-side yields in d+Au and Au+Au collisions, we observe a strong increase by a factor of 3-4 going from d+Au to central Au+Au collisions. Studying separately the jet and ridge contributions to the near-side yield (Fig. 1), we find that the ridge yield rises with centrality and is responsible for the observed strong increase of the near-side yield with centrality. The jet yield is independent of centrality and consistent with that in d+Au collisions. No significant baryon/meson or particle/anti-particle differences are observed.

Next, we study the dependence of the near-side yield on the transverse momentum of the trigger particle,  $p_T^{trig}$ , shown in Fig. 2. While the ridge yield increases with  $p_T^{trig}$  and flattens off for  $p_T^{trig} > 3.0$  GeV/c, the jet yield keeps increasing with  $p_T^{trig}$  as expected. The jet yield for  $\Lambda$  triggers is systematically below that of charged hadron and  $K_S^0$  triggers. Remaining effects of artificial track merging, which are found to affect more V0s than charged tracks, are currently under investigation.

We have also measured the invariant  $p_T$  spectra of associated charged particles for




**Figure 2.** Dependence of the ridge yield (a) and jet yield (b) on  $p_T^{trig}$  for various trigger species in central (0-10%) Au+Au collisions. The bands indicate systematic errors on the ridge yield due to the  $v_2$  subtraction.

$p_T^{trig} = 3-6$  GeV/c (not shown) and extracted the inverse slope,  $T$ . The ridge spectra have, for all studied trigger species,  $T \sim 400$  MeV, close to that of 'the bulk', while the jet spectra have  $T \sim 450$  MeV. The 0-10%/40-80% centrality ratio of the near-side yields is about 3 at  $p_T^{assoc} = 1$  GeV/c and decreases with  $p_T^{assoc}$ . As shown in Fig. 1, this large ratio is due to the correlations at large  $|\Delta\eta|$  because the jet yield is independent of centrality. Our results qualitatively agree with the parton recombination model [10] which points toward a significant role of thermal-shower recombination in Au+Au collisions. To draw quantitative conclusions, the calculation must be done for the same centrality and  $p_T^{trig}$  selection and also reproduce properties of the measured  $\Delta\eta$  correlations.

#### 4. Correlations with multiply-strange trigger baryons

Our study of correlations with multiply-strange baryons, especially  $\Omega$ , has been stimulated by the predictions from the recombination model of [11]. Contrary to the production of  $K$  and  $\Lambda$ , for particles created exclusively from strange quarks, such as  $\phi$  and  $\Omega$ , the contribution from shower  $s$  quarks should be negligible for  $p_T$  up to 8 GeV/c. Consequently, there should be no associated particles for  $\phi$ - and  $\Omega$ -triggered correlations with  $p_T > 3$  GeV/c because they cannot be distinguished from background.

Figure 3(a) shows the azimuthal correlations for strange trigger baryons with increasing strangeness content:  $\Lambda$ ,  $\Xi$  and  $\Omega$ . Due to limited statistics, the trigger particles have been selected with  $p_T^{trig} = 2.5-4.5$  GeV/c. Clearly, there is a remaining near-side peak above the elliptic flow contribution for all discussed trigger species. The near-side peak persists even if the larger  $v_2$ , determined from the event plane method, is used. Moreover, the strength of the near-side peak is, within errors, independent of the strangeness content in the trigger particle. This is further confirmed by the study of



**Figure 3.** (a) Azimuthal correlations with  $\Lambda$ ,  $\Xi$  and  $\Omega$  for  $p_T^{\text{trig}}=2.5$ - $4.5$  GeV/ $c$  in 0-10% central Au+Au collisions. The lines indicate the  $v_2$  contributions. (b) Dependence of the near-side yield on  $p_T^{\text{trig}}$  for various strange trigger particles in central (0-10%) Au+Au collisions. The bands represent systematic errors due to the  $v_2$  subtraction. No  $\Delta\eta$  acceptance correction has been applied.

the  $p_T^{\text{trig}}$  dependence of the near-side yield shown in Fig. 3(b). Within errors, the yield is consistent for all studied baryon and meson species.

## 5. Conclusions

We have reported results on two-particle correlations with strange trigger particles at intermediate- $p_T$  at RHIC. The correlations reveal a strong contribution from the long-range  $\Delta\eta$  correlations in the near-side. We do not observe any significant baryon/meson differences. The behavior of the central-to-peripheral ratio of the near-side yields agrees qualitatively with a recombination model. However, for the  $\Omega$ -triggered correlations there exist associated particles in the near-side, which disagrees with the model. Studies of jet and ridge yields for  $\Omega$ -triggered correlations as well as correlations with identified associated particles are expected to constrain the origin of these correlations.

## References

- [1] Adler S S *et al* (PHENIX) 2003 *Phys. Rev. Lett.* 91 172301
- [2] Adams J *et al* (STAR) 2004 *Phys. Rev. Lett.* 92 052302
- [3] Adams J *et al* (STAR) nucl-ex/0601042
- [4] Abelev B *et al.* (STAR) 2006 *Phys. Rev. Lett.* 97 152301
- [5] Fries R J, Muller B, Nonaka C and Bass S A 2003 *Phys. Rev.* C68 044902
- [6] Greco V, Ko C M and Levai P 2003 *Phys. Rev.* C68 034904
- [7] Greco V, Ko C M and Levai P 2003 *Phys. Rev. Lett.* 90 202302
- [8] Hwa R C and Yang C B 2003 *Phys. Rev.* C67 034902
- [9] Putschke J (STAR) these proceedings
- [10] Hwa R C and Tan Z 2005 *Phys. Rev.* C72 057902
- [11] Hwa R and Yang C B 2006 nucl-th/0602024