# Nucleon number dependence of the onset of anomalous $J/\psi$ suppression and the dynamics of nuclear collisions

Anna Nogová<sup>a</sup>, Neva Pišútová<sup>b</sup>, and Ján Pišút<sup>b</sup>

 <sup>a)</sup> Institute of Physics of the Slovak Academy of Science, SK-84228 Bratislava,Slovakia
 <sup>b)</sup> Department of Physics, Comenius University, SK-84248 Bratislava, Slovakia

Abstract We point out that the data on the onset of anomalous  $J/\psi$  suppression as a function of nucleon numbers A and B could provide information on the dynamics of nuclear interactions. In particular the models of anomalous  $J/\psi$  suppression by Blaizot and Ollitrault (BO) and by Kharzeev, Lourenco, Nardi and Satz (KLNS) are based on different assumptions on the dynamics of nuclear collisions and lead to different predictions of the dependence of the onset of anomalous  $J/\psi$  suppression on nucleon numbers of colliding nuclei. The data on this onset as function of A and B could become a tool for the study of the dynamics of nuclear collisions and bring further evidence on  $J/\psi$  suppression by new form of hadronic matter, possibly Quark-Gluon Plasma. In particular we propose to study  $J/\psi$  suppression in A+Pb interaction with nucleon number of A between 58 and 73 or a bit higher.

### 1 Introduction

Experimental data of the NA38 and NA50 Collaborations at the CERN SPS [1, 2, 3, 4] on  $J/\psi$  suppression has shown that the anomalous  $J/\psi$  suppression is observed in near central Pb+Pb collisions but it is absent in peripheral Pb+Pb and in interactions induced by ions lighter (or equal) to Sulphur.

Non- anomalous suppression of  $J/\psi$  has been phenomenologically well described by the desintegration of  $J/\psi$  by nucleons present in nuclei [5, 6].

There have appeared numerous attempts to describe the anomalous suppression of  $J/\psi$ . We shall not try to make here a complete list of these attempts. A review of approaches based on  $J/\psi$  suppression by comovers can be found in Refs.[7, 8]. For studies based on the initial state interactions see Refs.[9, 10].

A rather abrupt increase of  $J/\psi$  suppression with increasing transverse energy  $(E_T)$  in Pb-Pb interactions observed by the NA50 Collaboration [4] can also be interpreted as due to the formation of an "anomalous" state of hadronic matter, possibly of the QGP. Models of this type have been suggested by Blaizot and Ollitrault (referred to below as BO) [11] and by Kharzeev, Lourenço, Nardi and Satz (KLNS) [12]. Both models have built in the onset of the formation of matter in the "anomalous" state so they are both able to describe the onset of the anomalous  $J/\psi$  suppression at a given value of  $E_T$  in Pb-Pb interactions at the CERN SPS.

The purpose of this note is to point out that BO and KLNS models are based on different assumptions on dynamics of production of matter with high energy density in nuclear collisions and if they were the only possibilities it would be possible to learn which one gives the correct description by studying the occurence of anomalous  $J/\psi$  suppression as function of nucleon numbers of colliding nuclei. Since, in fact, BO and KLNS are not the only possibilities, the data on the dependence of anomalous  $J/\psi$  suppression on nucleon numbers of colliding nuclei, if available, would provide an information on the mechanism of transverse energy production in nuclear collisions. In the Sect.2 we shall discuss this point in more detail, in Sect.3 we shall analyze the possibility of discriminating between the BO and KLNS models by using suitable experimental data and in Sect.4 we shall present conclusions and comments.

## 2 Nucleon number dependence of the onset of anomalous $J/\psi$ production in BO and KLNS models

In the present paper we shall use a simple model of nuclei as spheres with constant nucleon densities and radii of  $r_A = 1, 2A^{1/3}$  fm where A is the nucleon number. In this simplified description some results can be obtained in a transparent analytical form.

Both BO [11] and KLNS [12] assume that the mechanism of anomalous

 $J/\psi$  suppression observed by the NA50 Collaboration at the CERN SPS is due to the dissolution of  $J/\psi$  by the Quark- Gluon Plasma (QGP). The time spent by the  $J/\psi$  in the QGP does not enter into these models what means that the dissolution is assumed to be immediate. The dynamics of at least the first decisive stage of the nuclear collision is assumed to be longitudinal in the sense that the criterion for the formation of QGP depends only on the particular tube- on- tube interaction. Following the picture of the Wounded Nucleon Model [13] BO introduce density of participating nucleons per unit transverse area

$$n_p(\vec{s}, \vec{b}) = T_A(\vec{s}) [1 - exp(-\sigma_N T_B(\vec{s} - \vec{b}))] + T_B(\vec{s} - \vec{b}) [1 - exp(-\sigma_N T_A(\vec{s}))]$$
(1)

where  $\sigma_N$  is the non-diffractive nucleon- nucleon cross- section,  $\vec{b}$  is the impact parameter and  $\vec{s}$  gives the transverse position of a nucleon with respect to the center of the nucleus A. Functions  $T_A(\vec{s})$  and  $T_B(\vec{s}-\vec{b})$  give the nucleon densities per unit area in the transverse plane with

$$T_A(s) = \int_{-\infty}^{+\infty} \rho_A(s, z) dz \tag{2}$$

The condition for the formation of QGP at a particular value of  $\vec{b}$  and  $\vec{s}$  is then stated in Ref.[11] as

$$n_p(\vec{s}, \vec{b}) \ge n_{BO,c} \tag{3}$$

where  $n_{BO,c}$  is supposed to be just equal to the maximal possible value reached in S + U collisions. Note that the condition in Eq.(3) makes no explicit mention of the role possibly played by the formation times of the secondary hadrons or partons, which may be of some importance, see e.g. Ref.[14].

Neglecting exponential factors in Eq.(1), using the sharp sphere approximation and multiplying  $n_p(\vec{s}, \vec{b})$  by  $\sigma_N = 30$ mb to get a dimensionless quantity we obtain Eq.(3) in the following form

$$\kappa_{BO} \equiv \rho_0 \sigma_N 2 \sqrt{R_A^2 - s^2} + \rho_0 \sigma_N 2 \sqrt{R_B^2 - (\vec{s} - \vec{b})^2} \ge \kappa_{BO,c} \tag{4}$$

where  $\kappa_{BO}$  is defined as  $n_p$  multiplied by  $\sigma_N$  and  $\rho_0 = 0.138 f m^{-3}$ . Following their earlier work [15] BO assume in Ref.[11] that all  $J/\psi$ 's produced at values of  $\vec{b}, \vec{s}$  satisfying Eq.(4) are completely suppressed (independently of whether they are produced directly or via e.g. radiative decays of  $\chi$ ). When discussing the onset of QGP formation and thus also the onset of anomalous suppression of  $J/\psi$  as function of nucleon numbers A and B we have to consider the collision of tubes with b = 0 and s = 0, obtaining thus

$$2\rho_0 \sigma_N 1.2 (A^{1/3} + B^{1/3}) \ge \kappa_{BO,c} \tag{5}$$

which in variables  $x = A^{1/3}$ ,  $y = B^{1/3}$  gives

$$x + y = 1.006\kappa_{BO,c} \tag{6}$$

In their model of the formation of QGP KLNS [12] use also the picture of tube- on- tube collisions, but in contradistinction to BO [11] they use as a criterion the interaction density [16] defined (in the sharp sphere model) as

$$\kappa_K = \frac{N_c}{N_w} = \frac{\rho_0 \sigma_N 2 \sqrt{R_A^2 - s^2} . \rho_0 \sigma_N 2 \sqrt{R_B^2 - (\vec{s} - \vec{b})^2}}{\rho_0 \sigma_N 2 \sqrt{R_A^2 - s^2} + \rho_0 \sigma_N 2 \sqrt{R_B^2 - (\vec{s} - \vec{b})^2}}$$
(7)

The onset of  $J/\psi$  suppression at the highest available  $E_T$  (neglecting fluctuations in  $E_T$ ) corresponds again to b = 0, s = 0 and Eq.(7) in this case leads to

$$\frac{xy}{x+y} = 1.006\kappa_{K,c}$$

what is equivalent to

$$y = \frac{a^2}{x-a} + a, \qquad a = 1.006\kappa_{K,c}$$
 (8)

As pointed out by KLNS the  $\kappa_K$  as defined by the first part of Eq.(7) can in principle be determined experimentally. The number of nucleon- nucleon collision  $N_c$  is proportional to the Drell- Yan (DY) pair production, whereas the number of secondary hadrons produced is proportional to the number of participating (wounded) nucleons. At the CERN SPS this statement is only approximately correct, since the data on transverse energy  $(E_T)$  distributions and in particular the data on  $E_T$  distributions associated to DY production do require some contribution to  $E_T$  proportional to  $N_c$ , for details see the paper by Armesto, Capella, and Ferreiro [8] based on the Dual Parton Model [17]. We would like to stress that the dynamics responsible for energy density as given by Eq.(7) is rather different from that corresponding to Eq.(4). When considering the A+B interaction in the cms of individual nucleon- nucleon collisions, the numerator in Eq.(7) is proportional to the number of nucleon- nucleon collisions (within a given tube-on-tube collision) whereas the denominator is proportional to the sum of the lengths (and volumes) of the two interacting tubes. So one of possible interpretations of Eq.(7) is that  $E_T$  relevant for the QGP formation is due to semihard gluon- gluon interactions, since the repetition of soft ones would be damped by the Landau-Pomeranchuk effect. The ratio in Eq.(7) is then proportional to the total transverse energy over the interaction volume and thus to energy density.

For the sake of completeness we shall now present a few details concerning the calculations. When using the Gerschel and Hüfner [5, 6] mechanism we have taken  $\sigma_{abs} = 7$ mb and we have not introduced the effects due to the fluctuations of the  $E_T = E_T(b)$  dependence. The calculations has been performed by using the expression

$$S = \frac{N}{N_0} \tag{9}$$

where

$$N_{0} = \int_{0}^{R_{A}} sds \int_{0}^{2\pi} d\theta 2\sqrt{R_{A}^{2} - s^{2}} 2L_{B}(\vec{b}, \vec{s}, \theta)$$
$$L_{A}(s) = \sqrt{R_{A}^{2} - s^{2}}, \quad L_{B}(\vec{b}, \vec{s}, \theta) = \sqrt{R_{B}^{2} - b^{2} - s^{2} + 2bscos(\theta)}$$

when the expression under the square-root in  $L_B$  is positive, and  $L_B(\vec{b}, \vec{s}, \theta) = 0$  when this expression is negative.

The numerator is given as

$$N = \int_{0}^{R_{A}} sds \int_{0}^{2\pi} d\theta \Theta(R_{B}^{2} - b^{2} - s^{2} + 2bscos(\theta))$$
$$\int_{-L_{A}(s)}^{L_{A}(s)} dz_{A} \int_{-L_{B}(s,\theta)}^{L_{B}(s,\theta)} dz_{B}e^{-\rho_{A}\sigma_{a}[z_{A}+L_{A}(s)]}$$
$$e^{-\rho_{B}\sigma_{a}[z_{B}+L_{B}(s,\theta)]}\Theta(\kappa_{c} - \kappa)$$

where we have put  $\sigma_a = 7$ mb. The last term is to be interpreted as  $\Theta(\kappa_{c,K}^{\chi} - \kappa_K^{\chi})$  or  $\Theta(\kappa_{c,BO} - \kappa_{BO})$  respectively, or in the case of the two threshold scheme we have to make the replacement, see Ref. [12]

$$\Theta(\kappa_c - \kappa) \to 0.4\Theta(\kappa_{K,c}^{\chi} - \kappa_K) + 0.6\Theta(\kappa_{K,c}^{\psi} - \kappa_K)$$

where the first threshold corresponds to the dissolution of  $\chi$  by QGP and the second one to the dissolution of  $J/\psi$ .

### 3 On the possibility of discriminating between BO and KLNS models by data

In this section we shall show first that KLNS and BO schemes cannot be made equivalent in what concerns the dependence of anomalous  $J/\psi$  suppression on  $E_T$  and on nucleon numbers A,B of colliding nuclei. As an illustration we shall consider a situation in which the onset of anomalous  $J/\psi$  suppression is assumed to occur (without taking into account the fluctuations of  $E_T$ ) just in the interaction of the two longest tubes in S+U collision. In this academic example we take U-nucleus as spherical. In this case  $\kappa_{BO,c} = 9.312$  and  $\kappa_{K,c} = 2.086$  and the curves for the onset of the anomalous  $J/\psi$  suppression in the  $x = A^{1/3}$  and  $y = B^{1/3}$  plane are given by Eqs.(6,8) and this leads to

$$(BO): \quad y = 9.37 - x, \qquad (KLNS): \quad y = \frac{4.41}{x - 2.1} + 2.1 \qquad (10)$$

The two curves are shown in Fig.1. As expected the two curves cross each other for values of x and y corresponding to S and U nuclei.

With  $\kappa_{BO,c}=9.312$  and  $\kappa_{K,c}=2.08$  fixed by the assumption that anomalous  $J/\psi$  suppression sets on just at the end of the  $E_T$  distribution for S-U interactions we can now study the onset of anomalous  $J/\psi$  suppression in Pb-Pb collisions. In Fig.2 we present  $J/\psi$  suppression calculated via Eq.(9) in Pb-Pb interactions. We have calculated  $E_T(b)$  by using the relationship  $E_T(b) = 0.325N_w(b)$ GeV, where  $N_w(b)$  is the number of wounded (participating) nucleons at a given value of b.

Fig.2 demonstrates an interesting, although very simple effect. When  $\kappa_{BO,c}$  and  $\kappa_{K,c}$  are fixed by the condition related to the interaction of the two longest tubes in S-U interaction, the onset of the anomalous  $J/\psi$  suppression in Pb-Pb collisions appear at different values of  $E_T$  in BO and KLNS schemes.

Since, in fact, the anomalous  $J/\psi$  suppression has not been observed in S-U interactions we find it more appropriate to determine the values  $\kappa_{BO,c}$  and  $\kappa_{K,c}$  by the data [4] on  $J/\psi$  suppression in Pb-Pb collisions.

In Fig.3 we present the NA50 Collaboration data [4] on the survival probability of  $J/\psi$  as a function of the transverse energy  $E_T$  in Pb-Pb interactions at 158 GeV per nucleon and three curves giving  $J/\psi$  suppression under different conditions: the curve denoted as "no anom." corresponds to pure Gerschel- Hüfner mechanism with  $\sigma_{abs} = 7$ mb, the curve "BO" is given by the BO scheme with  $\kappa_{BO,c} = 9,75$  and with all  $J/\psi$ 's with  $\kappa_{BO} \geq \kappa_{BO,c}$  completely absorbed, the curve "KLNS" corresponds to the KLNS scheme with  $\kappa_{K,c} = 2,43$  and with all  $J/\psi$ 's with  $\kappa_K \geq \kappa_{K,c}$  completely absorbed.

In their work KLNS [12] take into account that about 40% of  $J/\psi$ 's observed in nuclear collisions is due to radiative  $\chi$ - decays and assume that  $\chi$  is suppressed at a different threshold in  $\kappa_K$  than directly produced  $J/\psi$ . This permits them to describe also the second decrease of survival probability of  $J/\psi$  in the region of  $E_T > 100$ GeV indicated by the data.

The same two threshold mechanism could also be implemented into the BO scheme and in this way both BO and KLNS schemes could describe the Pb+Pb data in a similar way. Within our simple model this could be achieved by increasing the value of  $\sigma_{abs}$  and lowering the tresholds given by  $\kappa_{K,c}$  and  $\kappa_{BO,c}$ . We shall not follow this path here since the purpose of this note is to study only the onset (the first one) of the anomalous  $J/\psi$  suppression and not the detailed shape of the survival probabability, including the possibility of a second threshold (the possible onset of  $\chi$  suppression).

As can be seen in Fig.3 shapes of  $S(E_T)$  obtained within BO and KLNS schemes in Pb-Pb interactions are rather similar when  $\kappa_{BO,c}$  and  $\kappa_{K,c}$  are fixed by the condition that the onset of anomalous suppression starts at the same value of  $E_T$ . It seems therefore that by using only the Pb-Pb data it would be very difficult to discriminate between the two schemes.

In the rest of this section we shall try to find out whether one can discriminate between the BO and KLNS by studying the onset of anomalous  $J/\psi$  suppression as the function of atomic numbers A and B of the colliding nuclei. Note that we are neglecting the role of possible fluctuations in the energy density of matter formed in nuclear collisions, see Ref.[18].

In Fig.4 we show curves giving the onset of anomalous  $J/\psi$  suppression in the  $x = A^{1/3}$  and  $y = B^{1/3}$  plane corresponding to values of  $\kappa_{K,c} = 2.43$ and  $\kappa_{BO,c} = 9.75$  (the values used also in calculations leading to Fig.3).

As can be seen in Fig.4 for A+Pb collisions with  $57 \le A \le 73$  the survival probability as function of  $E_T$  shows the presence of anomalous suppression within the BO scheme and no anomaly for the KLNS one. For nuclei with  $74 \le A$  both KLNS and BO show the presence of anomaly but the anomaly sets on at different values of  $E_T$ . As an illustration we show in Fig.5 the results obtained again via Eq.(9) with  $\kappa_{BO,c} = 9.75$  and  $\kappa_{K,c} = 2.43$  for the  $As^{75}$ -Pb interactions, still using  $E_T(b) = 0.325 \text{GeV} N_w(b)$ .

#### 4 Comments and conclusions

As can be seen in Fig.4 there do exist regions in the A,B plane where the BO picture of  $J/\psi$  suppression based on the Wounded Nucleon Model [13] predicts anomalous  $J/\psi$  suppression whereas the KLNS scheme [12] predicts its absence. For Pb as one of colliding nuclei this region for the other nucleus A contains nuclei with

$$57 \le A \le 73 \tag{11}$$

including nuclei as  $Ni_{59}$ ,  $Cu_{64}$  and  $Zn_{65}$ . For A  $\geq$  74 the anomalous suppression is present in both schemes but sets on at lower value of  $E_T$  in the BO case.

To conclude: we propose to study experimentally anomalous  $J/\psi$  suppression at the CERN SPS in collisions of A-Pb or Pb-A with the nucleon number A within the interval given by Eq.(11). The presence of anomalous  $J/\psi$  suppression for A-Pb collisions with A within the interval given by Eq.(11) would support the scheme of high energy density formation suggested by Blaizot and Ollitrault [11], the absence of it would give evidence in favour of the Kharzeev, Lourenço, Nardi and Satz model [12].

A study of the onset of anomalous  $J/\psi$  suppression in a series of A-Pb interactions including the available data on S-Pb and Pb-Pb, and some of A-Pb with A given by Eq.(11) (or with A only a bit higher) at 158Gev per nucleon could bring interesting evidence on the dynamics of production of critical energy density for QGP formation in this energy range. Moreover, since at present the anomalous  $J/\psi$  suppression has been observed in only one case, the second observation in A-Pb with a lighter A would support the idea of anomalous  $J/\psi$  suppression by an anomalous state of matter, possibly by QGP. As seen in Fig.5 the difference between predictions of BO and KLNS schemes is not very large and some indeterminacy will be introduced by  $E_T$  fluctuations [12, 18, 19] not considered here we still believe that a study of A-Pb interactions with A given by Eq.(11) or nearby could bring in a very useful information.

Acknowledgements: the authors are indebted to A.Capella and R.Hwa for useful discussions on the problem of dynamics of nuclear collisions and of anomalous  $J/\psi$  suppression. The present work has been supported in part by the U.S.- Slovakia Science and Technology Program, under Grant No. 003-95 (INT 9319091) and by the Slovak Grant Agency under Grant V2F13-G.

#### **Figure Captions**

**Fig.1** Curves denoted as KLNS and BO give lines of the onset of anomalous  $J/\psi$  suppression in BO [11] and KLNS [12] schemes as functions of nucleon numbers A and B. The curves are given explicitly in Eq.(10). The critical values are  $\kappa_{BO,c}=9.312$  and  $\kappa_{K,c}=2.086$ .

**Fig.2** Dependence of the survival probability  $S(E_T)$  on transverse energy  $E_T$  calculated via Eq.(9) for KLNS and BO schemes with  $\kappa_{BO,c}$ =9.312 and  $\kappa_{K,c}$ =2.086 (corresponding to the onset of anomalous suppression just above the S-U interactions) in the space of nucleon numbers A and B. The curve denoted as "no anom." gives the case with no anomalous suppression.

**Fig.3** Comparison of survival probabilities of  $J/\psi$  in Pb-Pb collisions in cases of no anomalous suppression ("no anom.") and BO and KLNS schemes. More details are given in the text. The data are taken from Ref.[4] and w.m.b means "with minimum bias". The critical values of  $\kappa$ 's are  $\kappa_{BO,c}=9.75$  and  $\kappa_{K,c}=2.43$ 

**Fig.4** Curves giving the onset of anomalous  $J/\psi$  suppression within KLNS and BO schemes with values  $\kappa_{BO,c}=9.75$  and  $\kappa_{K,c}=2.43$  obtained in comparison of KLNS and BO schemes with data in Fig.3.

**Fig.5** Survival probability  $S(E_T)$  calculated with  $\kappa_{BO,c}$ =9.75 and  $\kappa_{K,c}$ =2.43 for As<sup>75</sup>-Pb interactions for BO and KLNS schemes.

### References

- M.C.Abreu et al., NA50 Collab., Phys.Lett. B410 (1997) 337; M.Gonin et al., NA50 Collab., Nucl.Phys. A610 (1996) 404c
- [2] C.Lourenco, NA50 Collab., Nucl. Phys. A610 (1996) 552c
- [3] M.C.Abreu et al., NA50 Collab., Nucl. Phys. A638 (1998) 261c
- [4] M.C.Abreu et al., NA50 Collab., Phys. Lett. **B477** (2000) 28

- [5] C.Gerschel, J.Hüfner, Zeit.f.Phys. C56 (1992) 171
- [6] C.Gerschel, J.Hüfner, Phys.Rev. **B207** (1988) 253
- [7] R.Vogt, Phys. Reports **310** (1999) 197
- [8] N.Armesto, A.Capella and E.G.Ferreiro, Phys. Rev. C59 (1999) 395
- [9] R.C.Hwa, J.Pišút, N.Pišútová, Phys.Rev. C58 (1998) 434
- [10] C.Gale, S.Jeon, and J.Kapusta, Phys. Rev. Lett. 82 (1999) 1639;
  E.Marco, and E.Oset, Nucl. Phys. A 645 (1999) 303
- [11] J.-P.Blaizot and J.-Y.Ollitrault, Phys. Rev. Lett. 77 (1996) 1703
- [12] D.Kharzeev, C.Lourenço, M.Nardi and H.Satz, Zeit.f.Phys. C74 (1997) 307
- [13] A.Bialas, M.Bleszynski and W.Czyz, Nucl. Phys. B111 (1976) 461; for review see A.Bialas, in Proceedings of the XIIIth Int. Symposium on Multiparticle Dynamics, edited by W.Kittel, W.Metzger and A.Stergion, World Scientific, Singapore, 1983, p.328
- [14] J.Pišút, N.Pišútová and P.Závada, Zeit. f. Phys. C67 (1995) 467
- [15] J.-P.Blaizot and J.-Y.Ollitrault, Phys.Rev. **D39** (1989) 232
- [16] D.Kharzeev, Nucl. Phys. A610 (1996) 418c
- [17] A.Capella, V.P.Sukhatme, C.-I.Tan, and J.Tran Than Van, Physics Reports, 236 (1994) 225
- [18] J.-P.Blaizot, P.M.Dinh, and J.-Y.Ollitrault, Transverse energy fluctuations and the pattern of  $J/\psi$  suppression in Pb-Pb collisions, nuclth/0007020
- [19] A.Capella, E.G.Ferreiro, A.B.Kaidalov, Phys. Rev. Lett. 85 (2000) 2080









