

THEORY OF THE NUCLEON SPIN-POLARIZABILITIES II

THOMAS R. HEMMERT

*IKP (Theorie), FZ Jülich, D-52425 Jülich, Germany**

Email: th.hemmert@fz-juelich.de

I discuss a generalization of the concept of nucleon spin-polarizabilities and present an overview of the available theoretical predictions up to now for these elusive but interesting structure parameters of the nucleon.

The concept of nucleon spin-polarizabilities has been discussed in full detail in the plenary talk contribution by Ulf Meißner ¹. In this brief contribution I want to point out that an interpretation of the 4 so called Ragusa spin-polarizabilities γ_i ² in terms of multipole excitations and de-excitations can be given, if one resorts to a different basis R_i for the excitation part of the 6 structure functions A_i of nucleon Compton scattering discussed in Meißner's talk. For example ³, choosing the following linear combinations of the γ_i

$$\gamma_{E1} = -\gamma_1 - \gamma_3 \quad (E1 \rightarrow E1) \quad (1)$$

$$\gamma_{M1} = \gamma_4 \quad (M1 \rightarrow M1) \quad (2)$$

$$\gamma_{E2} = \gamma_2 + \gamma_4 \quad (M1 \rightarrow E2) \quad (3)$$

$$\gamma_{M2} = \gamma_3 \quad (E1 \rightarrow M2), \quad (4)$$

the resulting spin-polarizabilities $\gamma_{E1(M1)}$ can be interpreted as the nucleon's spin response to an electric (magnetic) dipole excitation via the incoming Compton photon and a successive dipole de-excitation through the outgoing Compton photon. Likewise, the other 2 linear combinations $\gamma_{E2(M2)}$ describe the response to an electric (magnetic) dipole excitation and a successive magnetic (electric) quadrupole de-excitation. The 4 Ragusa spin-polarizabilities can therefore be expressed in a basis where their physical meaning becomes clear—these polarizabilities are just parameterizing the leading spin response of the nucleon for dipole excitations in (spin-dependent) Compton scattering. As discussed by Meißner ¹, none of these (dipole) spin-polarizabilities has been measured directly up to now—results of future polarized Compton scattering experiments on the proton are eagerly awaited. In chiral perturbation theory the dipole spin-polarizabilities have now been calculated to next-to-leading order, the results and a comparison with recent dispersion theoretical calculations are given in table 1.

*New address: Physik Department T39, TU München, D-85747 Garching, Germany.

Table 1. Predictions for the isoscalar and isovector dipole spin-polarizabilities of the nucleon found to $\mathcal{O}(p^4)$ in heavy baryon chiral perturbation theory (GHM), to $\mathcal{O}(\epsilon^3)$ in the small scale expansion (SSE1) and in dispersion analyses (DKH,DGPV,BGLM). All results are given in the units 10^{-4} fm^4 . (*: $+2.5 \times 10^{-4} \text{ fm}^4$ from $\Delta(1232)$ pole still missing)

$\gamma_i^{(N)}$	GHM ⁵	DKH ⁶	DGPV ⁷	BGLMN ³	SSE1 ⁸
$\gamma_{E1}^{(s)}$	-3.0	-5.0	-5.2	-4.5	-5.2
$\gamma_{M1}^{(s)}$	+0.4*	+3.4	+3.4	+3.3	+1.4
$\gamma_{E2}^{(s)}$	+2.0	+2.4	+2.7	+2.4	+1.0
$\gamma_{M2}^{(s)}$	+0.6	-0.6	-0.5	-0.2	+1.0
$\gamma_{E1}^{(v)}$	+1.2	+0.5	+0.8	+1.1	-
$\gamma_{M1}^{(v)}$	+0.0	+0.0	-0.5	-0.6	-
$\gamma_{E2}^{(v)}$	-0.2	-0.2	-0.5	-0.5	-
$\gamma_{M2}^{(v)}$	+0.1	-0.0	+0.5	+0.5	-

After realizing the possibility of such a multipole interpretation for the Ragusa spin-polarizabilities, one of course does not have to stop classifying the nucleon's spin-dependent response in external electromagnetic fields at the level of electric or magnetic dipole excitations. Recently, Holstein et al.⁴ have generalized such an analysis to classify spin structure responses related to low energy quadrupole excitations of the nucleon. A straightforward derivation of these higher order spin-polarizabilities can be given via a multipole expansion of the 6 Compton structure amplitudes $R_i(\omega, \theta)$, $i = 1 \dots 6$, where ω corresponds to the photon energy in the center-of-mass system and θ denotes the cms scattering angle. For example, for $i = 6$ one finds³:

$$\begin{aligned}
R_6 = \sum_{l \geq 1} \{ & [f_{MM}^{l+} - f_{MM}^{l-}] (l P_l'' + P_{l-1}''') - [f_{EE}^{l+} - f_{EE}^{l-}] P_l''' \\
& - [f_{EE}^{l+} - f_{EE}^{l-}] P_l''' + f_{ME}^{l+} [(3l+1) P_l'' + 2 P_{l-1}'''] \\
& - 2 f_{EM}^{l+} [(l+1) P_{l+1}'' + 2 P_l'''] \}
\end{aligned}$$

where the $P_i^a(z)$ denote the a -th derivative of Legendre-Polynomial P_i and $f_{XY}^l(\omega)$ are the energy-dependent expansion parameters of the multipole series. If one now maps the one-particle-irreducible or excitation part of R_6 onto the corresponding A_6 structure amplitude discussed by Meißner¹ and stops the multipole series at $l = 1$, one obtains

$$\begin{aligned}
A_6^{1PI} &= -c R_6^{1PI} = +c 12 f_{EM}^{1+} + (l \geq 2) \\
&= 4 \pi \gamma_3 \omega^3 + \dots,
\end{aligned} \tag{5}$$

where c corresponds to an amplitude normalization constant and γ_3 denotes one of the spin-polarizabilities discussed by Meißner. We can therefore make the identification

$$\rightarrow \gamma_3 = \frac{3c f_{EM}^{1+}(\omega)}{\omega^3 \pi} \Big|_{\omega \rightarrow 0} \equiv \gamma_{M2} \quad (E1 \rightarrow M2),$$

which reproduces Eq.(4). Generalizing this analysis to $l = 2$ one obtains a set of 12 spin-polarizabilities defined via the expansion parameters $f_{XY}^l(\omega)$ introduced above:

$$\begin{aligned} 4\pi\omega^3 (\gamma_{E1} + \omega^2 \gamma_{E1,\nu}) &= f_{EE}^{1+}(\omega) - f_{EE}^{1-}(\omega) & (E1 \rightarrow E1) \\ 4\pi\omega^3 (\gamma_{M1} + \omega^2 \gamma_{M1,\nu}) &= f_{MM}^{1+}(\omega) - f_{MM}^{1-}(\omega) & (M1 \rightarrow M1) \\ 4\pi\omega^3 (\gamma_{E2} + \omega^2 \gamma_{E2,\nu}) &= 6 f_{ME}^{1+}(\omega) & (M1 \rightarrow E2) \\ 4\pi\omega^3 (\gamma_{M2} + \omega^2 \gamma_{M2,\nu}) &= 6 f_{EM}^{1+}(\omega) & (E1 \rightarrow M2) \\ 4\pi\omega^5 \gamma_{ET} &= 3 [f_{EE}^{2+}(\omega) - f_{EE}^{2-}(\omega)] & (E2 \rightarrow E2) \\ 4\pi\omega^5 \gamma_{MT} &= 3 [f_{MM}^{2+}(\omega) - f_{MM}^{2-}(\omega)] & (M2 \rightarrow M2) \\ 4\pi\omega^5 \gamma_{E3} &= 15 f_{ME}^{2+}(\omega) & (M2 \rightarrow E3) \\ 4\pi\omega^5 \gamma_{M3} &= 15 f_{EM}^{2+}(\omega) & (E2 \rightarrow M3). \end{aligned}$$

The 4 dipole spin-polarizabilities $\gamma_{E1}, \gamma_{M1}, \gamma_{E2}, \gamma_{M2}$ have already been discussed in Eqs.(1-4). When extending the multipole series to $l = 2$, one in addition also picks up the 4 slopes $\gamma_{E1,\nu}, \gamma_{M1,\nu}, \gamma_{E2,\nu}, \gamma_{M2,\nu}$ of their energy-dependence, which are usually called dispersion spin-polarizabilities. Finally, there are 4 proper higher order spin-polarizabilities $\gamma_{ET}, \gamma_{MT}, \gamma_{E3}, \gamma_{M3}$, which correspond to quadrupole excitations and successive quadrupole order even octupole de-excitations. Results of recent predictions for these essentially unknown higher order spin structure parameters of the nucleon obtained in chiral effective field theories and via dispersion analyses are given in table 2. It is interesting to note that all predictions agree that the dispersive effects of the dipole spin-polarizabilities should give the dominant effect among the $l = 2$ spin-polarizabilities—quadrupole and octupole de-excitation only seems to play a minor role. This is another manifestation of the rather rigid behavior of the nucleon under deformation forces as in the case of the small quadrupole $N\Delta$ -transition moment or the rather small electric (magnetic) polarizabilities $\bar{\alpha}_E, (\bar{\beta}_M)$. Furthermore, the next-to-leading order chiral perturbation theory calculation gives the first indication that as in the case of the dipole spin-polarizabilities (table 1) also for the higher order spin-polarizabilities the isoscalar components clearly dominate over the isovector part (table 2), making proton and neutron rather similar objects as far as their spin-structure at

Table 2. Status of the theoretical predictions as of summer 2000 for the higher order isoscalar and isovector spin-polarizabilities as given by next-to-leading order chiral perturbation theory, by dispersion analyses and to leading order in the small scale expansion. All results are given in the units 10^{-4} fm^6 . Note that the chiral next-to-leading calculations for the higher order isovector spin-polarizabilities find $\gamma_{ET}^{(v)} = \gamma_{MT}^{(v)} = \gamma_{E3}^{(v)} = \gamma_{M3}^{(v)} = 0$, while no numbers from dispersion theory are available yet. (*: $+1.20 \times 10^{-4} \text{ fm}^6$ from $\Delta(1232)$ pole still missing)

$\gamma_i^{(N)}$	$\mathcal{O}(p^4)$ ChPT ⁹	Disp. Th. ⁴	$\mathcal{O}(\epsilon^3)$ SSE ⁴
$\gamma_{ET}^{(s)}$	+0.08	-0.15	-0.28
$\gamma_{MT}^{(s)}$	+0.06	-0.09	-0.03
$\gamma_{E3}^{(s)}$	+0.03	+0.06	+0.11
$\gamma_{M3}^{(s)}$	+0.03	+0.09	+0.11
$\gamma_{E1,\nu}^{(s)}$	-3.40	-3.42	-5.16
$\gamma_{M1,\nu}^{(s)}$	+0.63*	+2.23	+0.83
$\gamma_{E2,\nu}^{(s)}$	+1.68	+1.30	+0.28
$\gamma_{M2,\nu}^{(s)}$	-0.12	-0.60	-0.22
$\gamma_{E1,\nu}^{(v)}$	+0.97	?	-
$\gamma_{M1,\nu}^{(v)}$	+0.00	?	-
$\gamma_{E2,\nu}^{(v)}$	-0.09	?	-
$\gamma_{M2,\nu}^{(v)}$	+0.05	?	-

low energies is concerned.

I would like to thank the organizer of GDH2000 for the opportunity to present these new results and I am grateful for their financial support.

References

1. U.-G. Meißner, “Theory of the Nucleon Spin-Polarizabilities I”, these proceedings, [hep-ph/0008060].
2. S. Ragusa, Phys. Rev. **D47** (1993) 3757.
3. D. Babusci et al., Phys Rev. **C58** (1998) 1013.
4. Holstein et. al, Phys. Rev. **C61** (2000) 034316.
5. G.C. Gellas, T.R. Hemmert and U.-G. Meißner, Phys. Rev. Lett. **85** (2000) 14.
6. D. Drechsel, O. Krein, O. Hanstein, Phys.Lett. **B420** (1998) 248.
7. D. Drechsel et al., Phys. Rev. **C61** (2000) 015204.
8. T.R. Hemmert et al., Phys. Rev. **D57** (1998) 5746.
9. G.C. Gellas et al., “Multipole Analysis of the Nucleon’s Polarizabilities,

forthcoming.