## Photon production in relativistic heavy-ion collisions using rates with two-loop calculations from quark matter

Dinesh Kumar Srivastava

Variable Energy Cyclotron Centre, 1/AF Bidhan Nagar, Calcutta 700 064, India

Erratum: Eur. Phy. J. C 10 (1999) 487.

It has been shown recently [1] shown that the values of  $J_T$  and  $J_L$  which appear in Eq.(2) and (5) of the above paper and which are taken from the work of Aurenche et al[2] are too large by a factor of 4. Correcting for this changes the Fig. 1–4 (see below). The basic result of the paper, that the emissions from the quark matter can outshine those from the hadronic matter when the photon rates upto two-loop level are used, remains valid though the range of  $p_T$  over which it happens is reduced to  $p_T < 0.5$ , 1, and 2 GeV/c respectively at SPS, RHIC, and LHC energies.

## References

- D. Dutta et al. and P. Aurenche et al. Private Communication, F. D. Steffen and M. H. Thoma, hep-ph/0103044.
- [2] P. Aurenche, F. Gelis, H. Zaraket, and R. Kobes, Phys. Rev. D. 58, 085003 (1998).




Figure 1: Radiation of photons from various processes in the quark matter at  $T=250~\mathrm{MeV}$ 




Figure 2: Radiation of photons from central collision of lead nuclei at SPS energies from the hadronic matter (in the mixed phase and the hadronic phase) and the quark matter (in the QGP phase and the mixed phase). The contribution of the quark matter while using the rates obtained by Kapusta et al and Aurenche et al, and those from hard QCD processes are shown separately




Figure 3: Same as Fig. 2 for RHIC energies.




Figure 4: Same as Fig. 2 for LHC energies.