

A simple functional form for proton- ^{208}Pb total reaction cross sections

S. Majumdar, P. K. Deb, and K. Amos

School of Physics, University of Melbourne, Parkville, Victoria, Australia, 3010

(November 21, 2018)

Abstract

A simple functional form has been found that gives a good representation of the total reaction cross sections for the scattering from ^{208}Pb of protons with energies in the range 30 to 300 MeV.

arXiv:nucl-th/0103066v1 26 Mar 2001

The values of total reaction cross sections of nucleon scattering from nuclei (stable as well as radioactive) are required in a number of fields of study. Some of quite current interest concern the transmutation of long-lived radioactive waste into shorter-lived products using accelerator-driven systems (ADS) and in predicting dosimetries for patients in radiation therapy. To be able to specify those total reaction cross sections in a simple functional form then has great utility for any associated evaluation such as of nucleon production in spallation calculations.

Recently [1] nucleon-nucleus total reaction cross sections to 300 MeV for ^{12}C and ^{208}Pb have been predicted in good agreement with data. Those cross sections were evaluated using g -folding optical potentials formed by full folding realistic two nucleon (NN) interactions with credible nucleon-based structure models of those nuclei. The effective interactions at each energy in the range to 300 MeV were defined from the NN g matrices (solutions of the Bruckner-Bethe-Goldstone equations) of the free NN (BonnB) interactions; and so vary with energy and the medium. The ground state of ^{12}C was specified from a complete $(0 + 2)\hbar\omega$ shell model evaluation [1], while that of ^{208}Pb was obtained from a Skyrme-Hartree-Fock calculation [2]. The resulting nA optical potentials are complex, energy dependent and very non-local. All details of the approach and numerous demonstrations of its successful use with targets spanning the entire mass range (3 to 238) are given in the recent review [3]. But each such prediction of nucleon scattering is an involved calculation that culminates with use of large scale computer programs, and those of DWBA98 [3,4] in our recent studies. It would be very utilitarian if aspects of such scattering were indeed well approximated by a simple convenient functional form. We demonstrate herein that for the total reaction cross sections such a form may exist at least for heavy nuclei such as ^{208}Pb .

We have used the g -folding optical potential calculations to specify proton- ^{208}Pb scattering S matrices at energies $E \propto k^2$,

$$S_l^\pm \equiv S_l^\pm(k) = e^{2i\delta_l^\pm(k)} = \eta_l^\pm(k) e^{2i\Re[\delta_l^\pm(k)]}, \quad (1)$$

where $\delta_l^\pm(k)$ are the (complex) scattering phase shifts and $\eta_l^\pm(k)$ are the moduli of the S matrices. The superscript designates $j = l \pm 1/2$. Total reaction cross sections then are predicted by

$$\begin{aligned} \sigma_R(E) &= \frac{\pi}{k^2} \sum_{l=0}^{\infty} \left\{ (l+1) \left[1 - (\eta_l^+)^2 \right] + l \left[1 - (\eta_l^-)^2 \right] \right\} \\ &= \frac{\pi}{k^2} \sum_{l=0}^{\infty} \sigma_l^{(R)}(E). \end{aligned} \quad (2)$$

The partial reaction cross sections $\sigma_l^{(R)}(E)$ so found we now take as 'data'. The values from calculations of the scattering of protons from ^{208}Pb are displayed in the Fig. 1 by the filled circles for energies 30, 40, 50, 65, 100, 160, 200, 250, and 300 MeV as the peak magnitudes increase respectively. They suggest to us that the partial total reaction cross sections $\sigma_l^{(R)}(E)$ can be described by the simple functional form,

$$\sigma_l^{(R)}(E) = (2l+1) \left[1 + e^{\frac{(l-l_0)}{a}} \right]^{-1} + \epsilon(2l_0+1) e^{\frac{(l-l_0)}{a}} \left[1 + e^{\frac{(l-l_0)}{a}} \right]^{-2} \quad (3)$$

with $l_0(E, A)$, $a(E, A)$, and $\epsilon(E, A)$ varying smoothly with energy and mass.

The summation giving the total reaction cross section can be limited to a value l_{max} and the associated form tends appropriately to the known high energy limit. With increasing energy, l_{max} becomes so large that the exponential fall-off of the functional form, Eq. 3, can be approximated as a straight vertical line ($l_0 = l_{max}$). In that case, the total reaction cross section equates to the area of a triangle, and

$$\sigma_R \Rightarrow \frac{\pi}{2k^2} l_{max} (2l_{max} + 1) \approx \frac{\pi}{k^2} l_{max}^2 . \quad (4)$$

Then with $l_{max} \sim kR$, at high energies

$$\sigma_R \Rightarrow \pi R^2 ; \quad (5)$$

the geometric cross section as required.

The fits of this functional form to the "data" from ^{208}Pb are displayed by the solid curves in Fig. 1 with the values of the three parameters given at the specific energies connected by straight lines in Fig. 2. These values are plotted as functions of \sqrt{E} (effectively k) and it is clear that they are smooth variations with energy. Of these the linearity of l_0 with \sqrt{E} is most evident and we find

$$l_0(E, ^{208}\text{Pb}) \sim kR - 3 ; \quad R \sim 1.6A^{-\frac{1}{3}} . \quad (6)$$

The best fit values of these parameters are listed in Table I.

For any dynamical situation in which numerous partial reaction cross sections contribute non-negligibly to the sum in Eq. 2, one can use the limit form

$$\sigma_R(E) = \frac{\pi}{k^2} \sum_{l=0}^{\infty} \sigma_l^{(R)}(E) \Rightarrow \frac{\pi}{k^2} \int_0^{\infty} \sigma^{(R)}(E; \lambda) d\lambda . \quad (7)$$

Then with $x = \frac{\lambda - l_0}{a}$ and using the basic integrals

$$\begin{aligned} \int_{-l_0/a}^{\infty} \frac{1}{1 + e^x} dx &= \left[\int_0^{l_0/a} \frac{1}{1 + e^{-x}} dx + \int_0^{\infty} \frac{e^{-x}}{1 + e^{-x}} dx \right] = \frac{l_0}{a} + \sum_{n=1}^{\infty} (-)^{n+1} e^{-\frac{l_0}{a}n} \\ \int_{-l_0/a}^{\infty} \frac{x}{1 + e^x} dx &= \left[- \int_0^{l_0/a} \frac{x}{1 + e^{-x}} dx + \int_0^{\infty} \frac{x e^{-x}}{1 + e^{-x}} dx \right] \\ &= -\frac{1}{2} \frac{l_0^2}{a^2} + 1.645 + \sum_{n=1}^{\infty} (-)^n \left(1 + \frac{1}{n^2} \right) e^{-\frac{l_0}{a}} , \end{aligned} \quad (8)$$

the integral observable is

$$\begin{aligned} \sigma^{(R)}(E) &= \frac{\pi}{k^2} \sum_l \sigma_l^{(R)}(E) \rightarrow \frac{\pi}{k^2} \int_0^{\infty} \sigma^{(R)}(\lambda; E) d\lambda \\ &= \frac{\pi}{k^2} \left\{ l_0(l_0 + 1) + 3.29a^2 + a\epsilon(2l_0 + 1) \right. \\ &\quad \left. + \sum_{n=0}^{\infty} e^{-n\frac{l_0}{a}} (-)^n \left[2a^2 \left\{ 1 + \frac{1}{n^2} \right\} + (2l_0 + 1) \left\{ a\epsilon - \frac{1}{n} \right\} \right] \right\} \\ &\rightarrow \frac{\pi}{k^2} \left\{ l_0(l_0 + 1) + 3.29a^2 + a\epsilon(2l_0 + 1) + \left[(2l_0 + 1) \left\{ \frac{1}{n} - a\epsilon \right\} - 4a^2 \right] e^{-\frac{l_0}{a}} \right\} \quad (9) \end{aligned}$$

Thus if l_0 , a , and ϵ are well behaved functions from matching to observed data, one can have confidence that these simple prescriptions give reliable estimates of reaction cross section values that have not been measured as yet but may be required as input in, for example, spallation production calculations. The ratios of the total reaction cross sections calculated under this approximation to those determined from the microscopic optical model potentials are listed in the last column of the Table. The results compare well to within a few percent.

TABLES

TABLE I. Parameter values giving fits to $\sigma_l^{(R)}(E)$

E (MeV)	l_0	a	ϵ	$\sigma_{n=1}^{(R)}/\sigma^{(R)}(E)$
30	10.2	1.123	-1.19	0.99
40	11.9	1.02	-1.13	1.02
50	13.1	0.89	-0.81	1.01
65	14.6	0.96	-0.63	1.00
100	18.6	1.59	-0.92	0.99
160	24.4	2.65	-1.25	0.97
200	27.5	3.06	-1.35	0.96
250	30.8	3.41	-1.48	0.98
300	34.0	3.75	-1.67	0.97

FIGURES

FIG. 1. The total partial reaction cross sections for proton scattering from ^{208}Pb .

FIG. 2. The parameters of the simple functional form for the p- ^{208}Pb partial total cross sections.

REFERENCES

- [1] P. K. Deb, K. Amos, S. Karataglidis, M. B. Chadwick, and D. G. Madland, Phys. Rev. Lett. **86**, issue 15 (2001).
- [2] B. Alex Brown, Phys. Rev. Lett. **85**, 5296 (2000).
- [3] K. Amos, P. J. Dortmans, H. V. von Geramb, S. Karataglidis, and J. Raynal, Adv. in Nucl. Phys. **25**, 275 (2000).
- [4] J. Raynal, *computer code DWBA98* NEA 1209/05, 1999.



