

# Pion Deuteron Scattering Length in Effective Field Theory

Buğra Borasoy<sup>1</sup> and Harald W. Griesshammer<sup>2</sup>

*Institut für Theoretische Physik (T39), Physik-Department,  
Technische Universität München, D-85747 Garching, Germany*

## Abstract

The S wave pion deuteron scattering length is presented in an effective field theory approach of the two-nucleon system. We include pions as dynamical particles which allows us to calculate pion re-scattering contributions inside the deuteron. Two-nucleon two-pion contact interactions with unknown parameters not determined by chiral symmetry need to be introduced in order to renormalise the appearing divergences. By choosing their values accordingly we are able to accommodate the available experimental data.

Suggested PACS numbers: 13.75.G, 14.20.Dh, 21.30.Fe, 25.80.Dj, 27.10.+h

Suggested Keywords: Effective Field Theory, pion nucleon and pion deuteron scattering length, iso-spin even pion nucleon scattering

---

<sup>1</sup>Email: borasoy@physik.tu-muenchen.de

<sup>2</sup>Email: hgrie@physik.tu-muenchen.de

# 1 Motivation

A systematic framework to calculate low-energy scattering processes of hadrons is provided by Chiral Perturbation Theory ( $\chi$ PT), the effective field theory of QCD at low energies. For example, the determination of the pion nucleon scattering length has drawn a lot of attention and still remains a topic of interest. The chiral corrections to the Weinberg-Tomozawa current algebra theorem according to which the iso-scalar (iso-spin even) S wave pion nucleon scattering length  $a^+ := \frac{1}{2}(a_{\pi p} + a_{\pi n})$  is zero at leading order (LO) have been calculated by Bernard, Kaiser and Meißner [1, 2]. Their next-to-leading order (NLO) prediction has recently been refined by determining the unknown parameters in a comparison of non-zero energy  $\pi N$  scattering at the chiral orders  $Q^3$  [3] and  $Q^4$  [4] to Koch's Karlsruhe [5], Matsinos' EM98 [6] and the VPI/GW group's SP98 [7] partial wave analyses. The obtained range  $a_{\text{HB}\chi\text{PT}}^+ = [-0.01 \dots +0.006] m_\pi^{-1}$  is still compatible with zero. Recently, Gasser et al. [8] pointed out that electro-magnetic iso-spin breaking effects could provide corrections of up to 7% to the LO result. Because  $a^+$  is very small, such effects are stronger than the naïve expectation that they are naturally suppressed by powers of the fine structure constant  $\alpha = 1/137$ , contributing at most 1%. Another recent HB $\chi$ PT analysis of pion nucleon scattering to order  $Q^4$  attempting to connect threshold and near-threshold parameters with the low energy theorems of chiral symmetry found on the basis of [5] that  $a^+ = -0.008 m_\pi^{-1}$  [9]. Direct extractions of  $a^+$  from the same  $\pi N$  partial wave analyses yield, on the other hand,  $(+0.0041 \pm 0.0009) m_\pi^{-1}$  [6] and  $+0.002 m_\pi^{-1}$  [7].

On the experimental side, the best extraction of  $a^+$  comes from measurements of elastic scattering and single charge exchange in pionic atoms. In contradistinction to phase shift analyses, no extrapolation to zero energy scattering is necessary, and electro-magnetic effects are considered small. Most recently, the ETHZ-Neuchâtel-PSI [10] collaboration found  $a_{\text{exp}, \pi N}^+ = (-0.0022 \pm 0.0043) m_\pi^{-1}$  directly from the line shift and width change in pionic hydrogen, assuming iso-spin symmetry. This is compatible with zero and with the HB $\chi$ PT result, but with smaller error bars. The ongoing experiment R-98.01 at the Paul Scherer Institute aims to reduce the error by an order of magnitude [11].

The value of the pion deuteron scattering length, which was measured by the same method as for pionic hydrogen to be  $a_{\pi d} = [(-0.0261 \pm 0.0005) + i(0.0063 \pm 0.0007)] m_{\pi}^{-1}$  [12], has also been used to constrain  $a^+$ . However, since – as mentioned above – the deuteron is not a purely iso-scalar nucleon target, binding and especially pion re-scattering effects between the two nucleons have to be accounted for. Conventional potential model approaches which combine deuteron and hydrogen data have been utilised: Baru and Kudryatsev, e.g., used multiple scattering methods and quote a value  $a_{\text{phen, BK}}^+ = (-0.0015 \pm 0.0009) m_{\pi}^{-1}$  [13], however, their errors are substantially underestimated as shown in [14]. An analysis by Landau and Thomas [15] gives  $a_{\text{phen, LT}}^+ = (+0.0016 \pm 0.0013) m_{\pi}^{-1}$  which recently has been updated by Ericson, Loiseau and Thomas [14] to  $a_{\text{phen, ELT}}^+ = (-0.0017 \pm 0.0002(\text{stat}) \pm 0.0008(\text{sys})) m_{\pi}^{-1}$ . In that work it was also mentioned that  $a^+$  is also a major source of uncertainty in the extraction of the coupling constant  $g_{\pi NN}$  [14]. With the negative value for  $a^+$  given in [14],  $g_{\pi NN}^2/(4\pi) = 14.17 \pm 0.20$  is extracted in contrast to most other approaches which favour  $g_{\pi NN}^2/(4\pi) \approx 13.7 \pm 0.1$ , see Table I in [14].

Given the unsatisfactory experimental situation for  $a^+$ , it seems preferable to take into account binding and pion re-scattering effects between two nucleons in a more model-independent fashion, in order to constrain  $a^+$  from pion deuteron scattering. Such a framework is provided by  $\chi$ Pt which incorporates the chiral symmetry of low-energy QCD. Employing the so-called Weinberg counting [16] in which the pions are treated non-perturbatively and iterated to infinite order in ladder exchange diagrams, Beane et al. [17] calculated  $a_{\pi d}$ . Their result for  $a_{\pi d}$  does not involve any undetermined parameters, however, the non-perturbative effects responsible for nuclear binding are accounted for using phenomenological deuteron wavefunctions which introduces an inevitable model dependence. It is not clear to what extent the phenomenologically based deuteron wavefunctions employed in [17] are constrained from chiral symmetry, and one may therefore pose the question:

Does chiral symmetry predict the pion deuteron scattering length or is additional phenomenological input needed in order to fix new unknown parameters?

Clearly, this cannot be answered within the Weinberg framework which always employs deuteron wavefunctions in order to account for the non-perturbative behaviour of pion ex-

change. One must resort to a different approach which is given by the method of Kaplan, Savage and Wise [18] (ENT(KSW)) where, in contradistinction to the Weinberg scheme, pion exchange between the two nucleons is treated perturbatively. It has been illustrated in a number of papers that ENT(KSW) calculations at next-to-leading order (NLO) are – for small momenta – in good agreement with experimental data. At higher orders, however, the perturbative expansion is expected to fail in the deuteron channel [19]. Since our primary interest is to clarify the existence and importance of contributions to the pion deuteron scattering length not fixed by chiral symmetry, we will restrict ourselves to a next-to-leading calculation in the KSW scheme. As we will see, this issue can already be investigated at next-to-leading order so that we are not concerned with problems arising at higher orders.

The article is organised as follows: We start by presenting the Lagrangean (Sect. 2). In Sect. 3, the  $\pi d$  scattering amplitude at zero momentum and its scattering length is presented, followed by a discussion of our findings in Sect. 4, including the comparison to the EFT calculation of Beane et al. [17]. The final Section contains our conclusions. Two Appendices summarise details of the calculation and renormalisation procedure.

## 2 The Lagrangean

We now present the terms of the most general chirally invariant Lagrangean consisting of contact interactions between non-relativistic nucleons, and between nucleons and pions, which are relevant for our NLO calculation.

The pertinent terms satisfying the QCD symmetries are in the zero and one nucleon sector (see e.g. [20])

$$\begin{aligned}
\mathcal{L}_{\pi, \pi N} = & \frac{f_\pi^2}{8} \text{tr}[(D_\mu \Sigma^\dagger)(D^\mu \Sigma)] + \frac{f_\pi^2}{4} \omega \text{tr}[\mathcal{M}_q(\Sigma^\dagger + \Sigma)] + \\
& + N^\dagger(iD_0 + \frac{\vec{D}^2}{2M})N + g_A N^\dagger \vec{A} \cdot \vec{\sigma} N + \\
& + N^\dagger \left[ 2\omega c_1 \text{tr}[\mathcal{M}_q(\Sigma + \Sigma^\dagger)] + 4 \left( c_2 - \frac{g_A^2}{8M} \right) A_0^2 + 4c_3 A^\mu A_\mu \right] N + \dots ,
\end{aligned} \tag{2.1}$$

where  $N = \begin{pmatrix} p \\ n \end{pmatrix}$  is the nucleon doublet of two-component spinors,  $M = 938.918$  MeV the iso-scalar nucleon mass, and  $\sigma$  ( $\tau$ ) the Pauli matrices acting in spin (iso-spin) space. The

field  $\xi$  describes the relativistic pion, for which we choose the sigma gauge for convenience,

$$\Sigma(x) = \xi^2(x) = \sqrt{1 - 2 \frac{\pi^a \pi^a}{f_\pi^2}} + i\sqrt{2} \frac{\pi^a \tau^a}{f_\pi} . \quad (2.2)$$

$D_\mu$  is the chirally covariant derivative  $D_\mu = \partial_\mu + V_\mu$ , and the vector and axial currents are

$$V_\mu = \frac{1}{2}(\xi \partial_\mu \xi^\dagger + \xi^\dagger \partial_\mu \xi) , \quad A_\mu = \frac{i}{2}(\xi \partial_\mu \xi^\dagger - \xi^\dagger \partial_\mu \xi) . \quad (2.3)$$

The pion decay constant is normalised to be  $f_\pi = 130$  MeV,  $g_A = 1.27$ ,  $\mathcal{M}_q = \text{diag}(\hat{m}, \hat{m})$  is the quark mass matrix where we work in the iso-spin limit  $m_u = m_d = \hat{m}$ , and the constant  $\omega$  is chosen such that  $m_\pi^2 = 2\omega\hat{m}$  at the order examined here, where we use the iso-scalar value  $m_\pi = 138.039$  MeV for the pion mass. The coefficients  $c_1, c_2, c_3$  encode high-energy physics integrated out in HB $\chi$ PT and need at present to be determined by experiment. Since at the scales considered, the momenta of the nucleons are small compared to their rest mass, the nucleons are treated non-relativistically at leading order in the velocity expansion, with relativistic corrections systematically included at higher orders. Thus, the relativistic HB $\chi$ PT Lagrangean is reduced to the form shown above.

The germane terms in the two nucleon Lagrangean are (see also [19])

$$\begin{aligned} \mathcal{L}_{NN, \pi NN} = & - C_0 (N^T P^i N)^\dagger (N^T P^i N) + \frac{C_2}{8} \left[ (N^T P^i N)^\dagger (N^T P^i (\vec{D} - \overleftarrow{D})^2 N) + \text{H.c.} \right] - \\ & - \frac{\omega}{2} D_2 \text{tr}[\mathcal{M}_q(\Sigma + \Sigma^\dagger)] (N^T P^i N)^\dagger (N^T P^i N) + \\ & + \left( E_2 \text{tr}[A_0^2] - F_2 \text{tr}[\vec{A}^2] \right) (N^T P^i N)^\dagger (N^T P^i N) + \dots , \end{aligned} \quad (2.4)$$

where  $P^i$  is the projector onto the S wave of the iso-scalar-vector channel,

$$P_{a\alpha}^{i, b\beta} = \frac{1}{\sqrt{8}} (\sigma_2 \sigma^i)_\alpha^\beta (\tau_2)_a^b , \quad (2.5)$$

and the parameters of the Lagrangian  $C_i, D_2, E_2$  and  $F_2$  are not constrained by chiral symmetry, but can be extracted from experimental input or estimated with additional model dependent assumptions. E.g.,  $C_0$  can be related to the binding energy of the deuteron and  $C_2$  to the nucleon-nucleon scattering length [18]. The remaining parameters  $D_2, E_2$  and  $F_2$ , on the other hand, have not been determined yet.

### 3 Calculation

The pion deuteron scattering length  $a_{\pi d}$  follows from the amplitude  $\mathcal{A}_{\pi d}$  at zero momentum,

$$a_{\pi d} = \frac{1}{4\pi} \left[ 1 + \frac{m_\pi}{2M} \right]^{-1} \mathcal{A}_{\pi d} \quad , \quad (3.1)$$

which in turn is decomposed into a contribution in which the pion scatters off only one nucleon (Fig. 1), and a term with two nucleon interactions (Fig. 2),

$$\mathcal{A}_{\pi d} = \mathcal{A}_{\pi d}^{\text{1body}} + \mathcal{A}_{\pi d}^{\text{2body}} \quad . \quad (3.2)$$

The first amplitude starts at LO,  $\mathcal{O}(Q^2)$  after wave function renormalisation, while the second one is NLO,  $\mathcal{O}(Q^3)$ .

#### 3.1 One Body Contributions

We first discuss briefly how to embed the well known HB $\chi$ PT result for the iso-scalar S wave pion nucleon scattering amplitude  $\mathcal{A}^+$  into the deuteron. The one body contributions to  $\pi d$  scattering as shown in Fig. 1 consist of the LO and NLO iso-scalar pion nucleon amplitude  $\mathcal{A}^+$  at zero momentum (diagrams (1a) and (1b)), corrections arising from wave function renormalisation and NLO deuteron effects (diagrams (1c)). Since the latter diagrams can be absorbed into a re-definition of the deuteron source used, it is no surprise that their contribution to  $\mathcal{A}_{\pi d}$  at zero momentum cancels with the NLO wave function renormalisation of the LO amplitude.

As the pion can scatter off the proton or neutron inside the deuteron, the one body amplitude is twice the physical amplitude  $\mathcal{A}^+$  only, independent of any deuteron observable. This is formally confirmed in an exemplary calculation in App. A. Relativistic effects do not enter at the order we calculate. Therefore,

$$\mathcal{A}_{\pi d}^{\text{1body}} = \frac{4m_\pi^2}{f_\pi^2} \left[ \frac{\Delta_{\text{bare}}(\mu)}{2} - \frac{g_A^2}{8M} + \frac{g_A^2(3m_\pi - 2\mu)}{64\pi f_\pi^2} \right] = 2\mathcal{A}^+ \quad (3.3)$$

with the bare quantity  $\Delta_{\text{bare}}(\mu) := 2(c_{2,\text{bare}}(\mu) + c_{3,\text{bare}}(\mu) - 2c_{1,\text{bare}}(\mu))$ . Notice that both graphs (b1) and (b2) in Fig. 1 depend on the renormalisation scale  $\mu$ , as the nested integral




Figure 1: *The contributions from  $\mathcal{A}^+$ : (a): LO; (b): NLO in  $\mathcal{A}^+$ ; (c): NLO deuteron corrections. Shaded circles denote the deuteron source. Graphs obtained by permuting vertices or external lines are not displayed.*

is linearly divergent. In the  $\overline{\text{MS}}$  or  $\overline{\text{MS}}$  scheme usually chosen in  $\text{HB}\chi\text{PT}$ , the divergence is discarded ( $\Delta_{\text{bare}}(\mu) = \Delta^{\overline{\text{MS}}}$ ), but it is manifest in the PDS scheme which for consistency has of course to be used also in the one nucleon part of the calculation. Therefore, the  $\text{HB}\chi\text{PT}$  parameters  $c_1$ ,  $c_2$ ,  $c_3$  start to depend on the renormalisation procedure at NLO and are not observables. It is however obvious that the regulator dependence is absorbed into the NLO,  $\mathcal{O}(Q)$  part of the combination of coefficients  $\Delta(\mu) = \Delta^{(0)} + \Delta^{(1)}(\mu)$ , in agreement with the power counting. The LO part,  $\Delta^{(0)}$ , is  $\mu$  independent. The value for  $\Delta$  in the  $\overline{\text{MS}}$  scheme is hence easily translated into the PDS scheme as  $\Delta_{\text{bare}}(\mu) = \Delta^{\overline{\text{MS}}} + \frac{g_A^2 \mu}{16\pi f_\pi^2}$ , making the full, physical one body amplitude (3.3) explicitly  $\mu$  independent.

Recently, updated values for the parameters  $c_i$  have been determined in an  $\mathcal{O}(Q^3)$  and  $\mathcal{O}(Q^4)$   $\text{HB}\chi\text{PT}$  fit to three finite energy pion nucleon scattering analyses [3, 4]. As already noted in Refs. [1, 20], the iso-scalar S wave scattering length

$$a^+ := \frac{1}{4\pi} \left[ 1 + \frac{m_\pi}{M} \right]^{-1} \mathcal{A}^+ \quad (3.4)$$

can however not be determined precisely because of a numerical cancellation which may signal physics at small scales not yet understood. The  $\text{HB}\chi\text{PT}$  analysis predicts the range [3,

4]

$$a_{\text{HB}\chi\text{PT}}^+ = [-0.01 \dots + 0.006] m_\pi^{-1} = (-0.002 \pm 0.008) m_\pi^{-1} , \quad (3.5)$$

compatible with the various experimental and phenomenological extractions discussed in the Introduction. The comparatively large range comes from the use of different partial wave analyses for the  $\pi N$  amplitudes; the theoretical uncertainty from HB $\chi$ PT is considerably smaller.

### 3.2 Two Body Contributions

Processes involving deuteron correlations, Fig. 2, enter at NLO,  $\mathcal{O}(Q^3)$ . The sum of the two diagrams Fig. 2 (a1) and (a2) is independent of the parametrisation of the pion field. The two pion one nucleon vertices in (b) stem from the chirally covariant form of the time derivative in the kinetic energy term for the nucleon in (2.1).




Figure 2: *The two body contributions from pion re-scattering (a,b) and two body contact interactions (c) at NLO. Graphs obtained by permuting vertices or external lines are not displayed.*

The diagrams with a pion in the intermediate state, Figs. 2 (a) and (b), contain logarithmic divergences. In order to see how they manifest themselves in a more traditional setting, consider the graphs when a deuteron wave function is used instead of the point-like source with the correct quantum numbers which is the starting point of the standard field theoretical treatment. In this case, the contributions are proportional to the expectation values of two operators [17, 21]

$$\mathcal{A}_{\pi d, \text{trad}}^{\text{2body} (a)} \propto \left\langle \frac{(\vec{q} \cdot \vec{\sigma}_1)(\vec{q} \cdot \vec{\sigma}_2)}{(\vec{q}^2 + m_\pi^2)^2} \right\rangle ,$$



$$\mathcal{A}_{\pi d, \text{trad}}^{2\text{body}(b)} \propto \left\langle \frac{1}{\vec{q}^2} \right\rangle \quad (3.6)$$

between deuteron wave functions, where  $\vec{q}$  is the momentum transfer between the nucleons. Thus, the graphs seem to probe the deuteron only at momenta of the order of  $q \sim m_\pi$  in Fig. 2 (a) and  $q \sim 0$  in Fig. 2 (b). The Fourier transformation into position space reveals however that the expectation values of the operators  $\frac{1}{r}$  and  $\frac{e^{-m_\pi r}}{r}$  are probed, testing the deuteron wave function at arbitrarily short distance. Usually, the UV part of the deuteron wave function is parameterised by adding to the long range pionic potentials strong phenomenological terms mimicking short range repulsion. The deuteron wave function is thus suppressed at large momentum transfer at the price of adding some dependence on the shape and parameters of the short distance potential. Thus, the result for these amplitudes depends also on short distance physics subsumed into an arbitrary, unphysical cut-off parameter.

It is therefore no surprise that a logarithmic divergence appears in the pion exchange diagrams. The necessary integrals are tabulated in [22], so that the bare amplitudes are

$$\begin{aligned} \mathcal{A}_{\pi d, \text{bare}}^{2\text{body}(a)} &= \frac{g_A^2 m_\pi^2 \gamma}{12\pi f_\pi^4} \left( \Gamma - \frac{m_\pi}{m_\pi + 2\gamma} - 2 \ln \left[ \frac{m_\pi + 2\gamma}{\mu} \right] \right) \\ \mathcal{A}_{\pi d, \text{bare}}^{2\text{body}(b)} &= - \frac{m_\pi^2 \gamma}{\pi f_\pi^4} \left( \Gamma - 2 \ln \left[ \frac{2\gamma}{\mu} \right] \right) \end{aligned} \quad (3.7)$$

with a divergence  $\Gamma := \Gamma[4 - d] + \ln[\pi] + 1$  in  $d \rightarrow 4$  space-time dimensions<sup>1</sup>. Therefore, the bare two pion two nucleon contact interactions Fig. 2 (c1) and (c2) entering at the same order are also necessary in order to consistently remove all regulator dependence in the total, physical two body amplitude.

$$\mathcal{A}_{\pi d, \text{bare}}^{2\text{body}(c)} = \frac{m_\pi^2 \gamma}{\pi f_\pi^2} [D_{2, \text{bare}}(\mu) + E_{2, \text{bare}}(\mu)] (\gamma - \mu)^2 \quad (3.8)$$

They serve the same purpose as suppressing the deuteron wave function at short distance by a cut-off. As must be expected from the power counting, the combinations  $D_{2, \text{bare}}(\mu) (\gamma - \mu)^2$  and  $E_{2, \text{bare}}(\mu) (\gamma - \mu)^2$  scale as  $Q^0$ . The strengths of both two pion two nucleon contact

---

<sup>1</sup>Following [22], we prefer to work with a definition of  $\Gamma$  which is independent of the scale  $\mu$ , in contradiction to e.g. [20].

terms cannot be predicted at the present time. Albeit the parameter  $D_2$  is encountered as the chiral symmetry breaking contribution to  $NN$  scattering at NLO, in this process its renormalised value cannot be dis-entangled from the parameter  $C_0^{(0)}$  which respects chiral symmetry [19]. The strength  $E_2$  is not fixed, either. Unfortunately, we are also not able to extract their values from additional experimental input so that we cannot predict the pion deuteron scattering length.

Nonetheless, EFT allows to estimate the natural size of the combination  $D_2 + E_2$  after specifying a prescription to handle the divergences in three and four dimensions. Only after renormalisation is one justified to compare the sizes of the various contributions. In the PDS scheme, the theory is usually renormalised (and the power counting made manifest even before renormalisation) by choosing the renormalisation scale to be natural [18]:

$$\mu \rightarrow m_\pi, \quad \Gamma \rightarrow 2 \quad (3.9)$$

With the latter choice, the divergences in 4 dimensions are removed by demanding that the pionic contribution to zero momentum scattering between two nucleons disappears, see (B.1) in App. B. Power counting and dimensional analysis dictate then that

$$D_2 + E_2 = \frac{4\pi}{M \Lambda_{NN}} \frac{\mathcal{Z}}{(\mu - \gamma)^2}, \quad (3.10)$$

where the magnitude of the dimensionless parameter is of order unity,  $|\mathcal{Z}| \approx 1$ , if the naturalness assumption holds. Its sign remains undetermined. The scale  $\Lambda_{NN} \approx 300$  MeV enters in the chiral expansion for  $NN$  scattering [18] and subsumes all short distance physics, i.e. all effects of particles not contained as explicit degrees of freedom in the Lagrangean, like the  $\Delta$ ,  $\rho$  meson exchange etc.

However, it must be stressed that in the approach taken here, a decomposition of the physically observable  $\pi d$  scattering length into parts related to the pion re-scattering diagrams of Fig. 2 (a) and (b) separately is strictly speaking meaningless: None of these diagrams is renormalisation group invariant (i.e. cut-off independent) on its own, and only combinations of these diagrams with the two pion two nucleon contact diagrams of Fig. 2 (c) form observables free of divergences, i.e. independent of  $\mu$  and  $\Gamma$ . After renormalisation

performed in App. B, the scattering amplitude reads

$$\mathcal{A}_{\pi d}^{2\text{body}} = \frac{m_\pi^2 \gamma}{\pi f_\pi^4} \left[ 2 \ln \left[ \frac{2\gamma}{\Lambda^*} \right] - \frac{g_A^2}{6} \left( \frac{1}{2} \frac{m_\pi}{m_\pi + 2\gamma} + \ln \left[ \frac{(m_\pi + 2\gamma)}{\Lambda^*} \right] \right) \right] , \quad (3.11)$$

and contains only one un-determined, physical parameter  $\Lambda^*$ . This dimension-ful number parametrises the renormalisation group invariant strength of the contact interactions  $D_2$  and  $E_2$  between nucleons and subsumes effects from the deuteron wave function at short distances. It needs to be determined from experiment or from a microscopic calculation of  $NN$  scattering in QCD.  $\Lambda^*$  is expected to be of the order of the natural low energy scale ( $m_\pi$  or  $\Lambda_{\text{QCD}}$ ) since all dependence on higher scales integrated out has disappeared with renormalisation.

As mentioned above, relativistic effects do not enter at the order we are working. Nevertheless, it may be worthwhile to comment on their contributions. Relativistic corrections to the energy-momentum relation are accounted for by inserting perturbatively higher dimension operators, the lowest one being  $\vec{p}^4/(8M^3)$ . They start at N2LO and are suppressed by factors of the nucleon mass, and not by  $\Lambda_{NN} \approx 300$  MeV. Such effects are therefore small compared to other corrections which enter formally at the same order N2LO. Our calculation only includes NLO effects. For our purposes, they are therefore negligible, and may be regarded to lie within the given error bars.

## 4 Results and Discussion

With the data at hand, we cannot predict either the iso-scalar or the pion deuteron scattering length in a unique way from our calculation due to the unknown physical scale  $\Lambda^*$  stemming from the combination  $D_2 + E_2$  of counter terms. These unknown strengths of the two-pion two-nucleon couplings enter already at NLO, i.e. at the same order as pion re-scattering. In order to determine their sizes, we use as inputs the experimental value for the pion deuteron scattering length and a value for  $a^+$ . Clearly, requiring self-consistency forbids to consider phenomenological extractions of  $a^+$  into which  $a_{\pi d}$  entered. We use as choice either the value from the pionic hydrogen experiment,  $a_{\text{exp}, \pi N}^+ = (-0.0022 \pm 0.0043) m_\pi^{-1}$  [10], or

the HB $\chi$ PT prediction, Eq. (3.5) [3, 4]. Both are compatible with zero, and the HB $\chi$ PT result has an error bar accommodating also the phenomenological partial wave analysis extractions [5, 6, 7] cited in the Introduction. We find

$$\begin{aligned}\Lambda^* &= (262_{-109}^{+185}) \text{ MeV} && \text{when fitted to } a_{\text{exp}, \pi N}^+ \text{ [10]} \\ \Lambda^* &= (269_{-170}^{+458}) \text{ MeV} && \text{when fitted to } a_{\text{HB}\chi\text{PT}}^+ \text{ [3] ,}\end{aligned}\tag{4.1}$$

which contains a large error bar from the uncertainty in  $a^+$ . The value of  $\Lambda^*$  increases as  $a^+$  increases. The Logarithms in (3.11) are indeed of order 1.

We now consider the sizes of two renormalisation group invariant subsets constructed out of the two body scattering result (3.11): The first one is the term independent of  $g_A$ . The second one is quadratic in  $g_A$ , and is easily estimated to be suppressed by a factor 1/12. We reproduce the well known result that the dominant pion re-scattering contribution stems from physics unchanged by taking the chiral limit, and  $\Lambda^* \approx \Lambda_{E_2}^*$ . Albeit the definition of  $\Lambda^*$  depends also on  $g_A$  (B.8), this polynomial separation with respect to  $g_A$  is at present a good analogue to the values quoted in the Weinberg approach for the pion re-scattering contributions, Fig. 2 (a) and (b). Although much simpler, our results for these renormalisation group invariant combinations are close to the findings of the Weinberg approach, see Table 1. This should not be too much a surprise since it was shown above that the pion re-scattering diagrams test after renormalisation only the tail of the deuteron wave function, at momenta not larger than  $m_\pi$ . Figure 3 depicts the dependence of the two body contributions to the  $\pi d$  scattering length on  $\Lambda^*$ . We finally summarise all one and two body contributions to  $a_{\pi d}$  in Table 2 for the extraction using  $a_{\text{exp}, \pi N}^+$ .

A similar analysis can also be made for  $D_2 + E_2$ . Taking  $a_{\text{exp}, \pi N}^+$  from the pionic hydrogen experiment, we find using the prescription (3.10)

$$\mathcal{Z} = 1.0 \mp 1.5\tag{4.2}$$

with a large uncertainty due to the error bar in the value of  $a_{\text{exp}, \pi N}^+$ . For the central value  $\mathcal{Z} = 1.0$  the contributions from the individual diagrams are  $a_{\pi d}^{2\text{body}(a)} = +0.0005 m_\pi^{-1}$ ,  $a_{\pi d}^{2\text{body}(b)} = -0.028 m_\pi^{-1}$ ,  $a_{\pi d}^{2\text{body}(c)} = +0.006 m_\pi^{-1}$ . We note that choosing a different prescription for

approach	$a_{\pi d}^{2\text{body}}(g_A^2) [m_\pi^{-1}]$	$a_{\pi d}^{2\text{body}}(g_A^0) [m_\pi^{-1}]$
KSW , RG invariant	-0.0004	-0.0210
Weinberg	$-0.0007 \pm 0.0002$	$-0.0196 \pm 0.0005$

Table 1: Comparison of two body corrections to  $a_{\pi d}$  in the KSW power counting with  $\Lambda^*$  determined from  $a_{\text{exp}, \pi N}^+$  [10], and in the hybrid approach based on Weinberg's power counting taken from [17]. Error bars from the uncertainty in  $a_{\text{exp}, \pi N}^+$  omitted. The error bars in the Weinberg approach stem from using different model wave functions for the deuteron.




Figure 3: Sizes of the renormalisation group invariant contributions to  $a_{\pi d}$ , (3.11), depending on the physical parameter  $\Lambda^*$ . Dashed line: contribution binomial in  $g_A$ ; dash-dotted line: contribution independent of  $g_A$ ; solid line: total.

removing the 4 dimensional UV divergences,  $\Gamma \rightarrow 1$ , leads to pion re-scattering contributions numerically deviating by less than 8% from those obtained by Beane et al. [17]:  $a_{\pi d}^{2\text{body} (a)}(\Gamma \rightarrow 1) = -0.0008 m_\pi^{-1}$ ,  $a_{\pi d}^{2\text{body} (b)}(\Gamma \rightarrow 1) = -0.019 m_\pi^{-1}$ ,  $a_{\pi d}^{2\text{body} (c)}(\Gamma \rightarrow 1) = -0.002 m_\pi^{-1}$  with  $\mathcal{Z}(\Gamma \rightarrow 1) = -0.4 \mp 1.5$ . As the error bars indicate, the magnitude of  $D_2 + E_2$  increases as  $a^+$  increases. Again, the contribution from the counter terms is clearly not unnaturally large and is roughly 10 – 20% of the dominating pion re-scattering amplitude  $a_{\pi d}^{2\text{body} (b)}$ . This implies that a 10% uncertainty in the parameters  $D_2$  and  $E_2$  yields an error of about 1% in the total scattering length. The effect of the counter terms is thus even smaller than the

diagram	contribution to $a_{\pi d}$ [ $m_\pi^{-1}$ ]
1 body from $a_{\text{exp}, \pi N}^+$	$-0.005 \pm 0.009$
2 body, total	$-0.021 \mp 0.009$
2 body, $g_A^0$	$-0.021 \mp 0.009$
2 body, $g_A^2$	$-0.0004 \pm 0.0014$
$a_{\pi d}^{\text{exp}}$	$-0.0261 \pm 0.0005$

Table 2: *One and two body corrections to  $a_{\pi^0 d}$ , choosing as input the experimental extraction of  $a^+$  from pionic hydrogen experiments [10].*

power counting suggests.

One may also assume that the two body counter terms  $D_2$  and  $E_2$  are saturated by a mechanism which the phenomenological extraction of  $a^+$  from the pion deuteron scattering length by Ericson et al. [14] can capture correctly. Taking their value  $a_{\text{phen, ELT}}^+$ , we obtain  $\Lambda^* \approx 280$  MeV,  $\mathcal{Z} \approx 0.8$ . However, such an approach violates the spirit of self-consistency at the basis of our calculation.

## 5 Conclusions

We presented a calculation of the pion deuteron scattering length in Effective Field Theory with perturbative pions. In this scheme proposed by Kaplan, Savage and Wise, knowledge of the iso-scalar pion nucleon scattering length  $a^+$  does not suffice to determine  $a_{\pi d}$  directly due to two unknown short distance effective interactions coupling two nucleons to two pions. Their strengths  $D_2$  and  $E_2$  can be subsumed into one physical unknown which we estimated in the range  $\Lambda^* \sim 270$  MeV, but with sizeable error bars. Employing  $\Lambda^* \sim 270$  MeV, we can reproduce the numerical results for the pion re-scattering graphs in the work by Beane et al. [17] who use the Weinberg scheme. Unfortunately, the experimental error on  $a^+$  is not small enough to constrain  $\Lambda^*$  (or  $D_2 + E_2$ ) to a more precise value. The 10% accuracy

assigned to our NLO calculation is not only comparable to the uncertainty induced by experiment in the sizes of our counter terms. It also masks iso-spin breaking contributions to  $a^+$ , which were shown in [8] not to exceed 7% of the pion nucleon scattering length. A more accurate measurement of  $a^+$  not involving pion deuteron scattering data, as by the present PSI experiment R-98.01 [11], will substantially reduce the uncertainty.

We can therefore answer the question posed in the Introduction as follows: Chiral symmetry in combination with already known parameters does not suffice to determine the pion deuteron scattering length due to two new contact terms with unknown couplings. However, once these terms are fixed from the pion deuteron scattering length they may be used to predict pion deuteron scattering data at non-zero momentum transfer.

## Acknowledgements

We are grateful to S. Beane, T. Ericson, T. Hemmert, N. Kaiser, D. Phillips, A. Thomas, U. van Kolck and W. Weise for discussions and useful remarks. D. Gotta drew our attention to the ongoing experiments on pionic hydrogen at PSI. The INT (Seattle) provided generous hospitality during part of this work (H.W.G.). We also acknowledge financial support by the DFG Sachbeihilfen GR 1887/1-1 and GR 1887/2-1 (H.W.G.) and by the Bundesministerium für Bildung und Forschung and the Deutsche Forschungsgemeinschaft.

## A Calculating the One Body Scattering Amplitude

As the pion can scatter off the proton or neutron inside the deuteron, the one body amplitude is (up to relativistic corrections) expected to be given by twice the physical amplitude  $\mathcal{A}^+$  only, irrespective of the presence of nested loops at NLO. We can confirm this formally and turn as an example to Fig. 1 (b1). After the integration over the loop energy of the larger loop, the two loop integral to be performed is using the loop momentum assignment indicated in Fig. 1 (b1)

$$\int \frac{d^3q}{(2\pi)^3} \frac{d^4\ell}{(2\pi)^4} \frac{1}{(\vec{q}^2 + \gamma^2)^2} \frac{1}{Ml_0 - \gamma^2 - \frac{\vec{q}^2 + (\vec{\ell} - \vec{q})^2}{2}} \frac{\vec{\ell}^2}{(l_0^2 - \vec{\ell}^2 - m_\pi^2)^2} , \quad (\text{A.1})$$

where the second term comes from the nucleon propagator inside the nested loop, and the last term is the pion propagator. Following the threshold expansion technique [23, 24], we identify two poles in the energy integration of the nested loop: (1)  $Ml_0 \sim \vec{q}^2$ ,  $\vec{\ell}^2 \sim Q^2$  from the nucleon propagator, and (2)  $l_0 \sim |\vec{\ell}| \sim m_\pi \sim Q$  from the pion. In the first case, the pion propagator is expanded because  $l_0 \ll |\vec{\ell}|, m_\pi$ , so that the pion becomes instantaneous. Since the nucleon propagates forward in time, this contribution is expected to vanish. The dimensionally regularised loop integral over  $l_0$  is indeed zero as no external scale is present in the denominators. In the second case, the pion pole is picked, and the nucleon propagator is expanded into

$$\frac{1}{Ml_0 - \gamma^2 - \frac{\vec{q}^2 + (\vec{\ell} - \vec{q})^2}{2}} \rightarrow \frac{1}{Ml_0} \left[ 1 + \frac{\gamma^2 + \frac{\vec{q}^2 + (\vec{\ell} - \vec{q})^2}{2}}{Ml_0} + \dots \right] . \quad (\text{A.2})$$

The nucleon propagator becomes static as in HB $\chi$ PT, and the two integrals factorise at LO in the expansion. The (relativistic) corrections are suppressed by powers of  $\frac{Q}{M} \approx \frac{1}{7}$ , i.e. negligible in a NLO calculation.

We therefore find (3.3) as conjectured to NLO, independent of any deuteron observable:

$$\mathcal{A}_{\pi d}^{\text{1body}} = \frac{4m_\pi^2}{f_\pi^2} \left[ \frac{\Delta_{\text{bare}}(\mu)}{2} - \frac{g_A^2}{8M} + \frac{g_A^2(3m_\pi - 2\mu)}{64\pi f_\pi^2} \right] = 2\mathcal{A}^+ \quad (\text{A.3})$$



## B Renormalising the Two Body Scattering Amplitude

We will now construct the renormalisation group invariant (and hence physical) combinations of amplitudes in the two body sector of  $\pi d$  scattering at threshold. As mentioned above, the parameter  $D_2$  enters in  $NN$  scattering at NLO together with the parameter  $C_0^{(0)}$  which respects chiral symmetry. There, it is also needed to absorb the four dimensional UV divergence stemming from the one pion exchange diagram. We recall [18] that the pertinent




Figure 4: *The contributions to  $NN$  scattering in the  ${}^3S_1$  channel at NLO in  $ENT(KSW)$  which completely determine the renormalisation of  $D_2$ .*

terms to renormalise  $D_2$  in the triplet S wave channel of  $NN$  scattering at NLO, depicted in Fig. 4, are in the centre-of-mass frame between nucleons with relative momentum  $p$  from one pion exchange

$$\mathcal{A}_{NN, \text{bare}}^{\pi \text{ exchange}} = - \left( \frac{M \mathcal{A}_{-1}(p)}{4\pi} \right)^2 \frac{g_A^2}{4f_\pi^2} \left[ 2(\mu + ip)^2 - m_\pi^2 \left( \Gamma - 2 \ln \left[ \frac{m_\pi - 2ip}{\mu} \right] \right) \right] \quad (\text{B.1})$$

and from the contact interactions involving  $D_2$  and  $C_0^{(0)}$

$$\mathcal{A}_{NN, \text{bare}}^{\text{contact}} = - \left( m_\pi^2 D_{2, \text{bare}}(\mu) + C_{0, \text{bare}}^{(0)}(\mu) \right) \left( \frac{\mathcal{A}_{-1}(p)}{C_0^{(-1)}} \right)^2, \quad (\text{B.2})$$

where  $C_0^{(-1)}$  is the LO part of the strength of the two nucleon contact interaction  $C_0$ , and the connected piece of the LO  $NN$  scattering amplitude (lower line of Fig. 4) is

$$\mathcal{A}_{-1}(p) := - \frac{C_0^{(-1)}}{1 + \frac{C_0^{(-1)} M}{4\pi} (\mu + ip)}. \quad (\text{B.3})$$

The one pion exchange part involving powers of the arbitrary regularisation parameter  $\mu$  comes from the contact piece of one pion exchange which generates divergences in 3 dimensions manifest in the PDS scheme. It is independent of the pion mass, i.e. unchanged by the chiral limit, and its regulator dependence can hence be absorbed into the definition of the renormalised coupling strength  $C_{0, R}^{(0)} := C_{0, \text{bare}}^{(0)} + \frac{g_A^2}{2f_\pi^2}$  in accordance with the power counting [25]. On the other hand, the four dimensional divergence  $\Gamma$  giving rise to the logarithmic dependence on  $\mu$  is chiral symmetry breaking and hence needs to be balanced by the definition of the bare two nucleon coupling  $D_{2, \text{bare}}$ .  $D_2$  is renormalised by setting

$$D_{2, \text{bare}}(\mu) \left( \frac{4\pi}{MC_0^{(-1)}} \right)^2 := \frac{g_A^2}{4f_\pi^2} \left( \Gamma - 2 \ln \left[ \frac{\Lambda_{D_2}^*}{\mu} \right] \right) . \quad (\text{B.4})$$

At LO,  $C_0^{(-1)} = -\frac{4\pi}{M} (\mu - \gamma)^{-1}$  is determined by demanding the triplet S wave to exhibit a pole at the deuteron binding energy in (B.3). Therefore, one obtains finally

$$D_{2, \text{bare}}(\mu)(\mu - \gamma)^2 = \frac{g_A^2}{4f_\pi^2} \left( \Gamma - 2 \ln \left[ \frac{\Lambda_{D_2}^*}{\mu} \right] \right) . \quad (\text{B.5})$$

The combination of the divergent one pion exchange diagram and of the contact interaction depending on the renormalisation group variant parameter  $D_{2, \text{bare}}$  is traded for one, renormalisation group invariant parameter  $\Lambda_{D_2}^*$ . This dimension-ful, physical number parametrises the renormalisation group invariant strength of the chiral symmetry breaking contact interaction between nucleons which does not contain derivatives. It needs to be determined from experiment or from a microscopic calculation of  $NN$  scattering in QCD. As only a variation of the pion mass can dis-entangle the effects of  $C_0^{(0)}$  and  $D_2$  in  $NN$  scattering at NLO, this process cannot serve to determine  $\Lambda_{D_2}^*$  experimentally.

Inserting the definition (B.5) into the two body amplitude  $\mathcal{A}_{\pi d}^{2\text{body}}$ , (3.7/3.8) reveals that some dependence on  $\mu$  and on both logarithmic and power law divergences remains:  $D_2$  serves only as a partial counter term to the diagrams Fig. 2 (a) depending on  $g_A^2$ , and does not affect the divergence of the  $g_A^0$  diagram Fig. 2 (b). In the second, analogous step, the remaining divergences in the two body sector are easily shown to disappear by setting

$$E_{2, \text{bare}}(\mu)(\mu - \gamma)^2 := \frac{1}{f_\pi^2} \left( 1 - \frac{g_A^2}{3} \right) \left( \Gamma - 2 \ln \left[ \frac{\Lambda_{E_2}^*}{\mu} \right] \right) , \quad (\text{B.6})$$

where  $\Lambda_{E_2}^*$  is another dimension-ful quantity, parameterising the renormalisation group invariant strength of the quark mass independent coupling of two nucleons and two pions.

The two body sector of the renormalised amplitude of  $\pi d$  scattering at threshold,

$$\mathcal{A}_{\pi d}^{2\text{body}} = \frac{m_\pi^2 \gamma}{\pi f_\pi^4} \left[ 2 \ln \left[ \frac{2\gamma}{\Lambda_{E_2}^*} \right] - \frac{g_A^2}{6} \left( \frac{1}{2} \frac{m_\pi}{m_\pi + 2\gamma} + \ln \left[ \frac{(m_\pi + 2\gamma)(\Lambda_{D_2}^*)^3}{(\Lambda_{E_2}^*)^4} \right] \right) \right] , \quad (\text{B.7})$$

contains therefore two un-determined parameters. Pion deuteron scattering at non-zero momentum transfer will allow to separate the two, and hence also  $C_0^{(0)}$  from  $D_2$  in  $NN$  scattering, but at present we choose to represent the two by one common scale

$$(\Lambda^*)^{\frac{g_A^2}{12}-1} := (\Lambda_{D_2}^*)^{-\frac{g_A^2}{4}} (\Lambda_{E_2}^*)^{\frac{g_A^2}{3}-1} . \quad (\text{B.8})$$

The final answer for the two body scattering amplitude in terms of renormalised quantities is therefore given by (3.11)

$$\mathcal{A}_{\pi d}^{2\text{body}} = \frac{m_\pi^2 \gamma}{\pi f_\pi^4} \left[ 2 \ln \left[ \frac{2\gamma}{\Lambda^*} \right] - \frac{g_A^2}{6} \left( \frac{1}{2} \frac{m_\pi}{m_\pi + 2\gamma} + \ln \left[ \frac{(m_\pi + 2\gamma)}{\Lambda^*} \right] \right) \right] . \quad (\text{B.9})$$

The well known fact that dimensional regularisation (together with the employed renormalisation scheme) preserves chiral symmetry manifests itself in the observation that all appearing divergences are renormalised manifestly chirally invariant.

## References

- [1] V. Bernard, N. Kaiser and U.-G. Meißner: *Phys. Lett.* **B309**, 421 (1993) [hep-ph/9304275].
- [2] V. Bernard, N. Kaiser and U.-G. Meißner: *Phys. Rev.* **C52**, 2185 (1995) [hep-ph/9506204].
- [3] N. Fettes, U.-G. Meißner and S. Steininger: *Nucl. Phys.* **A640**, 199 (1998) [hep-ph/9803266].
- [4] N. Fettes and U.-G. Meißner: *Nucl. Phys.* **A676**, 311 (2000) [hep-ph/0002162].
- [5] R. Koch: *Nucl. Phys.* **A448**, 707 (1986).
- [6] E. Matsinos: *Phys. Rev.* **C56**, 3014 (1997).
- [7] SAID on-line programme, R. A. Arndt et al., <http://gwdac.phys.gwu.edu/>; M. M. Pavan and R. A. Arndt: *IN Newslett.* **15**, 171 (1999).
- [8] J. Gasser, M. A. Ivanov, E. Lipartia, M. Mojžiš and A. Rusetsky: hep-ph/0206068.
- [9] T. Becher and H. Leutwyler: *JHEP* **0106**, 017 (2001) [hep-ph/0103263].
- [10] H. C. Schröder *et al.*: *Phys. Lett.* **B469**, 25 (1999).
- [11] D. Gotta: “Measurement of the ground-state shift and width in pionic hydrogen to the 1% level: A new proposal at PSI”, in: *IN Newsletter* **15**, 276, proceedings of the 8th International Symposium on Meson-Nucleon Physics and the Structure of the Nucleon, Zuoz (CH), 15th – 21st August 1999; D. Gotta: priv. commun.
- [12] P. Hauser *et al.*: *Phys. Rev.* **C58**, 1869 (1998).
- [13] V. V. Baru and A. E. Kudryatsev: *Phys. Atom. Nucl.* **60**, 1475 (1997).
- [14] T. E. Ericson, B. Loiseau and A. W. Thomas: *Phys. Rev.* **C66**, 014005 (2002) [hep-ph/0009312].

- [15] A. W. Thomas and R. H. Landau: *Phys. Rept.* **58**, 121 (1980).
- [16] S. Weinberg: *Nucl. Phys.* **B363**, 3 (1991).
- [17] S. R. Beane, V. Bernard, T. S. Lee and U.-G. Meißner: *Phys. Rev.* **C57**, 424 (1998) [nucl-th/9708035].
- [18] D. B. Kaplan, M. J. Savage and M. B. Wise: *Phys. Lett.* **B424**, 390 (1998) [nucl-th/9801034]; *Nucl. Phys.* **B534**, 329 (1998) [nucl-th/9802075].
- [19] S. Fleming, T. Mehen and I. W. Stewart: *Nucl. Phys.* **A677**, 313 (2000) [nucl-th/9911001].
- [20] V. Bernard, N. Kaiser and U.-G. Meißner: *Int. J. Mod. Phys.* **E4**, 193 (1995) [hep-ph/9501384].
- [21] S. Weinberg: *Phys. Lett.* **B295**, 114 (1992) [hep-ph/9209257].
- [22] M. J. Savage and R. P. Springer: *Nucl. Phys.* **A686**, 413 (2001) [nucl-th/9907069].
- [23] M. Beneke and V. A. Smirnov: *Nucl. Phys.* **B522**, 321 (1998) [hep-ph/9711391].
- [24] H. W. Griebhammer: *Phys. Rev.* **D58**, 094027 (1998) [hep-ph/9712467].
- [25] P. F. Bedaque and H. W. Griebhammer: *Nucl. Phys.* **A671**, 357 (2000) [nucl-th/9907077].