

MANY-BODY THEORY AT EXTREME ISOSPIN

H. LENSKE, C.M. KEIL AND F. HOFMANN

*Institut für Theoretische Physik, Universität Giessen, Heinrich-Buff-Ring 16,
D-35392 Giessen, Germany*

E-mail: horst.lenske@theo.physik.uni-giessen.de

The structure of nuclei far off β -stability is investigated by nuclear many-body theory. In-medium interactions for asymmetric nuclear matter are obtained by (Dirac-) Brueckner theory thus establishing the link of nuclear forces to free space interactions. HFB and RPA theory is used to describe ground and excited states of nuclei from light to heavy masses. In extreme dripline systems pairing and core polarization are found to be most important for the binding, especially of halo nuclei. The calculations show that far off stability mean-field dynamics is gradually replaced by dynamical correlations, giving rise to the dissolution of shell structures.

1 Introduction

Nuclear many-body theory provides a very general scheme allowing to investigate the whole variety of nuclear systems from (purely theoretical) infinite nuclear matter and macroscopic objects as neutron star matter down to the microscopic level of finite nuclei. Testing the concepts and methods of nuclear many-body theory at the extremes as found in exotic nuclei is an important and challenging question for structure physics. The vast amount of new data on nuclei far off stability ¹ has initiated corresponding efforts on the theoretical side, ranging from new impacts on cluster models and *ab initio* shell model calculations for light nuclei to mean-field calculations over the entire mass range. New phenomena, like halo and skin formation, have been explained or predicted by nuclear theory, documenting the progress made over the last years.

Until very recently, data were mainly available for ground state properties of exotic nuclei. At present, the situation is changing because the progress in experimental techniques allows to obtain information also on dynamical processes in neutron- and proton-rich nuclei. For theory this means to extend the methods to a new sector of phenomena, e.g. inelastic and charge exchange response functions and the fragmentation patterns of single particle strength functions in breakup and transfer reactions. The description of such phenomena clearly requires new approaches, accounting properly for new effects as the coupling to the nearby continuum and extreme isospin. Since one is entering unexplored territories models providing intrinsically an

extrapolation scheme for interactions and the resulting nuclear dynamics are requested. Phenomenological approaches like Skyrme theory and relativistic mean-field theory are moderately successful in this respect, but their flexibility and generality is constrained by the assumed operator structure of the model Hamiltonian or Lagrangian and the ability to determine the model parameters by fits to data only.

An important alternative is to approach the problem microscopically. A clear advantage of such a program is that - at least in principle - investigations can be based on a systematic order-by-order hierarchy of interaction diagrams, as typical for many-body theory, thus avoiding ambiguities and taking advantage of results being tested independently in other regions of nuclear physics. Taking this as a guideline the approach presented in this contribution starts from calculating interactions in asymmetric nuclear matter by Brueckner theory. As discussed in section 2 applications to the equation of state of infinite matter, neutron star matter and neutron star structure calculations give information on the global properties of the interaction model, e.g. indicating the importance for going beyond the pure ladder approximation inherent to Brueckner theory.

Applications of the microscopic approach to finite nuclei are discussed in section 3, illustrating pairing in dripline nuclei for ^{11}Li , and core polarization far off stability is the subject of section 4 as examples for the transition into a new dynamical regime of correlation dynamics. A summary and an outlook are given in section 5.

2 Interactions in Asymmetric Nuclear Matter

Good confidence on the free NN-interaction has been obtained with field theoretical meson-exchange models like the Bonn potentials¹⁷, showing that the ladder approximation is an appropriate scheme in free space. In a medium, however, the situation is less certain in the sense that additional classes of diagrams will contribute, as illustrated in Fig.1. The in-medium pieces introduce an additional density dependence and contribute new types of operators, not found in free space interactions^{8,9}. Among those, the coupling of mesons to medium polarization modes and three body forces from intermediate excitations of nucleonic resonances (TBF) will alter also the isospin structure of interactions. Little is known about the dependence of these contributions on varying the proton-to-neutron composition.

In ref.² Dirac-Brueckner Hartree-Fock (DBHF) theory was used to investigate 2-body interactions in infinite matter over a large range of asymmetries. In-medium meson-nucleon coupling constants were extracted for the isoscalar




Figure 1. The in-medium NN scattering amplitude including 2- and 3-body contributions, respectively. The 3-body pieces, being of order $\sim \mathcal{O}(\rho)$, result from the coupling to nucleon resonances and polarization graphs.

σ and ω and the isovector ρ and δ mesons, respectively ². In all meson channels a pronounced dependence on the isoscalar bulk density is found while the dependence on the asymmetry is close to negligible. Hence, to a very good approximation in-medium strong interactions remain intrinsically independent of isospin thus conserving the fundamental isospin symmetry. Using these coupling constants in the density dependent relativistic hadron (DDRH) field theory ³ finite nuclei ⁴, hypernuclei ⁵ and neutron stars ⁶ are well described, thus underlining the success of a microscopic approach.

In Fig.2 the equation of state obtained in non-relativistic theory is shown, including results in ladder approximation only and with TBF. The TBF are seen to act in total *attractive* in the low density region, but turn to *repulsive* at high densities, reflecting their specific density dependence. The full D3Y interaction, including TBF, is used in the following sections.

3 Pairing at the Dripline

In nuclear matter pairing is found to be a low density phenomenon, having a maximum contribution to the energy density around $\frac{1}{3}$ of the saturation density $\rho_0 = 0.16 \text{ fm}^{-3}$. Therefore, it has to be expected that pairing is enhanced in nuclei with a pronounced low-density region. This is exactly the situation in 2- or multi-nucleon halo systems, and, to a lesser extent, in weakly bound skin nuclei. A prominent case is ^{11}Li which exist as a particle-stable system mainly because of the mutual interactions among the two valence neutrons. Their low separation energy, $S_{2n}=320 \text{ keV}$, points to the importance of continuum coupling. For a detailed description of the valence wave function the conventional BCS and HFB methods using representations in terms of mean-field wave functions are not suitable. Theoretically, continuum effects




Figure 2. Equation of state of symmetric nuclear matter in non-relativistic Brueckner theory without and with 3-body interactions. The saturation point are indicated by full circles. For comparison, variational result (dots) of the Urbana group ⁸ are also shown.

are properly described by solving the Gorkov-equations ¹⁴

$$(h - e_+) \Phi_+ - \Delta \Phi_- = 0 \quad , \quad (h - e_-) \Phi_- + \Delta \Phi_+ = 0 \quad (1)$$

for the hole and particle components ¹⁵ Φ_{\pm} , coupled by the pairing field Δ . The single particle energies are $e_{\pm} = \lambda \pm E$ where λ and E (≥ 0) denote the chemical potential and the quasiparticle energy, respectively. Φ_{\pm} will in general *not* coincide with the eigenfunctions of the mean-field Hamiltonian h , i.e. the states Φ_{\pm} are off the (mean-field) energy shell.

Particle-stability is given for $\lambda < 0$. Then, irrespective of the value of E , the hole-type solutions Φ_- are exponentially decaying for $r \rightarrow \infty$. For $E \leq |\lambda|$, an exponential asymptotic behaviour is also found for the particle-type components Φ_+ and a finite subset of discrete eigenvalues E is obtained. For $E > |\lambda|$, eq.(1) has to be solved with continuum wave boundary conditions for Φ_+ leading to single particle spectral functions being distributed continuously in energy. Hence the quasi-particle picture, underlying BCS theory and, to some degree, also discretized HFB theory, is replaced by a fully dynamical continuum description.

For like-particle ($S = 0, T = 1$) pairing protons ($q = p$) and neutrons ($q = n$) are paired only among themselves. The pairing fields are given in terms of the anomalous or pairing density matrices κ and the pairing interaction V_{SE} ¹⁸,

$$\Delta_q(r_1, r_2) = V_{SE}(r_1, r_2)\kappa_q(r_1, r_2) \quad . \quad (2)$$

If $\lambda < 0$, $\kappa_q(r)$ and $\Delta(r)$ are guaranteed to decay exponentially for $r \rightarrow \infty$. It is still an open question whether the free space or an in-medium singlet-even interaction should be used (see e.g.¹⁶).

Strength functions for ¹¹Li are displayed in Fig.3. The strong deviation from a pure mean-field or BCS description is apparent by observing that besides the expected s- and p-wave components also $d_{5/2,3/2}$ strength is lowered into the bound state region. Remarkably, the mean-field does not support neither bound $2s$ nor $1d_{5/2,3/2}$ single particle levels and their appearance is solely due to the continuum coupling introduced by pairing.

Theoretically, the proton ($q = p$) and neutron ($q = n$) densities in a systems like ¹¹Li are defined by

$$\rho_q(r) = \sum_{j\ell} \frac{2j+1}{4\pi} \int_{-\infty}^{\lambda} de_- v_{qj\ell}^2(e_-) |G_{qj\ell}(r, e_-)|^2 \quad , \quad (3)$$

where $\Phi_- = v^2 G$ was used¹⁸. The particle numbers are given by

$$N_q = \sum_{j\ell} (2j+1) \int_{-\infty}^{\lambda} de_- v_{qj\ell}^2(e_-) = \sum_{j\ell} (2j+1) n_{qj\ell} \quad , \quad (4)$$

The neutron partial wave occupation numbers $N_{j\ell} = (2j+1)n_{j\ell}$ for ¹¹Li are displayed in Fig.3. In stable nuclei discrete levels at $2\lambda < e_- < \lambda$ will contribute to eq.(3). In extreme dripline nuclei they are missing because of the smallness of $|\lambda|$. In Fig.4 the proton and neutron ground state densities are shown, multiplied by r^2 , emphasizing the differences in shape and the neutron halo component. High-energy elastic scattering of ¹¹Li on a proton target is well described with the HFB densities¹⁹. Measured ¹¹Li response functions were analyzed by QRPA calculations in²⁰.

4 Dynamical Core Polarization at the Dripline

Dynamical core polarization (DCP) is seen most clearly in nuclei with a single nucleon outside a core. Approaching the driplines the core nucleus by itself is already far off stability with a lower surface tension. Therefore, the restoring forces against external perturbations are reduced. Such "soft core" systems




Figure 3. Single particle spectral functions for s- and p-wave neutron states in ^{11}Li . The finite widths of the states are due to continuum coupling. The partial occupation numbers $N_{j\ell}$ are also shown, including d-wave contributions.

are, for example, found in the neutron-rich even-mass carbon isotopes. A good indicator is the existence of low-energy 2^+ states, decreasing in energy with increasing mass. Continuum QRPA calculations with a residual interaction derived in Landau-Fermi liquid theory from the D3Y in-medium interaction reproduce the systematics of 2^+ states rather well. This leads to the conclusion that they are mainly of vibrational nature rather than due to static




Figure 4. Proton and neutron ground state density distributions (weighted by r^2) for ^{11}Li . The pronounced neutron halo component formed by s-,p- and d-states is clearly visible.

deformation as assumed e.g. in ^{24}C .

The QRPA calculations predict a strong increase of the quadrupole polarizabilities in the carbon isotopes for increasing neutron excess with a maximum around $^{16,18}\text{C}$. This behaviour is due to the lowering of the first 2^+ state which in ^{12}C is located at $E_x = 4.44$ MeV and moves down to $E_x = 1.66$ MeV in ^{18}C . The dipole (or electric) polarizability, however, changes only within 10% over the $^{10-22}\text{C}$ isotopic chain.

In a system with core polarization the valence particle (or hole) obeys a non-static wave equation

$$(H_{MF} + \Sigma_{pol}(\varepsilon) - \varepsilon) \Psi = 0 \quad (5)$$

including the static (HFB) mean-field Hamiltonian H_{MF} and the non-local and energy-dependent polarization self-energy Σ_{pol} describing the rescattering of the nucleon off the core thereby exciting it into states of various multipoles and excitation energies followed by subsequent de-excitations back to the ground state (see Fig.5).

During these processes the particle can be scattered virtually into high lying orbitals. The quantum numbers of the intermediate $2p1h$ (or $1p2h$) configurations are only constrained by the requirement that spin and parity must match those of the leading particle configuration, given by the state moving with respect to the core ground state. In nuclear matter and finite nuclei these processes are known to give rise to a depletion of momentum space




Figure 5. Diagrammatic structure of mean-field (left) and core polarization interactions (right) of a nucleon in single particle state j . Interactions (meson exchange) are indicated by dashed lines. Core polarization leads to intermediate states ($j'J_c$) with a (QRPA) core excitation J_c and a single particle state j' .

Table 1. Energies, spins and ground state spectroscopic factors from core polarization calculations.

<i>Nucleus</i>	j^π	<i>Energy keV</i>	$S(j^\pi, g.s.)$
8B	$3/2^-$	130	0.73
${}^{11}Be$	$1/2^+$	510	0.74
${}^{17}C$	$5/2^+$	760	0.43
${}^{19}C$	$1/2^+$	263	0.41

ground state occupation probabilities^{22,23} from the step function momentum distribution of a Fermi gas of quasiparticles interacting only by a static mean-field. In practice, the core excitations are calculated in QRPA theory thus extending the static HFB picture in a consistent way to a dynamical theory.

Dynamical single particle self-energies Σ_{pol} affect separation energies and wave functions. In dripline nuclei these effects are strongly enhanced. In fact, dynamical self-energies provide the main source of binding for ${}^{19}C(1/2^+, g.s.)$ ¹⁸.

Theoretically, the DCP description amounts to use a multi-configuration ground state containing a single particle component - reminiscent of the static mean-field configuration - and a multitude of configurations where the valence particle is rescattered into other orbits by interactions with the core^{15,21}. Coupling to the lowest 2^+ and 3^- states only as advocated e.g. in²⁵ cannot account for the complexity of the process.

The DCP wave functions have been used to analyze longitudinal mo-

momentum distributions and one-nucleon removal cross sections in relativistic breakup reactions with secondary beams ^{26,27,28}, reproducing the data very satisfactorily. Results for energies and spectroscopic factors in the single neutron halo nuclei ¹¹Be and ¹⁹C and the single proton halo nucleus ⁸B ²⁶ are shown in Tab.1. In ¹¹Be core polarization is causing the reversal of 1/2⁺ and 1/2⁻ states by supplying an additional attractive self-energy in the 1/2⁺ channel. The single particle spectroscopic factor $S_n(1/2^*, g.s.)=0.75$ is in good agreement with recent transfer and breakup data. A more dramatic effect is found in ¹⁹C and also ¹⁷C where the g.s. components are strongly suppressed as seen from the small spectroscopic factors in Tab.1. Hence, the valence nucleon is no longer attached to a definite mean-field orbital but exists in a wave packet-like state spread over a certain range of shell model states. In other words, mean-field dynamics have ceased to be the dominant source of binding. Rather, the ¹⁷C and ¹⁹C results indicate a new type of binding mechanism in dripline nuclei where shell structures are dissolved. The mean-field has lost its dominant role and binding is produced by dynamical valence-core interactions.

5 Summary and Outlook

Asymmetric nuclear matter and nuclei far off stability were described by microscopic approaches. An important advantage over empirical models is the clean diagrammatic structure allowing to extend the theory in a well defined way by inclusion of an increasingly larger class of diagrams. This was illustrated on the example of 3-body contributions to in-medium interactions. Pairing and core polarization far off stability were discussed in some detail. The results show that close to the dripline nuclear dynamics change from mean-field to correlation dynamics. As an important consequence shell structures are dissolved. This reflects that the traditional understanding of nuclear structure as being dominated by a static potential plus some residual interactions cannot account for nuclear structure far off stability.

Acknowledgments

This work was supported in part by DFG, project Le 439/3, European Graduate School "Complex Systems of Hadrons and Nuclei", Giessen-Copenhagen, GSI Darmstadt and BMBF.

References

1. P.G. Hansen, A.S. Jensen, B. Jonson, *Ann. Rev. Phys. Sci. (N.Y.)* **45**, 591 (1995).
2. F. de Jong, H. Lenske, *Phys. Rev. C* **57**, 3099 (1998).
3. H. Lenske, F. Fuchs, *Phys. Lett. B* **345**, 355 (1995); C. Fuchs, H. Lenske, H. Wolter, *Phys. Rev. C* **52**, 3043 (1995).
4. F. Hofmann, C. Keil, H. Lenske, *Phys. Rev. C* , (publication scheduled for Aug. 2001); nucl-th/0007050.
5. C. Keil, F. Hofmann, H. Lenske, *Phys. Rev. C* **61**, 06401 (2000).
6. F. Hofmann, C. Keil, H. Lenske, *Phys. Rev. C* **64**, 025804 (2001); nucl-th/0008038.
7. P. Ring *et al.*, *Ann. Phys. (N.Y.)* **198**, 132 (1990).
8. V.J. Pandharipande *et al.*, *Phys. Rev. C* **58**, 1804 (1998).
9. S.A. Coon, M.T. Peña, *Phys. Rev. C* **48**, 2559 (1993); S.A. Coon, M.T. Peña, D.O. Riska, *Phys. Rev. C* **52**, 2925 (1995).
10. H. Lenske, G. Schrieder, *Euro. Phys. J. A* **2**, 41 (1998); C. Gund *et al.*, *Euro. Phys. J. A* **10**, 85 (2001), see also nucl-ex/0010005.
11. F. Hofmann, H. Lenske, *Phys. Rev. C* **57**, 183 (1998).
12. E. Chabanat *et al.*, *Nucl. Phys. A* **627**, 273 (1997).
13. F.T. Baker *et al.*, *Phys. Rep.* **289**, 235 (1997).
14. L.P. Gorkov, *Sov. Phys. JETP* **9**, 1364 (1959); see also ¹⁶.
15. H. Lenske, *J. Phys. G* **24**, 1429 (1998).
16. For a review see e.g. W. Nazarewicz, *Prog. Part. Nucl. Phys.* **46**, (2001); P.Ring, *ibid.*.
17. R. Machleidt, *Adv. Nucl. Phys. (N.Y.)* **19**, 189 (1989).
18. H. Lenske *et al.*, *Prog. Part. Nucl. Phys.* **46**, 187 (2001).
19. P. Egelhoff, *Prog. Part. Nucl. Phys.* **46**, (2001).
20. M. Zinser, H. Emling, H. Lenske *et al.*, *Nucl. Phys. A* **619**, 151 (1997).
21. F.J. Eckle *et al.*, *Phys. Rev. C* **39**, 1662 (1989) and *Nucl. Phys. A* **506**, 199 (1990).
22. F. de Jong, H. Lenske, *Phys. Rev. C* **54**, 1488 (1996).
23. J. Lehr *et al.*, *Phys. Lett. B* **483**, 324 (2000).
24. F. Nunes *et al.*, *Nucl. Phys. A* **596**, 171 (1996).
25. N. Vinh-Mau, *Nucl. Phys. A* **592**, 33 (1995).
26. M.H. Smedberg *et al.*, *Phys. Lett. B* **452**, 1 (1999).
27. T. Baumann *et al.*, *Phys. Lett. B* **439**, 256 (1998).
28. D. Cortina-Gil *et al.*, *Euro. Phys. J. A* **10**, 49 (2001).