

# A possible quark origin of two-pion emission in $e + N$ and $N + N$ collisions

I.T. Obukhovsky\*

Institute of Nuclear Physics, Moscow State University, Moscow 119899, Russia

December 10, 2018

## Abstract

The constituent quark model has been taken as a starting point for investigation of one- and two-pion emissions. A wide range of observables have been evaluated in the framework of the  ${}^3P_0$  model. Comparison of the results with the data (when it is possible) shows an approximate agreement.

## 1 Introduction

In recent works of the Moscow [1] and Moscow-Tuebingen [2] groups the  ${}^3P_0$  model of pion emission [3, 4] has been extended to the description of mesonic clouds in nucleonic systems. In this model the meson momentum distribution in the cloud replicates the quark momentum distribution in the nucleon or in the overlap region of a two-nucleon system, and thus the standard quark-shell-model technique can be used for evaluation of meson cloud contributions to varied processes. As a result the pion emission amplitudes of interest [e.g. the amplitudes  $M(N \rightarrow \pi + B)$  of virtual transitions  $N \rightarrow B + \pi$  for pion quasielastic knockout  $N + e \rightarrow e' + \pi + B$  with  $B = N, \Delta, N_{1/2+}(1440)$ , etc.] have been calculated by this technique without invoking the Lagrangian field theory coupling constants  $f_{\pi NB}$  and vertex form factors  $F_{\pi NB}(k^2)$  (mostly unknown). On the same footing the related two-nucleon vertices  $N + N \rightarrow {}^2 B + \pi$ , actual for quasielastic processes of a similar type (e.g.  $d + e \rightarrow e' + \pi + {}^2 B$ ), have been evaluated.

In reactions  $N + e \rightarrow e' + \pi + B_i$  ( $d + e \rightarrow e' + \pi + {}^2 B$ ) a pionic decay of the final baryon (dibaryon) leads to a second pion emission  $B_i \rightarrow N + \pi$  ( ${}^2 B \rightarrow N + N + \pi$ ) with a momentum distribution predicted by the  ${}^3P_0$  model as well. However, the predicted cross sections for respective coincidence experiments are too small [1] to be seen over the background. Nevertheless another two-pion processes, e.g. the virtual " $\sigma$ -meson" (correlated  $\pi\pi$  pair) emission, have been considered.

## 2 One-pion predictions

Table 1 shows the "spectroscopic factors" [1]  $S_N^{\pi B} = \int |\overline{\Psi_N^{\pi B}(\mathbf{k})}|^2 d^3 k$  for pion momentum distributions  $\Psi_N^{\pi B_i}(\mathbf{k})$  in varied channels  $N \rightarrow \pi + B_i$ ,  $B_i = N, \Delta, N^*$ , etc.

$$\Psi_p^{\pi B_i}(\mathbf{k}) \sim \frac{M(N \rightarrow \pi + B_i)}{E_N(\mathbf{k}) - M_N + \omega_\pi(\mathbf{k})}, \quad |\overline{M(N \rightarrow \pi + B_i)}|^2 = f_{\pi qq}^2 \frac{4M_N M_B}{m_\pi^2} C_{\pi N B_i}^2 \mathbf{k}^2 F_{\pi N B_i}(\mathbf{k}^2),$$

\*The talk at International Symposium MENU2001, the George Washington University, 26-31 July.

with coefficients  $C_{\pi NB_i}^2$  calculated algebraically in the framework of  ${}^3P_0$  model ( $f_{\pi qq} = 3/5 f_{\pi NN}$  is the effective  $\pi q\bar{q}$  coupling constant common for all the channels).

**Table 1.** Spectroscopic factors  $S_p^{B_i\pi}$  and pion decay widths  $\Gamma_{B_i}$  for some baryons.

$B_i$	$N$	$\Delta$	$N_{1/2-}$ (1535)	$N_{1/2+}$ (1440)
$C_{\pi NB_i}^2$	2	$\frac{64}{25}$	$\frac{16\omega_\pi(\mathbf{k}^2)}{75\mathbf{k}^2}$	$\frac{2}{27}$
$\Gamma_{B_i}$ (MeV)	-	63	52	47
$S_p^{B_i\pi}$	0.153	0.065	0.006	0.011

The calculated values  $\Gamma_{B_i}$  are realistic ones but they are no more than in a rough agreement with experimental data. Nevertheless the evaluations made on the basis of  ${}^3P_0$  model have some advantages over the standard phenomenological description. In the model all the form factors  $F_{\pi NB_i}(\mathbf{k}^2)$  depend only on two parameters:  $b$  and  $b_\pi/b$ , where  $b \approx 0.6$  fm and  $b_\pi \approx 0.3$  fm are standard free parameters of the constituent quark model (the baryon and pion radius correspondingly). It is interesting that the sum of all the calculated spectroscopic factors  $N_\pi = \sum_i S_N^{\pi B_i} = 0.24$ , which is said (by analogy with the nuclear cluster physics) to be "a total number of pions in the nucleon", shows an approximate agreement with the "experimental" value  $N_\pi = 0.18 \pm 0.06$  obtained previously [1] by fitting the old pion knock-out data [5].

### 3 Two-pion predictions

In the recent work [2] the  $\sigma$ -meson formation has been considered in both the nucleon and the  $NN$ -system. We start from the  ${}^3P_0$ -model effective  $\pi qq$  vertex [2, 4, 6]

$$H_\alpha^{(3)}(\boldsymbol{\rho}'_2, \boldsymbol{\rho}_2) = \frac{f_{\pi qq}}{m_\pi} \frac{e^{2i\mathbf{k}\cdot\boldsymbol{\rho}'_2/3}}{\sqrt{2\omega_\pi}(2\pi)^{3/2}} \hat{O}(\boldsymbol{\rho}'_2, \boldsymbol{\rho}_2) \tau_{-\alpha}^{(3)} \boldsymbol{\sigma}^{(3)} \cdot \left[ \frac{\omega_\pi}{2m_q} \left( \frac{2}{i} \nabla_{\rho_2} + \frac{2}{3} \mathbf{k} \right) + \left( 1 + \frac{\omega_\pi}{6m_q} \right) \mathbf{k} \right],$$

and an effective  $\sigma\pi\pi$  scalar coupling  $H_{\pi\pi\sigma} = g_{\pi\pi\sigma} \vec{\phi}_\pi \cdot \vec{\phi}_\pi \phi_\sigma$  ( $g_{\pi\pi\sigma} \approx 2 - 4$  GeV) with the vertex form factor  $F_{\pi\pi\sigma}(q^2)$  of a Gaussian form  $\sim \exp(-q^2 b_\sigma^2/2)$  (here  $\hat{O}$  is a non-local operator related to the pion  $q\bar{q}$  wave function and  $\boldsymbol{\rho}_2 = (\mathbf{r}_1 + \mathbf{r}_2)/2 - \mathbf{r}_3$  is a relative coordinate). The resulting expression for the  $\sigma NN$  coupling constant  $g_{\sigma NN} \equiv g_0 D_N(0)$  with  $g_0 = \frac{f_{\pi qq}^2}{m_\pi^2} \frac{g_{\sigma\pi\pi}}{m_q^2 b^3}$  depends only on the loop integral  $D_N(k)$  for the triangle diagram in Fig. 1. It has been shown that a realistic value  $g_{\sigma NN} \approx 5-10$  could be only obtained if one takes into account a *coherent superposition* of all the baryon intermediate states  $B_i^*$  in the triangle graph:  $N$ ,  $N_{1/2-}^*$ ,  $N_{3/2-}^*$ ,  $N_{1/2+}^{**}$ ,  $\Delta$ ,  $\Delta_{1/2-}^*$ ,  $\Delta_{3/2-}^*$ ,  $\Delta_{3/2+}^{**}$ . In the quark shell model these states correspond to the configurations up to  $2\hbar\omega$  h.o.  $s^3$ ,  $s^2p$ ,  $sp^2$ , and  $sp^2 - s^22s(2d)$  with all the possible values of spin  $S = 1/2, 3/2$ , isospin  $T = 1/2, 3/2$ , and Young tableau  $[f_X] = [3]$ , [21]. Summation over these states leads to the following expression:

$$D_N(k=0) = \frac{b}{256} \frac{1}{2\pi^2} \mathcal{P} \int_0^\infty q^2 dq \frac{e^{-\beta^2 q^2} f_0(q) m_\sigma}{\omega_\pi(q/2) (m_\sigma - 2\omega_\pi(q/2))} \left\{ \frac{5}{4} \frac{1 + q^2 b^2/12}{m_B - m_N + q^2/8m_B + \omega_\pi(q/2)} \right. \\ \left. - \frac{m_q^2}{\omega_\pi^2(q/2)} \frac{q^2 b^2}{2} \left[ \frac{1}{q^2/8m_N + \omega_\pi(q/2)} + \frac{4}{m_\Delta - m_N + q^2/8m_N + \omega_\pi(q/2)} \right] \right\}, \quad \beta^2 = \frac{b^2}{12} \left( 1 + \frac{3b_\sigma^2}{2b^2} \right),$$




Figure 1: Quark diagram of the  $N \rightarrow \sigma(\pi\pi) + N$  transition.




Figure 2: The  $N + N \rightarrow DB$  transition.

where the function  $f_0$  takes into account all the possible time orderings in the triangle diagram [2] (for simplicity the common mass  $m_B = 1500$  MeV is used for all the baryons except  $\Delta$  and  $N$ ). However, for realistic values of the free parameters  $b = 0.5$  fm,  $b_\sigma = 0.2$  fm and  $g_{\sigma\pi\pi} = 4$  GeV we obtain a relatively small value  $g_{\sigma NN} = 0.433$ . It shows that the high-momentum behaviour of vertex form factors  $N \rightarrow B^* + \pi$ ,  $B^* \rightarrow N + \pi$  and  $\pi + \pi \rightarrow \sigma$  plays an important role in deriving a realistic value of the coupling constant  $g_{\sigma NN}$ . The standard quark shell model with a characteristic (confinement) scale  $b \approx 0.5 - 1$  fm predicts too soft vertex form factors, and thus some generalization of the model is needed to incorporate a new scale parameter of about 0.2-0.3 fm (the inverse value of the characteristic chiral scale 0.6 - 1 GeV/c). The following modification of a simple Gaussian form:  $e^{-\beta^2 q^2} \rightarrow \frac{2}{3} e^{-\beta^2 q^2} + \frac{1}{3} e^{-(\beta/Z)^2 q^2}$ , with  $Z \simeq 2$  was used to imitate a power-like behaviour  $\sim \left(\frac{\Lambda^2}{q^2 + \Lambda^2}\right)$  of the vertices. At  $Z = 1.9$  (which corresponds to  $\Lambda = 0.9$  GeV/c) a more realistic value  $g_{\sigma NN} = 3.26$  has been obtained.

The developed model has been used for the two-nucleon system. A special feature of the above processes in the  $NN$  system (see Fig. 2) is that a sum over intermediate dibaryon six-quark states  ${}^2B_i^* + \pi$  should be taken into consideration (hand by hand with the sum  $\sum_i B_i^* + \pi$  for each nucleon). The quantum numbers of intermediate dibaryons are fixed by the selection rules. In the channel  $J^P = 0^+(ST = 01)$  the  ${}^2B^*$  has quantum numbers of the so-called  $d'$  dibaryon [6]  $d' = |s^5 p[51]_X L=1, [321]_{CS} ST = 10, J^P = 0^- \rangle$ , while in the "deuteron channel"  $J^P = 1^+(ST = 10)$  it has quantum numbers of the  $d''$  dibaryon (a counterpart of the  $d'$  for  $S \rightleftharpoons T$  interchange [2]). The calculated amplitude for the transition  $N + N \rightarrow {}^2B + \sigma(2\pi)$ , where  ${}^2B$  is a compact six-quark configuration  $s^6[6]_X$  (a "quark core" of a "dressed bag" state  $DB = {}^2B + \sigma(2\pi)$ ), leads to a non-diagonal potential  $V_{DB, N+N}$  for two coupled channels ( $DB$  and  $N+N$ ). The corresponding coupled-channel model of  $NN$  interaction proposed in Ref. [2] on the basis of the above mechanism leads to a very reasonable description of the  $NN$  scattering at intermediate energies [2]. The  $DB$  admixture in deuteron  $P_{DB}$  has been also calculated [2]:  $P_{DB} = 3.66\%$ . This small value seems to be very important for a high-precision description of the deuteron properties.

## Acknowledgments

The author gratefully acknowledges the contribution to the talk of all the collaborators of the Moscow and Moscow-Tuebingen groups: Profs. A. Faessler, V.I. Kukulin, V.G. Neudatchin, N.P. Yudin, and Drs. V.M. Pomerantsev and L.L. Sviridova. The work was supported in part by the Deutsche Forschungsgemeinschaft grant No. Fa-67/20-1 and the Russian Foundation for

Basic Research grants No. 01-02-04015 and 00-02-16117.

## References

- [1] V.G. Neudatchin, I.T. Obukhovsky, and N.P. Yudin, *Relativistic Nuclear Physics and Quantum Chromodynamics*, Proc. XIV Int. Seminar on High Energy Physics Problems (Editors: A.M. Baldin and V.V. Burov), JINR, Dubna (2000), vol. 1, p. 55; I.T. Obukhovsky, V.G. Neudatchin, and N.P. Yudin, the contributed paper in this issue, to be published.
- [2] V.I. Kukulín, I.T. Obukhovsky, V.N. Pomerantsev, and A. Faessler, *Physics of Atomic Nuclei* **64**, 1667 (2001).
- [3] A. LeYaouanc, Ll. Oliver, O. Pène, and J.-C. Raynal, *Hadron Transitions in the Quark Model*. Gordon and Breach Sci. Publ., N.Y., London, Paris. 1988.
- [4] F. Cano, P. Gonzalez, S. Noguera, and B. Desplanques, *Nucl. Phys.* **A343**, 331 (1996).
- [5] B. Brauel, T. Canzler, D. Cords, et al., *Z.Phys. C* **3**, 101 (1979).
- [6] I.T. Obukhovsky, A. Faessler, G. Wagner, and A. Buchmann, *Phys. Rev. C* **60**, 035207 (1999).