Structure of Sn isotopes beyond N = 82

L. Coraggio, A. Covello, A. Gargano, and N. Itaco

Dipartimento di Scienze Fisiche, Università di Napoli Federico II, and Istituto Nazionale di Fisica Nucleare. Complesso Universitario di Monte S. Angelo, Via Cintia, I-80126 Napoli, Italy (November 1, 2018)

We have performed shell-model calculations for 134,135 Sn using a realistic effective interaction derived from the CD-Bonn nucleon-nucleon potential. Comparison shows that the calculated results for 134 Sn are in very good agreement with recent experimental data. This supports confidence in our predictions of the hitherto unknown spectrum of ¹³⁵Sn.

PACS number(s): 21.60.Cs, 21.30.Fe, 27.60.+j

The Sn isotopes play an essential role in the shell-model description of nuclear structure, as they provide the opportunity to study the change of nuclear properties when varying the number of neutrons over a large range. From the experimental point of view, there is information for practically all isotopes with valence neutrons in the 50-82 shell, namely from doubly magic ¹³²Sn down to doubly magic ¹⁰⁰Sn. It is therefore of great relevance to try to go beyond 132 Sn, so as to test the shell-model predictions when having valence neutrons in the 82-126 shell. While this is not an easy task, since very neutron-rich Sn isotopes lie well away from the valley of stability, two very recent studies [1,2] have provided some experimental information on ¹³⁴Sn with two neutrons outside the doubly magic ¹³²Sn core.

During the past few years, we have studied several nuclei around 132 Sn [3] in terms of the shell model employing realistic effective interactions derived from modern nucleon-nucleon (NN) potentials. We have therefore found it interesting to perform calculations of this kind for ¹³⁴Sn and also to predict the spectroscopic properties of ¹³⁵Sn with three neutrons in the 82-126 shell, for which no experimental spectroscopic data are available as yet. In this paper, we present the results of our realistic shell model calculations starting with a brief description of how they have been obtained.

We consider ¹³²Sn as a closed core and let the valence neutrons occupy the six single-particle levels $h_{9/2}$, $f_{7/2}$, $f_{5/2}$, $p_{3/2}$, $p_{1/2}$, and $i_{13/2}$ of the 82-126 shell. The effective two-body matrix elements for the chosen model space have been derived from the CD-Bonn free NN potential [4]. This derivation has been performed within the framework of a G-matrix folded-diagram formalism, including renormalizations from both core polarization and folded diagrams. A detailed description of our derivation is given in Ref. [5]. We should point out, however, that the effective interaction used in the present work has been obtained including G-matrix diagrams through third order.

Let us now come to the single-particle (SP) energies. The value of $\epsilon_{f_{7/2}}$ has been fixed at -2.455 MeV, as determined from the experimental one-neutron separation energy for ¹³³Sn [6]. As regards the other five SP energies, our adopted values (relative to the $f_{7/2}$ level) are (in MeV): $\epsilon_{p_{3/2}} = 0.854$, $\epsilon_{h_{9/2}} = 1.561$, $\epsilon_{p_{1/2}} = 1.656$, $\epsilon_{f_{5/2}} = 2.055$, and $\epsilon_{i_{13/2}} = 2.694$. The positions of the three single-neutron states $p_{3/2}$, $h_{9/2}$, and $f_{5/2}$ are those determined in Ref. [7] by measuring γ rays in coincidence with delayed neutrons following the decay of ¹³⁴In. In that work, it was also tentatively proposed the $p_{1/2}$ assignment to the state at 1.656 MeV. The energy of the $i_{13/2}$ level has been taken from Ref. [8], where it was estimated from the position of the 2.434 MeV level in 134 Sb assumed to be a 10⁺ state of $\pi g_{9/2} \nu i_{13/2}$ nature. Following the same procedure, we have calculated $\epsilon_{i_{13/2}}$ using an effective neutron-proton interaction derived from the CD-Bonn potential. Our result comes very close to that of [8].

We now present the results of our calculations. The spectrum of 134 Sn proposed in the experimental studies of Refs. [1,2] is compared with the theoretical one in Fig. 1. In Tables I and II we report the calculated excitation energies up to about 2.8 and 4.0 MeV for the positive- and negative-parity states, respectively. We hope that these predictions may provide a guidance to the interpretation of the results of future studies on 134 Sn.

From the structure of our wave functions, it turns out that the 20 states of Table I essentially arise from the configurations $(f_{7/2})^2$, $f_{7/2}p_{3/2}$, $(p_{3/2})^2$, $f_{7/2}h_{9/2}$, and $f_{7/2}p_{1/2}$. The four members of the $(f_{7/2})^2$ multiplet all lie at an excitation energy smaller than 1.05 MeV and are separated from the other states by a pronounced gap (about 500 keV), above which we predict the existence of four seniority-two and one seniority-zero states coming from the $f_{7/2}p_{3/2}$ and $(p_{3/2})^2$ configurations, respectively. Just above these states we find the lowest- and highest-spin members (1⁺ and 8^+) of the $f_{7/2}h_{9/2}$ multiplet. All the states up to 2.4 MeV excitation energy are dominated by a unique configuration, the percentage of configurations other than the dominant one reaching at most 23% in the 2^+ state at 1.501 MeV. The other members of $f_{7/2}h_{9/2}$ and $(p_{3/2})^2$ multiplets as well as the two states arising from the $f_{7/2}p_{1/2}$ configuration are predicted in the energy region above 2.4 MeV. It should be mentioned that for some of these states a significant admixture of these three configurations is present in our calculated wave functions. As regards the negative-parity states, we have reported in Table II all the 8 states arising from the $f_{7/2i_{13/2}}$ configuration. They are practically

pure, the percentage of this configuration ranging from 87 to 100%.

Coming back to the comparison with experiment, we see from Fig. 1 that the calculated level scheme is in good agreement with the proposed one, supporting the interpretation given in Refs. [1,2]. Our calculated value of the ground-state binding energy relative to 132 Sn is 5.986 ± 0.064 MeV, to be compared with the experimental one 6.365 ± 0.104 MeV [6]. Note that the error on the calculated value arises from the experimental error on the neutron separation energy of 133 Sn.

As far as the electromagnetic observables are concerned, only the $B(E2; 6^+ \rightarrow 4^+)$ is known with a measured value of 0.88 ± 0.17 W.u. [1]. To reproduce this experimental E2 transition rate an effective neutron charge of $0.70 \pm 0.06e$ is needed in our calculation. This value is significantly smaller than 1.01e, which was determined by Zhang *et al.* in [1], but comes close to the value 0.62e of Ref. [9]. This is essentially due to the fact that in [1] the 4^+ and 6^+ states are interpreted as pure $(f_{7/2})^2$ states. By making use of $e_n^{\text{eff}} = 0.70e$, we have calculated the E2 rates for transitions involving all the observed states. The values are reported in Table III.

Shell-model calculations on ¹³⁴Sn have been performed in some previous studies [2,9,10], where use was made of a two-body interaction constructed by Chou and Warburton [10] starting from the Kuo-Herling effective interaction for the ²⁰⁸Pb region [11]. A comparison between experimental and calculated spectra is shown in Refs. [2,9], but not in [10] since no experimental data where available at that time. It should be mentioned, however, that very different conclusions are given in [2] and [9] as regards the adopted effective interaction. In fact, while the comparison between theory and experiment of Ref. [2] shows a very good agreement, the calculated excitation energies reported in [9] are all too high. Note that the results of Ref. [9] coincide with those of [10], both studies making use of the same set of SP energies, which is different from that of Ref. [2]. However, we have verified that the differences in the SP energies do not completely account for the results obtained in [2]. This point has already been evidenced in both [2] and [9], but no explanation was given.

Based on the good agreement between theory and experiment obtained for ¹³⁴Sn, we have found it interesting to predict some spectroscopic properties of ¹³⁵Sn. This nucleus is expected to have a $\frac{7}{2}^{-}$ ground state [12,13] with a binding energy of 8.393 ± 0.401 MeV [14], as derived from systematic trend. No other experimental information on its spectrum is presently available. In Table IV we report the calculated excitation energies of ¹³⁵Sn up to about 1.5 MeV. Our calculations confirm the $\frac{7}{2}^{-}$ nature of the ground state and predict for the binding energy the value 8.396 ± 0.078 MeV. As regards the excited states, we find that the six members of the $f_{7/2}^3$ multiplet all lie at an excitation energy smaller than 1.0 MeV. From Table IV it can be seen that in this energy region we predict the existence of a second $\frac{3}{2}^{-}$ state (0.643 MeV) of seniority-one nature, which is dominated by the $f_{7/2}^2 p_{3/2}$ configuration. All the levels between 1.0 and 1.5 MeV arise from the $f_{7/2}^2 p_{3/2}$ configuration with only two exceptions, the $\frac{1}{2}^{-}$ and $\frac{9}{2}^{-}$ states at 1.221 and 1.331 MeV, respectively. The latter is, in fact, the seniority-one state of the $f_{7/2}^2 h_{9/2}$ configuration while the former contains almost the same percentage of the $f_{7/2}^2 p_{3/2}$ and $f_{7/2}^2 p_{1/2}$ configurations.

In Fig. 2 we show the behavior of the experimental energies [15,16] of the lowest-lying states with $J^{\pi} = \frac{3}{2}^{-}$, $\frac{5}{2}^{-}$, $\frac{7}{2}^{-}$, $\frac{9}{2}^{-}$, $\frac{11}{2}^{-}$, and $\frac{15}{2}^{-}$ in the N = 85 isotones with $52 \leq Z \leq 64$, as well as our predictions for Z = 50. From this figure we see that our calculated energies are quite consistent with the systematic trends. Note that they refer to states arising from the $f_{7/2}^3$ configuration, which, as mentioned above, we have found to be the lowest-lying ones. In the heavier isotones the lowest-lying states with $J^{\pi} = \frac{3}{2}^{-}$, $\frac{5}{2}^{-}$, $\frac{7}{2}^{-}$, $\frac{11}{2}^{-}$, and $\frac{15}{2}^{-}$ have been interpreted as having this nature, whereas the $\frac{9}{2}^{-}$ states were supposed to arise from the $f_{7/2}^2h_{9/2}$ configuration [16]. However, only for the $Z \geq 60$ isotones, which were studied by means of transfer reaction experiments, there is evidence of the nature of these states, while no information is available for the lighter isotones. In this situation, one cannot exclude the possibility that the lowest-lying $\frac{9}{2}^{-}$ state belonging to the $f_{7/2}^2h_{9/2}$ configuration and that arising from the $f_{7/2}^2h_{9/2}$ configuration lie at about 600 and 400 keV above the first one, respectively. In this connection it is worth mentioning that for 137 Te, with two protons in the 50-82 shell, we predict three $\frac{9}{2}^{-}$ states in a even smaller energy range (from about 0.6 to 1.0 MeV), the lowest-lying one containing almost the same percentage of the three-neutron configurations, respectively. Two excited states with $J^{\pi} = \frac{11}{2}^{-}$ and $\frac{15}{2}^{-}$ have been observed in 137 Te (see Fig. 2) at 0.608 and 1.141 MeV, respectively. These energies are very well reproduced by our calculations (0.597 and 1.072 MeV).

In summary, we have shown that our realistic effective interaction derived from the CD-Bonn NN potential leads to a very good description of the experimental data recently become available for ¹³⁴Sn. We are therefore confident in the predictive power of our calculations and hope that this work may stimulate further experimental efforts to gain information on the structure of Sn isotopes beyond N = 82.

ACKNOWLEDGMENTS

This work was supported in part by the Italian Ministero dell'Università e della Ricerca Scientifica e Tecnologica (MURST). NI thanks the European Social Fund for financial support.

- [1] C. T. Zhang *el al.*, Z. Phys. A **358**, 9 (1997).
- [2] A. Korgul *el al.*, Eur. Phys J. A 7, 167 (2000).
- [3] A. Covello, L. Coraggio, A. Gargano, and N. Itaco, Acta Phys. Pol. B 32, 871 (2001), and references therein.
- [4] R. Machleidt, Phys. Rev. C 63, 024001 (2001).
- [5] A. Covello, F. Andreozzi, L. Coraggio, A. Gargano, T. T. S. Kuo, and A. Porrino, Prog. Part. Nucl. Phys. 38, 165 (1997).
- [6] B. Fogelberg, K. A. Mezilev, H. Mach, V. I. Isakov, J. Slivova, Phys. Rev. Lett. 82, 1823 (1999).
- [7] P. Hoff *et al.*, Phys. Rev. Lett. **77**, 1020 (1996).
- [8] W. Urban el al., Eur. Phys J. A 5, 239 (1999).
- [9] S. Sarkar and M. Saha Sarkar, Phys. Rev. C 64, 014312 (2001).
- [10] W. T. Chou and E. K. Warburton, Phys. Rev. C 45, 1720 (1992).
- [11] G. H. Herling and T. T. S. Kuo, Nucl. Phys. A 181, 113 (1972).
- [12] A. Korgul, H. Mach, B. Fogelberg, W. Urban, W. Kurcewicz, and V. I. Isakov, Phys Rev. C 64, 021302(R) (2001).
- [13] J. Shergur et al., Nucl. Phys. A 682, 493c (2001).
- [14] G. Audi and A H. Wapstra, Nucl. Phys. A 595, 409 (1995).
- [15] W. Urban et al., Phys. Rev. C 61, 041301(R) (2000).
- [16] Data extracted using the NNDC On-Line Data Service from the ENSDF database, file revised as of December 19, 2001.

FIG. 1. Experimental and calculated spectrum of 134 Sn.

FIG. 2. Energy systematics for the levels of the $(f_{7/2})^3$ multiplet in the N = 85 isotones. All energies are relative to the $\frac{7}{2}^-$ state. Data for $Z \ge 52$ (solid symbols) are taken from [15,16] while for Z = 50 our calculated values are reported (open symbols).

J	$E({ m MeV})$
0	0.0
2	0.640
4	0.936
6	1.050
2	1.501
4	1.765
3	1.890
5	1.944
0	2.167
1	2.304
8	2.382
2	2.490
6	2.512
4	2.515
2	2.545
3	2.635
4	2.703
5	2.733
3	2.740
7	2.744

TABLE II. Calculated excitation energies for the negative-parity states of 134 Sn up to about 4.0 MeV.

J	E(MeV)
3	3.247
5	3.621
4	3.684
7	3.713
9	3.752
6	3.848
8	3.925
10	3.973

TABLE III. Calculated B(E2) (in W.u.) in ¹³⁴Sn.

$\overline{J_i^{\pi} o J_f^{\pi}}$	B(E2)
$2^+ \rightarrow 0^+$	1.72
$4^+ \rightarrow 2^+$	1.71
$6^+ \rightarrow 4^+$	0.88
$8^+ \rightarrow 6^+$	0.12

J^{π}	τ	E(MeV)
$\frac{7}{2}$	-	0.0
$\frac{5}{2}$ -	-	0.226
$\frac{\tilde{3}}{2}$ -	-	0.356
$\frac{11}{2}$ -	-	0.611
$\frac{\tilde{3}}{2}$ -	-	0.643
$\frac{\bar{9}}{2}$ -	-	0.706
$\frac{15}{2}$ -	-	0.911
$\frac{1}{9}$ -	-	1.093
$\frac{1}{2}$	-	1.192
$\frac{1}{2}$ -	-	1.221
$\frac{\tilde{3}}{2}$ -	-	1.264
$\frac{5}{2}$ -	-	1.295
$\frac{\frac{7}{7}}{2}$ -	-	1.298
$\frac{\bar{9}}{2}$ -	-	1.331
$\frac{5}{2}$ -	-	1.430

TABLE IV. Calculated excitation energies for states of $^{135}\mathrm{Sn}$ up to about 1.5 MeV.



