

The three-nucleon bound state using realistic potential models

A. Nogga¹, A. Kievsky^{2,3}, H. Kamada⁴, W. Glöckle⁵, L.E. Marcucci^{2,3}, S. Rosati^{2,3}, M. Viviani^{2,3}

¹*Department of Physics, University of Arizona, Tucson, Arizona 85721, USA*

²*Istituto Nazionale di Fisica Nucleare, Via Buonarroti 2, 56100 Pisa, Italy*

³*Dipartimento di Fisica, Università di Pisa, Via Buonarroti 2, 56100 Pisa, Italy*

⁴*Department of Physics, Faculty of Engineering,*

Kyushu Institute of Technology, Kitakyushu 804-8550, Japan

⁵*Institut für theoretische Physik II, Ruhr-Universität Bochum, D-44780 Bochum, Germany*

The bound states of ${}^3\text{H}$ and ${}^3\text{He}$ have been calculated using the Argonne v_{18} plus the Urbana three-nucleon potential. The isospin $T = 3/2$ state have been included in the calculations as well as the n - p mass difference. The ${}^3\text{H}$ - ${}^3\text{He}$ mass difference has been evaluated through the charge dependent terms explicitly included in the two-body potential. The calculations have been performed using two different methods: the solution of the Faddeev equations in momentum space and the expansion on the correlated hyperspherical harmonic basis. The results are in agreement within 0.1% and can be used as benchmark tests. Results for the CD-Bonn interaction are also presented. It is shown that the ${}^3\text{H}$ and ${}^3\text{He}$ binding energy difference can be predicted model independently.

In the last years great efforts have been made to improve the description of the nucleon-nucleon (NN) interaction. A new generation of potentials including explicitly charge independence and charge symmetry breaking (CIB,CSB) terms appeared. These interactions describe the NN scattering data below $T_{lab} = 300$ MeV with a nearly perfect $\chi^2/\text{datum} \approx 1$. The CD-Bonn [1] and Argonne v_{18} (AV18) [2] interactions also provide a neutron-neutron (nn) force, which has been adjusted to the experimental nn scattering length, whereas the Nijmegen interactions [3] are fitted only to proton-proton and proton-neutron data. Recently, the CD-Bonn potential has been updated to CD-Bonn 2000 [4]. In this paper we only present results for the AV18 and CD-Bonn 2000 interactions. Both are quite different from each other in their functional form, but their description of the NN data is almost equally accurate. Therefore a comparison of the results will give insights into the model dependence of our understanding of the three-nucleon ($3N$) bound states.

Following for example the notation of Ref. [2], all these NN potentials can be put in the general form

$$v(NN) = v^{EM}(NN) + v^\pi(NN) + v^R(NN). \quad (1)$$

The short range part $v^R(NN)$ of all of these interactions includes a certain number of parameters (around 40), which are determined by a fitting procedure to the NN scattering data and the deuteron binding energy (BE), whereas the long range part is represented by the one-pion-exchange potential $v^\pi(NN)$ and an electromagnetic part $v^{EM}(NN)$.

For AV18, $v^{EM}(pp)$ consists of the one- and two-photon Coulomb terms plus the Darwin-Foldy term, vacuum polarization and magnetic moment interactions. The $v^{EM}(np)$ interaction includes a Coulomb term due to the neutron charge distribution in addition to the magnetic moment interaction. Finally, $v^{EM}(nn)$ is given by the magnetic moment interaction only. All these terms take into account the finite size of the nucleon charge distributions. The $v^{EM}(NN)$ for CD-Bonn is much simpler:

$v^{EM}(pp)$ is given by the Coulomb force of point protons, whereas $v^{EM}(np) = v^{EM}(nn) = 0$.

As it is well known, when these interactions are used to describe the $3N$ bound state, an underbinding of about 0.5 MeV to 0.9 MeV depending on the model is obtained (see for example Ref.[5]). The local potentials lead to less binding than the non-local ones, a characteristic related to the bigger D -state probability predicted for the deuteron. Hence, it seems to be not possible to describe the $A > 2$ systems without the inclusion of three-nucleon interaction (TNI) terms in the nuclear Hamiltonian. Several TNI models have been studied in the literature mostly based on the exchange of two pions with an intermediate Δ excitation (for a recent review see Ref. [6]). These interactions include a certain number of parameters not completely determined by theory, therefore some of them can be used to reproduce, for example, the triton BE.

In the following we show BE results for ${}^3\text{H}$ and ${}^3\text{He}$. The AV18 interaction in conjunction with the Urbana IX (UIX) TNI [7] has recently been discussed by the Argonne group in a study of several bound states with $3 \leq A \leq 8$ [8]. Here we perform a detailed calculation of the $A = 3$ system including total isospin states $T = 1/2$ and $3/2$. In addition, particular attention will be given to the BE difference $D = B({}^3\text{H}) - B({}^3\text{He})$ as a test of the CSB terms present in the interaction. The experimental value of this quantity is 764 keV, from which only 85% correspond to the standard Coulomb potential [9, 10, 11]. The remaining 15% should come from other CSB terms. A previous analysis of the contributions to D has been performed a decade ago [10, 11] before the construction of the new interactions, which include for the first time the CSB terms in the fit to the NN data. Therefore a reanalysis is in order and might remove uncertainties due to an inaccurate description of the NN data.

Because the Coulomb energy scales with the BE of ${}^3\text{H}$ [9], we need $3N$ Hamiltonians, which predict this observable accurately. This can be achieved with properly adjusted TNI's. Then the calculation of D requires reliable

Hamiltonian	$ E $	B	T	$P_{S'}$	P_P	P_D	$P_{T=3/2}$
AV18 ($T = 1/2$)	—	7.618	46.714	1.295	0.066	8.510	—
AV18 ($T = 1/2, 3/2$)	—	7.624	46.727	1.293	0.066	8.510	0.0025
AV18+UIX($T = 1/2$)	—	8.474	51.262	1.055	0.135	9.301	—
AV18+UIX($T = 1/2, 3/2$)	—	8.479	51.275	1.054	0.135	9.301	0.0025
AV18 ($T = 1/2$)	7.622	7.616	46.73	1.290	0.066	8.510	—
AV18 ($T = 1/2, 3/2$)	7.621	7.621	46.73	1.291	0.066	8.510	0.0025
AV18+UIX($T = 1/2$)	8.477	8.470	51.28	1.051	0.135	9.302	—
AV18+UIX($T = 1/2, 3/2$)	8.476	8.476	51.28	1.052	0.135	9.302	0.0025

TABLE I: ^3H BE B , mean value of the kinetic energy T , S' -, P - and D -probabilities and the probability of the $T = 3/2$ state. The Pisa results are displayed in the first four rows. The last four rows show the Bochum results, in this case the modulus of the eigenvalue E is also given. All energies are given in MeV. The probabilities are given in %.

Hamiltonian	$ E $	B	T	$P_{S'}$	P_P	P_D	$P_{T=3/2}$
AV18 ($T = 1/2$)	—	6.917	45.669	1.531	0.064	8.468	—
AV18 ($T = 1/2, 3/2$)	—	6.925	45.685	1.530	0.065	8.467	0.0080
AV18+UIX($T = 1/2$)	—	7.742	50.194	1.242	0.131	9.249	—
AV18+UIX($T = 1/2, 3/2$)	—	7.750	50.211	1.242	0.132	9.248	0.0075
AV18 ($T = 1/2$)	6.936	6.915	45.70	1.515	0.065	8.465	—
AV18 ($T = 1/2, 3/2$)	6.923	6.923	45.68	1.524	0.065	8.466	0.0081
AV18+UIX($T = 1/2$)	7.759	7.738	50.23	1.229	0.132	9.248	—
AV18+UIX($T = 1/2, 3/2$)	7.746	7.746	50.21	1.235	0.132	9.248	0.0075

TABLE II: Same as Table I for ^3He .

Hamiltonian	^3H	^3He	Interaction term	D
AV18	6 keV	-6 keV	nuclear CSB	65 keV
AV18+UIX	7 keV	-7 keV	point Coulomb	677 keV
AV18	6 keV	-6 keV	full Coulomb	648 keV
AV18+UIX	7 keV	-7 keV	magnetic moment	17 keV
			n - p mass difference	14 keV
			total (theory)	744 keV
			experiment	764 keV

TABLE III: Contribution of the proton and neutron mass difference to the ^3H and ^3He BE. The Pisa results are displayed in the first two rows. The last two rows show the Bochum results.

solutions of the $3N$ Schrödinger equation including these TNI's. For this reason we performed, as a by-product of our analysis, a benchmark calculation for the bound states of ^3H and ^3He using the AV18+UIX potential model. The Bochum group solves the Faddeev equation in momentum space [5], whereas the Pisa group uses a decomposition of the wave function in pair-correlated hyperspherical basis functions [12, 13]. Both methods were used to find BE's to an accuracy of 2 keV, which means an accuracy better than 0.1%. Such a level of accuracy is nowadays routinely achieved for the $3N$ system by several methods using only NN interactions [8, 14, 15, 16]. Here we show that the same level of accuracy is obtained, when TNI terms are taken into account.

We start considering ^3H . The calculations have been done for three equal fermions using the isospin formalism. We used an averaged nucleon mass M with the value $\hbar^2/M = 41.471 \text{ MeV fm}^2$ (the contribution of the

TABLE IV: Contributions of the various terms of the interaction to the ^3H - ^3He mass difference D . The AV18+UIX potential has been used

	^3H		^3He		D
	$ E $	T	$ E $	T	
CD-Bonn 2000	8.005	37.64	7.274	36.81	0.731
CD-Bonn 2000+TM	8.482	39.39	7.732	38.54	0.750
Exp.	8.482	—	7.718	—	0.764

TABLE V: $3N$ BE's $|E|$ for CD-Bonn 2000 with and without TM-TNI compared to the experimental values. Results are shown for ^3H , ^3He and their BE difference D . Additionally, we show the kinetic energies T . All results are given in MeV

n - p mass difference will be given separately). The AV18 and AV18+UIX have been used to calculate the BE B , the mean value of the kinetic energy T as well as the

λ	$ E $	D	a_{nn}
0.9990	8.474	742	-18.75
0.9995	8.478	746	-18.86
1.0000	8.482	750	-18.97
1.0005	8.486	754	-19.08
1.0010	8.491	759	-19.19
1.0020	8.499	767	-19.42

TABLE VI: Strength factor λ for the 1S_0 nn force, resulting ^3H BE $|E|$ in MeV, the BE difference of ^3He and ^3H D in keV and nn scattering length a_{nn} in fm. The calculations are based on the CD-Bonn 2000 potential modified by the strength factor in the 1S_0 partial wave and the TM-TNI.




FIG. 1: Difference of the ^3He and ^3H BE's D dependent on the nn scattering lengths a_{nn} . The crosses are based on the calculations shown in Table VI and the solid line is a linear fit to the crosses. The dashed, the dotted and the dash-dotted lines mark pairs of D and a_{nn} , which belong to the experimental D (a_{nn} is an estimation based on the linear fit in this case), the predictions of CD-Bonn 2000 and AV18, respectively.

S' , P - and D -state probabilities. The results are given in Table I corresponding to two different calculations: *i*) for total isospin limited to $T = 1/2$ and *ii*) including also $T = 3/2$. The occupation probability $P_{T=3/2}$ of this state is given in the last column of Table I. The first four rows of the table show the Pisa group results, whereas the last four show the Bochum group results. In the later case, the BE $|E|$ is determined from the eigenvalue spectrum of the Faddeev equations. Additionally, we present the absolute value of the expectation value of the Hamiltonian B . One sees that both values agree within the numerical accuracy. In Table II the same set of results are given for ^3He . For the $T = 3/2$ calculations we find good agreement for the BE results and the wave function properties for both nuclei. The BE's are in agreement within 2 keV or 0.05 %. The deviations for the wave function properties, especially for $P_{S'}$, are slightly bigger, but remain below 0.4 %. This is below

our numerical error bounds and confirms the reliability of both methods, even in presence of a TNI. The tables also reveal a small, but appreciable, contribution of the $T = 3/2$ state to the BE. Its inclusion produces 5 keV (8 keV) more binding in ^3H (^3He). It should be noted that the $T = 1/2$ results depend on the numerical method. The truncation of the Hilbert space to $T = 1/2$ leads to average pp (nn) and np matrix elements in the isospin $t = 1$ NN channels. This averaging is performed for the potential matrix elements in case of the Pisa calculations, but for the t -matrix in case of the Bochum scheme. The small, but visible differences show that benchmarks to this accuracy require the comparison of fully charge-dependent calculations. This inconsistency also shows up in a visible deviation of $|E|$ and B for the $T = 1/2$ Faddeev calculations, because B is based on matrix elements of the potential whereas $|E|$ is based on the t -matrix.

The contribution of the n-p mass difference is visible, but sufficiently small to be treated perturbatively. Therefore we show only perturbative estimates in Table III. The positive sign in the tritium case indicates a slightly more bound system, conversely the ^3He results slightly less bound. Again we find an encouraging agreement between the Pisa and Bochum results.

Taking into account the contribution of the n-p mass difference and averaging the Pisa and Bochum results, the final values of the BE's for the AV18+UIX are: $B(^3\text{H}) = 8.485(2)$ MeV and $B(^3\text{He}) = 7.741(2)$ MeV. This is to be compared to the experimental values: $B_{exp}(^3\text{H}) = 8.482$ MeV and $B_{exp}(^3\text{He}) = 7.718$ MeV. Therefore, the AV18+UIX potential overbinds the tritium only by 3 keV, whereas the ^3He is overbound by 23 keV. This can be better analyzed looking at the predicted BE difference $D = 744$ keV, which is 20 keV smaller than the experimental value.

The calculations have been performed using the static Coulomb potential, i.e. the momentum dependence through α' has been disregarded [17], so that our results correspond to $\alpha' = \alpha$. It has been estimated that the momentum dependence of α might contribute about half of the 20 keV shortage found in the value of D [11, 18]. Therefore our results show that there is room from other contributions as for example CSB of TNI terms and a refinement of relativistic effects on D .

The contributions to D of different parts of the interaction have been studied calculating the ^3H and ^3He BE's omitting these parts and comparing to the full calculations. Note that this is not perturbative. The results for the AV18+UIX potential including isospin states $T = 1/2$ and $3/2$ states are collected in Table IV. We distinguish *i*) the nuclear CSB terms, *ii*) the point Coulomb interaction, *iii*) the complete pp and np Coulomb interaction, which includes the finite size charge distributions, the one- and two-photon terms and the Darwin-Foldy and vacuum polarization interactions, *iv*) the magnetic moment interaction, *v*) the n-p mass difference.

One sees that the electromagnetic interaction on top of point Coulomb visibly contributes to D and cannot be

neglected. This raises the interesting question, whether CD-Bonn 2000, coming without an elaborate electromagnetic force, can also describe D .

To this aim we performed $3N$ BE calculations using the CD-Bonn 2000 interaction. The results are given in Table V. Again, the NN interaction underbinds the $3N$ nuclei. Therefore we augmented the Hamiltonian by the Tucson-Melbourne (TM) TNI [19, 20, 21]. The strength of the original model has been adjusted to reproduce the experimental ${}^3\text{H}$ BE as described in [22]. It results the πNN cut-off value $\Lambda = 4.795 m_\pi$. Again ${}^3\text{He}$ is overbound. The mass difference D , of 750 keV is slightly improved compared to our result for AV18 and UIX.

In Ref. [23] it has been observed that D is only sensitive to CSB in the S -wave. Unfortunately, nn scattering is only poorly known experimentally. There is only one datum for the scattering length, which is still controversial [24, 25, 26, 27, 28, 29, 30, 31]. AV18 is adjusted to $a_{nn} = -18.82$ fm, whereas for CD-Bonn 2000 a_{nn} results in -18.97 fm. Its charge-dependence is based on theoretical predictions of the full Bonn model [4]. To pin down the origin of the difference of the predictions of both models, we modified the 1S_0 nn interaction of CD-Bonn 2000 by a factor λ and calculated the resulting nn scattering length a_{nn} , the ${}^3\text{H}$ BE and D . We found a strong linear correlation of a_{nn} and D shown in Table VI and Fig. 1. Moreover, the prediction of AV18+UIX perfectly fit into the results obtained from CD-Bonn 2000 and TM (see the dashed-dotted marks in the figure). This shows that the dependence of D on the interaction can be traced back

to different predictions for the nn scattering length. The very different treatment of electromagnetic interactions and the differences of the CSB in higher partial waves do not affect appreciable D . The deviation of the NN +TNI force prediction of D to the experimental value might be caused, as stated before, by CSB TNI terms not considered in the present description or by relativistic effects. As a consequence we cannot estimate a_{nn} from our calculations of D . However, we would like to note that the $a_{nn} \approx -16.3$ fm found in Refs. [29, 30] would worsen our description of the $3N$ BE difference significantly.

In summary we have calculated the ${}^3\text{H}$ and ${}^3\text{He}$ BE's based on modern NN interaction models including TNI terms using two different numerical methods. Our results showed the stability and reliability of both schemes. Using only NN forces the BE's are too small, calling for TNI terms. These lead by construction to the experimental ${}^3\text{H}$ BE. We found that the BE difference of ${}^3\text{H}$ and ${}^3\text{He}$ is predicted nearly model independently. We could trace back the remaining model sensitivity to the differences in the predictions for the nn scattering length. However, uncertainties arising from CSB TNI terms and relativity do not allow us to extract the scattering length from the ${}^3\text{H}$ and ${}^3\text{He}$ BE difference. The model dependence arising from the different electromagnetic parts of the interactions is extremely small.

A.N. acknowledge partial support from NSF grant# PHY0070858. Parts of the numerical calculations have been performed on the Cray T3E of the NIC in Jülich, Germany.

-
- [1] R. Machleidt, F. Sammarruca, and Y. Song, Phys. Rev. C **53**, R1483 (1996).
- [2] R.B. Wiringa, V.G.J. Stoks, and R. Schiavilla, Phys. Rev. C **51**, 38 (1995).
- [3] V.G.J. Stoks, R.A.M. Klomp, C.P.F. Terheggen, and J.J. de Swart, Phys. Rev. C **49**, 2950 (1994).
- [4] R. Machleidt, Phys. Rev. C **63**, 024001 (2001).
- [5] A. Nogga, H. Kamada, W. Glöckle, Phys. Rev. Lett. **85**, 944 (2000).
- [6] J. Carlson and R.Schiavilla, Rev. Mod. Phys. **70**, 743 (1998).
- [7] B.S. Pudliner, V.R. Pandharipande, J. Carlson, and R.B. Wiringa, Phys. Rev. Lett. **74**, 4396 (1995).
- [8] S.C. Pieper, V.R. Pandharipande, R.B. Wiringa, and J. Carlson, Phys. Rev. C **64**, 014001 (2001).
- [9] J.L. Friar, B. F. Gibson, and G.L. Payne, Phys. Rev. C **35**, 1502 (1987).
- [10] Y. Wu, S. Ishikawa, and T. Sasakawa, Phys. Rev. Lett. **64**, 1875 (1990).
- [11] Y. Wu, S. Ishikawa, and T. Sasakawa, Few-Body Systems **15**, 145 (1993).
- [12] A. Kievsky, M. Viviani, and S. Rosati, Nucl. Phys. **A551**, 241 (1993).
- [13] A. Kievsky, Nucl. Phys. **A624**, 125 (1997).
- [14] Y. Suzuki and K. Varga, *Stochastic variational approach to Quantum-Mechanical Few-Body Problems*, vol. m54 of *Lecture Notes in Physics* (Springer-Verlag, Berlin, 1998).
- [15] H. Kameyama, M. Kamimura, and Y. Fukushima, Phys. Rev. C **40**, 974 (1989).
- [16] P. Navrátil and B.R. Barrett, Phys. Rev. C **57**, 562 (1998).
- [17] G.J.M. Austen and J.J. de Swart, Phys. Rev. Lett. **50**, 2039 (1983).
- [18] S.C. Pieper, private communication.
- [19] S.A. Coon, M.D. Scadron, P.C. McNamee, B.R. Barrett, D.W.E. Blatt, and B.H.J. McKellar, Nucl. Phys. **A317**, 242 (1979).
- [20] S.A. Coon and W. Glöckle, Phys. Rev. C **23**, 1790 (1981).
- [21] S.A. Coon and H.K. Han, Few-Body Systems **30**, 131 (2001).
- [22] A. Nogga, H. Kamada, W. Glöckle, and B.R. Barrett, nucl-th/0112026.
- [23] R. Machleidt and H. Mütter, Phys. Rev. C **63**, 034005 (2001).
- [24] B. Gabioud *et al.*, Phys. Rev. Lett. **42**, 1508 (1979).
- [25] D.E. González Trotter *et al.*, Phys. Rev. Lett. **83**, 3788 (1999).
- [26] O. Schori *et al.*, Phys. Rev. C **35**, 2252 (1987).
- [27] I. Šlaus, Y. Akaishi, and H. Tanaka, Phys. Rep. **173**, 259 (1989).
- [28] C. R. Howell *et al.*, Phys. Lett. **B444**, 252 (1998).
- [29] V. Huhn, L. Wätzold, Ch. Weber, A. Siepe, W. von Witsch, H. Witała, and W. Glöckle, Phys. Rev. C **63**,

- 014003 (2000).
- [30] V. Huhn, L. Wätzold, Ch. Weber, A. Siepe, W. von Witsch, H. Witała, and W. Glöckle, Phys. Rev. Lett. **85**, 1190 (2000).
- [31] G.A. Miller, B.M.K. Nefkens, and I. Šlaus, Phys. Rep. **194**, 1 (1990).