Deform ed Boson Scheme including Conventional q-Deform ation in Time-Dependent Variational Method. IV

— The su $(2)_q$ – and su $(1;1)_q$ – Abgebras in Four K inds of Boson Operators —

Atsushi Kuriyama, ¹ Constanca Providência, ²
Joao da Providência, ² Yasuhiko Tsue ³ and Masatoshi Yamamura ¹

¹Faculty of Engineering, Kansai University, Suita 564-8680, Japan ²Departamento de Fisica, Universidade de Coimbra, P-3000 Coimbra, Portugal

³Physics Division, Faculty of Science, Kochi University, Kochi 780–8520, Japan

A bstract

Basic idea presented in Parts (I) (III) for the deform ed boson scheme is applied to the case of the su (2)—and su (1;1)—algebras for describing many-body systems consisting of four kinds of boson operators. A possible form of the coherent state given by the present authors is generalized and the su $(2)_q$ —and the su $(1;1)_q$ —algebras are obtained in the form expressed in terms of four kinds of boson operators. As an illustrative example of the application, the framework for describing thermale exts observed in two-level shell model under the pairing correlation is proposed.

x1. Introduction

The present paper is a continuation of the last three papers (I), (II) and (III) for the deform ed boson scheme. 1) In Part (I), we focused our attention on a possible generalization of the conventional boson coherent state for the trial function in the time-dependent variational m ethod. The scheme was formulated for the systems consisting of one kind of boson operator. The comparison of Part (I) with the paper by Penson and Solom on 2) may be interesting. Part (II) was devoted to presenting the deformed boson scheme for the systems consisting of two kinds of boson operators. Through a certain procedure generalized from that in (I), we obtained the deform ation of the su (2) - and su (1;1) -algebras given by Schwinger. 3) Various forms of the q-deformed algebra were derived and, of course, one of the examples is based on the most popular deformation which is characterized by the function $[x]_q$ = $(q^x - q^x) = (q - q^1)$. Furtherm ore, we obtained a transparent understanding for the treatment of a boson system interacting with an external harmonic oscillator in the fram ework of a certain pure state in the statistical sense. 4) The idea of Ref. 4) comes from the investigation on the su (1;1)-spin like behavior in the su (2)-spin system .⁵⁾ Part (III) was aim ed at form ulating the q-deformation of the su (2;1)-algebra in the case where the generators are expressed in term s of the bilinear form s for three kinds of boson operators. 6) W ith the help of this work, we were able to obtain a transparent understanding for the description of a boson system interacting with an external harm onic oscillator in terms of statistically mixed state. 7) The above works 4); 5); 6); 7) are sum marized in Ref.8). In addition to the above three papers of this series about the deformed boson scheme, we published three short notes 91;101 on some topics supplem enting (I).

Under the above-mentioned situation, one of our next important tasks may be to form ulate the su $(2)_q$ - and the su $(1;1)_q$ -algebras for the system's described in terms of four kinds of boson operators. There exist two reasons why we perform this task. As was already mentioned, in (II), we formulated the su $(2)_q$ - and the su $(1;1)_q$ -algebras in terms of two kinds of boson operators. Then, it may be interesting for formal theoretical aspects of the deformed boson scheme to investigate which parts of the framework given in (II) conserve their forms and which parts should be added newly to the forms in (II). The above is the rst point. The second is related with physical problems. We know that the damped- and amplied-oscillation in one-dimensional harmonic oscillator can be described in terms of the su (1;1)-algebra in the Schwinger boson representation. In this case, the phase space doubling is essential and, two kinds of boson operators are necessary. The details can be found in Refs.11) and 12), together with Ref.8). By extending the above idea, the present authors gave a possible method to describe therm all and dissipative properties appearing in

the su (2)-boson models consisting of one kind of boson operator. (13) The typical example of the above su (2)-boson model is the shell model in which many nucleons move in single orbit under the pairing correlation. This model is treated in terms of the Holstein-Primako boson representation of the su (2)-algebra and the pairing rotation is the object of the investigation. Furthermore, on the basis of the viewpoint slightly dierent from that in Ref.13), we made analysis for the su (2)-algebraic model such as that appearing in quantum optics. (14) We showed that the deformed boson scheme given in Ref.9) reproduces the description quite transparently. The above mentioned situation suggests us the following point: The deformed boson scheme may be helpful to the description of two-level shell model under the pairing correlation. In this model, not only the pairing rotation but also the pairing vibration is the object of the investigation and for the use of the Holstein-Primako boson representation, two kinds of boson operators are necessary. If we adopt the idea of the phase space doubling for the description of thermal properties, totally, four kinds of boson operators are necessary. The above is the second point.

The main aim of this paper is, as was already mentioned, to formulate the deformed boson scheme for treating the systems consisting of four kinds of boson operators. The fundam ental elem ent com m on to (I) (III) is to generalize the coherent state to the form appropriate for each case. In the exponential type, the coherent state is constructed in term s of som e generators belonging to the algebra under consideration, which can be called the coherent state generating operators. Then, introducing certain function characterizing the deform ation, we are able to give the generalized coherent state. With the use of these functions, the coherent state generating operators are deformed, and by calculating the commutation relations for these deformed operators, the deformed forms of the remaining generators are obtained. In Part (IV), ie., this paper, this method is developed. In Ref.15), the present authors investigated, in the frame of four kinds of boson operators, a possible form of the coherent state which is closely related to the su(2)-and the su(1;1)-algebras. This investigation is based on a generalized Schwinger representation containing the su (M + 1)and the su (N; 1)-algebra. 16 Since we are treating the case of four kinds of bosons, possible types of the deform ation increase compared with the types shown in (II). One of the possible deform ations gives us the Holstein-Primako boson representation of the su (3) algebra in a certain symmetric representation and it is powerful for the understanding of therm ale exts in two-level shellm odel under the pairing correlation. In this paper, the su (2)and the su(1;1)-algebras appear in various notations, for example, $(\hat{S}^0; \hat{S}_0)$ and $(\hat{T}^0; \hat{T}_0)$. They obey the commutation relations

$$[\hat{S}_{+}^{0}; \hat{S}^{0}] = +2\hat{S}_{0}; \qquad [\hat{S}_{0}; \hat{S}^{0}] = \hat{S}^{0}; \qquad (1.1a)$$

$$[\hat{\Gamma}_{+}^{0};\hat{\Gamma}^{0}] = 2\hat{\Gamma}_{0}; \qquad [\hat{\Gamma}_{0};\hat{\Gamma}^{0}] = \hat{\Gamma}^{0}: \qquad (1.1b)$$

The Casim ir operators are given as

$$\hat{S}^{2} = \hat{S}_{0}^{2} + (\hat{S}_{+}^{0} \hat{S}^{0} + \hat{S}^{0} \hat{S}_{+}^{0}) = 2 = \hat{S} (\hat{S} + 1);$$
 (1.2a)

$$\hat{T}^2 = \hat{T}_0^2 \qquad (\hat{T}_+^0 \hat{T}^0 + \hat{T}_-^0 \hat{T}_+^0) = 2 = \hat{T} (\hat{T} \quad 1) :$$
 (1.2b)

The next section is devoted to recapitulating the q-deformation of the su (2)—and the su (1;1)—algebras in the case of two kinds of boson operators which was presented in (II). In x3, the su (2)—and the su (1;1)—algebras in the boson representation are recapitulated for the case of four kinds of bosons shown in Ref.15). In x4, the deformation of the coherent state is discussed. The su $(2)_q$ —and the su $(1;1)_q$ —algebras are constructed in x5. In x6, an illustrative example of the application is investigated. Finally, in x7, as the concluding remarks, it is shown that the q-deformed algebra adopted in x6 is nothing but the Holstein—Primako representation of the su (3) su (3)—algebra in a certain symmetric representation. In Appendix, the explicit expression of the operator \hat{T} deformed in the relation (12b) is given.

x2. The su (2) - and the su (1;1)-algebras for the case of the systems composed of two kinds of boson operators

For convenience of later discussion, we rst recapitulate the basic idea of the deformed boson scheme for the su (2)—and the su (1;1)—algebras for the case of two kinds of bosons. This scheme was developed in (II). In this section, we consider many-body systems composed of two kinds of boson operators (\hat{a} ; \hat{a}) and (\hat{b} ; \hat{b}). In this system, two sets of the operators (\hat{S}^0 ; \hat{S}_0) and (\hat{T}^0 ; \hat{T}_0) are introduced in the form

$$\hat{S}^{0}_{+} = \hat{a} \hat{b}; \qquad \hat{S}^{0} = \hat{b} \hat{a}; \qquad \hat{S}_{0} = (1=2) \qquad (\hat{a}\hat{a} \quad \hat{b} \hat{b}); \qquad (2\cdot 1a)$$

$$\hat{T}_{+}^{0} = \hat{a} \hat{b} ; \qquad \hat{T}^{0} = \hat{b} \hat{a} ; \qquad \hat{T}_{0} = (1=2) \quad (\hat{a} \hat{a} + \hat{b} \hat{b} + 1) : \qquad (2.1b)$$

The operators \hat{S} and \hat{T} , which commute with $(\hat{S}^0; \hat{S}_0)$ and $(\hat{T}^0; \hat{T}_0)$, respectively, are dened as

$$\hat{S} = (1=2) \hat{g}(\hat{b} + \hat{a} \hat{a});$$
 (2.2a)

$$\hat{T} = (1=2) \hat{g}(\hat{b} \hat{a} \hat{a} + 1) :$$
 (2.2b)

If we intend to investigate the su (2)-algebraic aspect of the system s, our main concern is the set $(\hat{S}^0; \hat{S}_0)$. Naturally, the case of the su (1;1)-algebraic aspect is concerned with the set $(\hat{\Gamma}^0; \hat{\Gamma}_0)$.

The eigenstates of $(\hat{S} + \hat{S}_0; \hat{S})$ and $(\hat{T}_0 + \hat{T}; \hat{T})$ with the eigenvalues $(m_S; n=2)$ and $(m_T; n=2)$ are, respectively, given by

$$\underline{\mathbf{j}}_{\mathbf{m}}, \mathbf{n} = \begin{array}{cccc}
\mathbf{q} & & & & & \\
\underline{\mathbf{m}}_{\mathbf{T}}, \mathbf{n} & & & & \\
\mathbf{m}_{\mathbf{T}}, \mathbf{n} & & & \\
\mathbf{m}_{\mathbf{T}, \mathbf{n} & & \\
\mathbf{m}_{\mathbf{T}}, \mathbf{n}$$

$$\dot{n}i = \begin{array}{ccc}
\dot{p} & -1 & (\dot{b})^n \dot{p}i : & (\dot{S}^0 \dot{p}i = \dot{T}^0 \dot{p}i = 0)
\end{array}$$
(2.4)

If the eigenvalues of $(\hat{S}; \hat{S}_0)$ and $(\hat{T}; \hat{T}_0)$ are expressed as $(s; s_0)$ and $(t; t_0)$, respectively, the states (2·3a) and (2·3b) can be rewritten in the form

$$j_{n}$$
; $ni = j_{t}$; $t_{0}i$

$$= (t_{0} t)!(t_{0} + t 1)! (a)^{t_{0}} (b)^{t_{0} + t 1} j_{0}i; (2.5b)$$

$$m_S = s + s_0$$
; $n = 2s$; (2.6a)

$$m_T = t_0 t; n = 2t:$$
 (2.6b)

It should be noted that, in the state (2.3b), the eigenvalue of \hat{T} is positive. Under the superposition of the states (2.3a) and (2.3b), we are able to de ne the following wave packets:

$$\dot{\mathbf{p}}_{S}^{0} \dot{\mathbf{i}} = \begin{pmatrix} \mathbf{q} & \mathbf{q} & \mathbf{q} \\ \mathbf{0} & \mathbf{q} & \mathbf{q} \\ \mathbf{S} & \mathbf{q} & \mathbf{q} \end{pmatrix} \qquad (2.7a)$$

$$\dot{\mathbf{j}}_{\mathrm{T}}^{0} \dot{\mathbf{i}} = \begin{pmatrix} \mathbf{q} & \mathbf{m} & \mathbf{1} \\ \mathbf{0} & \mathbf{p} & \mathbf{T} & \mathbf{T}^{0} \\ \mathbf{1} & \mathbf{q} & \mathbf{T} & \mathbf{T}^{0} \end{pmatrix} \dot{\mathbf{I}} \dot{\mathbf{I}}$$

$$j i = \exp \hat{b} \quad j0i : \quad (\hat{S}^0 j i = \hat{T}^0 j i = 0)$$
 (2.8)

Here, $_{\rm S}$, $_{\rm T}$ and denote complex parameters and $_{\rm S}^{0}$ and $_{\rm T}^{0}$ are the normalization constants given by

$$_{S}^{0} = \exp (1 + j_{S} j) j^{2} ;$$
 (2.9a)

$$_{T}^{0} = (1 \quad j_{T} \hat{j})^{1} \exp (1 \quad j_{T} \hat{j})^{1} j^{2} :$$
 (2.9b)

In (II), we introduced the following operators:

$$^{\circ}_{S} = \hat{S}^{0} (\hat{N}_{b} + 1 +)^{1} ; \quad ^{\circ}_{S} = \hat{b} ;$$
 (2.10a)

$$^{\circ}_{T} = (\hat{N}_{b} + 1 +)^{1} \hat{T}^{0}; \quad ^{\circ}_{T} = [1 \ \hat{N}_{a} (\hat{N}_{b} + 1 +)^{1}] \hat{b}; \quad (2.10b)$$

denotes an in nitesimal parameter. The operators $\hat{N_a}$ and $\hat{N_b}$ mean the boson num bers and they are related with the generators in the following form:

$$\hat{N}_{a} = \hat{a} \hat{a}; \qquad \hat{N}_{b} = \hat{b} \hat{b};$$
 (2.11)

$$\hat{N}_{a} = \hat{S} + \hat{S}_{0}; \qquad \hat{N}_{b} = \hat{S} \quad \hat{S}_{0}; \qquad (2.12a)$$

$$\hat{N}_{a} = \hat{T}_{0} \quad \hat{T} ; \quad \hat{N}_{b} = \hat{T}_{0} + \hat{T} \quad 1 :$$
 (2.12b)

Then, the form (2.10) can be rewritten as

$$^{\circ}_{S}^{0} = (\hat{S} \quad \hat{S}_{0} +)^{1} \hat{S}^{0} ; \quad ^{\circ}_{S}^{0} = \hat{b};$$
 (2.13a)

$$^{\circ}_{T} = (\hat{T}_{0} + \hat{T} +)^{1} \hat{T}^{0}; \quad ^{\circ}_{T} = \hat{b} \quad \hat{a} (\hat{T}_{0} + \hat{T} +)^{1} \hat{T}^{0}: \quad (2.13b)$$

As was shown in (II), we have the relations

$$^{0}_{S}\dot{p}_{S}^{0}\dot{i} = _{S}[1 \quad (S \quad S_{0} +)^{1}]\dot{p}_{S}^{0}\dot{i}; \quad ^{0}_{S}\dot{p}_{S}^{0}\dot{i} = \dot{p}_{S}^{0}\dot{i}; \quad (2.14a)$$

$$^{0}_{T}\dot{y}_{T}^{0}\dot{z}_{T}^{0}\dot{z}_{T}=^{T}\dot{y}_{T}^{0}\dot{z}_{T}$$
; $^{0}_{T}\dot{y}_{T}^{0}\dot{z}_{T}=^{0}\dot{y}_{T}^{0}\dot{z}_{T}$: (2.14b)

The above relations tell us that the states \dot{y}_s^0 i and \dot{y}_T^0 i can be regarded as the su (2) – and the su (1;1)-spin coherent states, respectively.

As a possible deform ation of \dot{y}_S^0 i and \dot{y}_T^0 i, in (II), we investigated the following form :

$$\dot{p}_{T} \dot{i} = \int_{T}^{0} dt \hat{N}_{a} g_{T} \hat{N}_{b} \dot{p}_{T}^{0} \dot{i} :$$
 (2.15b)

Here, $_{\rm S}$ and $_{\rm T}$ denote the normalization constants. The deformation of $\dot{\bf j}_{\rm S}^0\dot{\bf i}$ and $\dot{\bf j}_{\rm T}^0\dot{\bf i}$ are characterized by the functions $f_{\scriptscriptstyle S}$, $g_{\scriptscriptstyle S}$, $f_{\scriptscriptstyle T}$ and $g_{\scriptscriptstyle T}$ and the details of these functions were mentioned in (II).W ith the use of the relations (2·12a) and (2·12b), $\dot{p}_{\! ext{S}}$ i and $\dot{p}_{\! ext{T}}$ i can be expressed as

$$\dot{p}_{S} i = \frac{q}{s} = \frac{1}{s} = \frac{1}{s} = \frac{1}{s} \left(\hat{S} ; \hat{S}_{0} \right) \dot{p}_{S}^{0} i;$$

$$s (\hat{S}; \hat{S}_{0}) = f_{S} (\hat{S} + \hat{S}_{0}) g_{S} (\hat{S} + \hat{S}_{0})^{1};$$

$$\dot{p}_{T} i = \frac{q}{s} = \frac{1}{s} = \frac{1}{s} \left(\hat{T}; \hat{T}_{0} \right) \dot{p}_{T}^{0} i;$$

$$f_{T} (\hat{T}; \hat{T}_{0}) = f_{T} (\hat{T}_{0} + \hat{T}_{0}) g_{T} (\hat{T}_{0} + \hat{T}_{0} + \hat{T}_{0}) g_{T} (\hat{T}_{0} + \hat{T}_{0} + \hat{T}_{0}) g_{T} (\hat{T}_{0} + \hat{T}_{0} + \hat{T}_{0} + \hat{T}_{0}) g_{T} (\hat{T}_{0} + \hat{T}_{0} + \hat{T}_{0} + \hat{T}_{0} + \hat{T}_{0} + \hat{T}_{0}) g_{T} (\hat{T}_{0} + \hat{T}_{0} + \hat{T}_{$$

In the present case, $^{\circ}_{S}$, $^{\circ}_{S}$, $^{\circ}_{T}$ and $^{\circ}_{T}$ can be de ned in the form

$$^{\circ}_{S} = {}_{S} (\hat{S}; \hat{S}_{0})^{\circ 0}_{S} {}_{S} (\hat{S}; \hat{S}_{0})^{1} ; \qquad ^{\circ}_{S} = {}_{S} (\hat{S}; \hat{S}_{0})^{\circ 0}_{S} {}_{S} (\hat{S}; \hat{S}_{0})^{1} ; \qquad (2.17a)$$

$$\hat{T} = T (\hat{T}; \hat{T}_0) \hat{T} T (\hat{T}; \hat{T}_0) \hat{T} ; \qquad \hat{T} = T (\hat{T}; \hat{T}_0) \hat{T} T (\hat{T}; \hat{T}_0) \hat{T}$$
(2.17b)

The operation of $\hat{\ }_S$, $\hat{\ }_S$, $\hat{\ }_T$ and $\hat{\ }_T$ on the states j_S i and j_T i gives us the same forms as those shown in the relations (2·14a) and (2·14b), respectively. In this sense, j_S i and j_T i are still coherent states for these operators.

W ith the use of the functions f_S , g_S , f_T and g_T , we are able to obtain the q-deform ation of the su (2)—and the su (1;1)—algebras. First, we de ness and \hat{T} in the form

$$\hat{S}_{+} = {}_{S} (\hat{S}; \hat{S}_{0})^{1} \hat{S}_{+}^{0} {}_{S} (\hat{S}; \hat{S}_{0}); \qquad \hat{S} = {}_{S} (\hat{S}; \hat{S}_{0}) \hat{S}^{0} {}_{S} (\hat{S}; \hat{S}_{0})^{1}; \qquad (2.18a)$$

$$\hat{T}_{+} = {}_{T} (\hat{T}; \hat{T}_{0})^{1} \hat{T}_{+}^{0} {}_{T} (\hat{T}; \hat{T}_{0}); \qquad \hat{T} = {}_{T} (\hat{T}; \hat{T}_{0}) \hat{T}^{0} {}_{T} (\hat{T}; \hat{T}_{0})^{1} : \qquad (2.18b)$$

Then, $[2\hat{S}_0]_q$ and $[2\hat{T}_0]_q$ are given as follows:

$$\begin{split} & [2\hat{S}_{0}]_{q} = [\hat{S}_{+}; \hat{S}_{0}] \\ & = [\hat{S}_{+}; \hat{S}_{0}]_{f_{S}} [\hat{S}_{0} + 1]_{g_{S}} \quad [\hat{S}_{+}; \hat{S}_{0} + 1]_{f_{S}} [\hat{S}_{0} + \hat{S}_{0}]_{g_{S}}; \qquad (2.19a) \\ & [2\hat{T}_{0}]_{q} = [\hat{T}_{0}; \hat{T}_{+}] \\ & = [\hat{T}_{0}, \hat{T}_{0} + 1]_{f_{T}} [\hat{T}_{0} + \hat{T}_{0}]_{g_{T}} \quad [\hat{T}_{0}, \hat{T}_{0}]_{f_{T}} [\hat{T}_{0} + \hat{T}_{0} + \hat{T}_{0}]_{g_{T}}; \qquad (2.19b) \end{split}$$

Here, $[\![x]\!]_{f_{\mathbb{S}}}$, $[\![x]\!]_{g_{\mathbb{S}}}$, $[\![x]\!]_{f_{\mathbb{T}}}$ and $[\![x]\!]_{g_{\mathbb{T}}}$ are de ned as

$$[x]_{f_S} = xf_S(x)^2 f_S(x - 1)^2;$$
 $[x]_{g_S} = xg_S(x)^2 g_S(x - 1)^2;$ (2.20a)
 $[x]_{f_T} = xf_T(x)^2 f_T(x - 1)^2;$ $[x]_{g_T} = xg_T(x)^2 g_T(x - 1)^2:$ (2.20b)

The details of the above treatment are shown in (II). Furthermore, the case in which the functions f(x) and g(x) have singularities was discussed in Ref.9). In this case, the use of a certain projection operator is essential.

x3. The su (2)-and the su (1;1)-algebras in the boson representation for four kinds of boson operators

As was mentioned in x1, the aim of this paper is to apply the basic idea of the deform ed boson scheme presented in the previous section to many-body systems consisting of four kinds of boson operators. For such boson systems, we can imagine various algebras. For example, the su (4)-algebra in a symmetric boson representation may be the most popular one. In the present paper, we investigate the su (2)-and the su (1;1)-algebras. For this aim, the formalism developed by the present authors in Ref.15) is useful. From the above reason, we, rst, recapitulate the main part of Ref.15) in a form slightly different from the original one with the new notations.

We denote the four kinds of boson operators as (\hat{a} ; \hat{a}) and (\hat{b} ; \hat{b}). With the use of these operators, (\hat{S}^0 ; \hat{S}_0) and (\hat{T}^0 ; \hat{T}_0) can be expressed as

$$\hat{S}^{0}_{+} = \hat{a}_{+} \hat{b}_{+} + \hat{a} \hat{b} ; \qquad \hat{S}^{0} = \hat{b}_{+} \hat{a}_{+} + \hat{b} \hat{a} ;$$

$$\hat{S}_{0} = (1=2) \quad (\hat{a}\hat{a}_{+} \quad \hat{b}_{+} \hat{b}_{+}) + (1=2) \quad (\hat{a}\hat{a} \quad \hat{b} \hat{b}) ; \qquad (3\cdot1a)$$

$$\hat{T}^{0}_{+} = \hat{a}_{+} \hat{b} \quad \hat{a} \hat{b}_{+} ; \qquad \hat{T}^{0} = \hat{b} \hat{a}_{+} \quad \hat{b}_{+} \hat{a} ;$$

$$\hat{T}_{0} = (1=2) \quad (\hat{a}\hat{a}_{+} + \hat{b} \hat{b} + 1) + (1=2) \quad (\hat{a}\hat{a} + \hat{b}_{+} \hat{b}_{+} + 1) ; \qquad (3\cdot1b)$$

Further, we introduce new su(2)-generators $(\hat{R}^0; \hat{R}_0)$ which play an auxiliary but essential role in the present form alism :

$$\hat{R}_{+}^{0} = \hat{a}_{+} \hat{a}_{+} + \hat{b}_{+} \hat{b}_{+}; \qquad \hat{R}^{0} = \hat{a}_{+} \hat{a}_{+} + \hat{b}_{+} \hat{b}_{+};$$

$$\hat{R}_{0} = (1=2) \quad (\hat{a}_{+} \hat{a}_{+} + \hat{a}_{+} \hat{a}_{+} + \hat{b}_{+} \hat{b}_{+} \hat{b}_{}_{+} \hat{b}_{+} \hat{b}_{+} \hat{b}_{+} \hat{b}_{+} \hat{b}_{+} \hat{b}_{+} \hat{b}_$$

It should be noted that we have the relation

[any of
$$(\hat{R}^0; \hat{R}_0)$$
; any of $(\hat{S}^0; \hat{S}_0)$] = 0;
[any of $(\hat{R}^0; \hat{R}_0)$; any of $(\hat{T}^0; \hat{T}_0)$] = 0;
[any of $(\hat{S}^0; \hat{S}_0)$; any of $(\hat{T}^0; \hat{T}_0)$] = 0: (3.2)

The Casim ir operators $\hat{R}^2 = \hat{R}(\hat{R} + 1)$, $\hat{S}^2 = \hat{S}(\hat{S} + 1)$ and $\hat{T}^2 = \hat{T}(\hat{T} + 1)$ are identical to each other and, then, as a possible choice, we can set up the relation

$$\hat{R} = \hat{S} = \hat{T} \quad 1 : \tag{3.3}$$

Since the eigenvalues of \hat{R} and \hat{S} are always positive, the eigenvalue of \hat{T} is always larger than 1. In Appendix, the explicit form of \hat{T} is given in terms of $(\hat{T}^0; \hat{T}_0)$, i.e., $(\hat{a}; \hat{a})$ and $(\hat{b}; \hat{b})$.

For the above system , we consider a set of the states f $j\! m$ $_R$ m $_S$ m $_T$;nig :

$$j_{m_{R}} m_{S} m_{T}; ni = \begin{cases} q & & \\ \hline N_{m_{R}m_{S}m_{T};n} \end{cases}$$

$$q & & \\ (m_{R}!)^{1} (m_{S}!)^{1} (m_{T}!)^{1} (\hat{R}_{+}^{0})^{m_{R}} (\hat{S}_{+}^{0})^{m_{S}} (\hat{T}_{+}^{0})^{m_{T}} jni;$$

$$j_{n}i = (n!)^{1} (\hat{b})^{n} j_{0}i : (\hat{R}^{0} j_{n}i = \hat{S}^{0} j_{n}i = \hat{T}^{0} j_{n}i = 0)$$

$$(3.4)$$

Here, N $_{\text{m}_{\,\text{R}}\,\text{m}_{\,\text{S}}\,\text{m}_{\,\text{T}}\,;n}$ denotes the norm alization constant :

$$N_{m_R m_S m_T, n} = [n \vdash (n m_R)!] [n \vdash (n m_S)!] [(n + 1 + m_T) \vdash (n + 1)!] :$$
 (3.6)

The state (3.4) is a simultaneous eigenstate of the operators $\hat{R} + \hat{R_0}$, $\hat{S} + \hat{S_0}$, $\hat{T_0}$ \hat{T} and $\hat{R} = \hat{S} = \hat{T}$ 1 which are given as

the eigenvalue of
$$\stackrel{8}{\underset{\stackrel{\times}{\times}}{\times}} \hat{R} + \hat{R}_{0} \quad \text{is} \quad m_{R};$$

$$\stackrel{5}{\underset{\stackrel{\times}{\times}}{\times}} \hat{S} + \hat{S}_{0} \quad \text{is} \quad m_{S};$$

$$\stackrel{1}{\underset{\stackrel{\times}{\times}}{\times}} \hat{T}_{0} \quad \hat{T} \quad \text{is} \quad m_{T};$$

$$\stackrel{1}{\underset{\stackrel{\times}{\times}}{\times}} \hat{R} = \hat{S} = \hat{T} \quad 1 \quad \text{is} \quad n=2:$$

Therefore, fig $_R$ $_R$ $_R$ $_T$; nig form s an orthogonal and complete set. Under the superposition of the states (34), we de ne the following wave packet:

$$jc^{0}i = \begin{array}{cccc} p_{-0} & 1 & \exp & {}_{R}\vec{R}^{0}_{+} & \exp & {}_{S}\vec{S}^{0}_{+} & \exp & {}_{T}\vec{T}^{0}_{+} & j i; \end{array}$$
 (38)

$$j i = \exp(\hat{b})j0i: (\hat{R}^{0} j i = \hat{S}^{0} j i = \hat{T}^{0} j i = 0)$$
 (3.9)

Here, $_{\rm R}$, $_{\rm S}$, $_{\rm T}$ and denote complex parameters and $^{\rm 0}$ is the normalization constant given as

$$^{0} = (1 \quad j_{T} f) \exp (1 + j_{R} f) (1 + j_{S} f) (1 \quad j_{T} f)^{1} j^{2} : \qquad (3.10)$$

In the same manner as that in x2, we can introduce the following operators:

They satisfy the relations

The relations (3·12) tell us that the state j_0^0i can be regarded as a coherent state for \hat{R} , \hat{S} , \hat{S} , \hat{R} , \hat{S} and \hat{T} in the denition (311) can be expressed in terms of (\hat{R}) and \hat{R}) and \hat{R}). The operators \hat{R}^0_+ , \hat{S}^0_+ and \hat{T}^0_+ can be regarded as operators generating the coherent state.

x4. Deform at ion of the coherent state \dot{r}^0 i

We are now in a stage to give a possible deform ation of the system which was discussed in x3. First, we note the following point: As can be seen in the forms (3.1a) and (3.1b), $(\hat{S}^0; \hat{S}_0)$ and $(\hat{T}^0; \hat{T}_0)$ consist of two su(2)- and the su(1;1)-spins, $(\hat{T}^0 + \hat{J}^0; \hat{T}_0 + \hat{J}_0)$ and $(\hat{K}^0 \hat{L}^0; \hat{K}_0 + \hat{L}_0)$, respectively. The four sets $(\hat{I}^0; \hat{I}_0)$, $(\hat{J}^0; \hat{J}_0)$, $(\hat{K}^0; \hat{K}_0)$ and $(\hat{L}^0; \hat{L}_0)$ can be expressed in term s of $(\hat{a}_+; \hat{a}_+; \hat{b}_+; \hat{b}_+)$, $(\hat{a}_+; \hat{a}_+; \hat{b}_+; \hat{b}_+)$, $(\hat{a}_+; \hat{a}_+; \hat{b}_+; \hat{b}_+)$, $(\hat{a}_+; \hat{a}_+; \hat{b}_+; \hat{b}_+)$, and $(\hat{a}_+; \hat{a}_+; \hat{b}_+; \hat{b}_+)$, respectively. Each is nothing but the Schwinger representation shown in x2. Therefore, following the relations (2.2a) and (2.2b), we are able to de ne the operators \hat{I} , \hat{J} , \hat{K} and \hat{L} in the form

$$\hat{I} = (1=2) \quad \hat{B}_{+} \hat{b}_{+} + \hat{a}_{+} \hat{a}_{+}); \qquad \hat{J} = (1=2) \quad \hat{B}_{+} \hat{b}_{+} + \hat{a} \hat{a}_{+}); \qquad (4\cdot1a)$$

$$\hat{K} = (1=2)$$
 \hat{B}_1 \hat{B}_2 \hat{B}_3 \hat{B}_4 \hat{B}

The properties of the above operators are self-evident. With the use of the above operators, we can set up the eigenvalue equations:

In x2, we restricted ourselves to the positive t. However, in this section, k and lare not restricted to the positive values. Since jm $_{\rm R}$ m $_{\rm S}$ m $_{\rm T}$; ni is also the eigenstate of $(\hat{1};\hat{J};\hat{S};\hat{S}_0)$ and (k^; L^; T^; T_0), we have the following relations :

$$j_{n_{R}} m_{S} m_{T}; ni = j_{L}j_{S}; ss_{0}i_{S} = X \\ hii_{0}j_{0}j_{S}s_{0}i_{S}j_{L}; i_{0}i_{L} \quad jj_{C}j_{0}i_{J};$$

$$j_{n_{R}} m_{S} m_{T}; ni = j_{C}l_{C}; tt_{0}i_{L} = X \\ hii_{0}j_{0}j_{S}s_{0}i_{S}j_{L}; i_{0}i_{L} \quad jj_{C}j_{0}i_{J}; j_{0}i_{J};$$

$$j_{R} m_{S} m_{T}; ni = j_{C}l_{C}; tt_{0}i_{L} = X \\ hii_{0}j_{0}j_{S}s_{0}i_{S}j_{L}; i_{0}i_{L} \quad jj_{C}j_{0}i_{J}; j_{0}i_{J}; j_{0}i_{J};$$

$$k_{0}i_{L} j_{C}i_{L} j_{C}i_{L};$$

$$k_{0}i_{L} j_{C}i_{L} j_{C}i_{L};$$

$$k_{0}i_{L} j_{C}i_{L} j_{C}i_{L};$$

$$(4.3b)$$

$$j_{R} m_{S} m_{T}; ni = kl; tt_{0}i_{T} = hkk_{0}ll_{0}jt_{0}i_{T}k; k_{0}i_{K} jl; l_{0}i_{L};$$
 (4.3b)

$$m_R = i j + s;$$
 $m_T = i + j s;$ $m_S = s + s_0;$ $n = 2s;$ (4.4a)

$$m_R = t k + 1 1; m_S = t k 1; m_T = t_0 t; n = 2t 2: (4.4b)$$

Here, the quantities hii, jj jss, is and hkk, ll_0 jt, ir denote the su (2) – and the su (1;1)-C lebsh – Gordan coe cients, respectively. The relations (4.4a) and (4.4b) tellus that the state (3.4) gives us the coupling rule of the su (2) - and the su (1;1)-spins:

$$s = s_m$$
; $s_m + 1$; $s_m + 2$; ; $(i+j)$; $m \neq ji$ jj;
 $s_0 = s$; $s + 1$; ; $s_m \neq j$; $s_m \neq$

The details can be seen in Ref. 15).

Under the above preparation, let us investigate the deformation of the coherent state \dot{z}^{0} i shown in the relation (3.8) and (3.9). By expanding \dot{z}^{0} i in terms of the orthogonal set fin $_R$ m $_S$ m $_T$; nig in the relation (3.4), we have the following form :

$$\dot{y}^{0} \dot{1} = \sum_{\substack{m_{R}, m_{S}, m_{T}, n \\ C_{m_{R}, m_{S}, m_{T}, n}}}^{X} C_{m_{R}, m_{S}, m_{T}, n} \dot{y}_{m_{R}, m_{S}, m_{T}, n}; \qquad (4.6)$$

$$C_{m_{R}, m_{S}, m_{T}, n} = P \xrightarrow{0} {}_{1} {}_{1} {}_{2} \frac{q}{m_{R}, m_{S}, m_{T}, n}; \qquad (4.6)$$

$$(4.7)$$

With the use of the relations (4.3a), (4.3b), (4.4a) and (4.4b), we have two form s, which are form ally dierent from each other, for j^0 i. We denote them as j^0 si and j^0 i:

$$\mathbf{j}_{S}^{0}\mathbf{i} = \overset{X}{\text{C}}_{ij;ss_{0}}(S)\mathbf{j}i\mathbf{j};ss_{0}\mathbf{i}_{S}$$
; (4.8a)

$$\dot{j}_{S}^{0} \dot{i} = X \quad C_{ij;ss_{0}} (S) jij; ss_{0} i_{S};$$

$$\dot{j}_{T}^{0} \dot{i} = C_{kl;tt_{0}} (T) kl; tt_{0} i_{T}:$$
(4.8a)

The above two states are essentially the same as each other. However, in the case where we consider the deform ation, the form aldi erence becomes essential. The coe cients $C_{ij;ss_0}$ (S) and $C_{kl;tt_0}$ (T) denote the amplitudes of the state with the su(2)-spin (s;s₀) coupled with two su (2)-spins (i; i_0) and (j; j_0) and of the state with the su (1;1)-spin (t; t_0) coupled with the su (1;1)-spins $(k;k_0)$ and $(1;l_0)$, respectively. Therefore, if we are interested in the su (2)algebraic aspect in the present system, we should perform the deformation of j^{μ} i based on the form (4.8a). On the other hand, if in the case of the su (1;1)-algebraic aspect, we should perform the deformation based on the form (4.8b). The two amplitudes $C_{ij;ss_0}$ (S) and $C_{kl;tt_0}$ (T) depend on the (ij;ss₀) and (kl;tt₀), respectively. Then, borrowing the idea in the case developed in x2, we make the following deform ation:

$$\dot{\mathbf{j}}_{S} \mathbf{i} = \begin{pmatrix} \mathbf{q} & \mathbf{1} & \mathbf{X} \\ \mathbf{s} & \mathbf{C}_{ij;ss_{0}} (S) & \mathbf{s}_{s} (ij;ss_{0}) \mathbf{j}_{ij} \mathbf{j}_{s} \mathbf{s}_{0} \mathbf{i}_{s} \mathbf{s}_{s} \end{pmatrix} \begin{pmatrix} \mathbf{q} & \mathbf{1} & \mathbf{X} \\ \mathbf{j}_{T} \mathbf{i} \mathbf{i} \mathbf{s} & \mathbf{C}_{kl;tt_{0}} (T) & \mathbf{K}_{s} \mathbf{j}_{s} \mathbf{j}$$

Here, $_{\rm S}$ and $_{\rm T}$ denote the normalization constants. The functions $_{\rm S}$ (ij;ss₀) and $_{\rm T}$ (kl;tt₀) are de ned as follows:

$$s(ij;ss_0) = d_s(i)e_s(j)f_s(s+s_0)g_s(s-s_0)^{-1};$$
 (4.10a)

$$_{T}$$
 (kl;tt₀) = d_{T} (k 1=2) e_{T} (1 1=2) f_{T} (t₀ t) g_{T} (t₀ + t 2): (4.10b)

The quantities i and j take the values 0; 1=2;1;3=2; . However, in the case of (k; 1), the negative integers and half-integers are also perm itted. The present deform ation is characterized

by the sets of the functions $(d_S; e_S; f_S; g_S)$ and $(d_T; e_T; f_T; g_T)$. We can see that the amplitudes change from $C_{ij;ss_0}$ (S) and $C_{kl;tt_0}$ (T) to $C_{ij;ss_0}$ (S) $_S$ (ij; ss₀) and $C_{kl;tt_0}$ (T) $_T$ (kl; tt₀), respectively. The states j_S i and j_T i can be rewritten as

$$\hat{f}_{S} = f_{S} (\hat{f} \hat{J}; \hat{S} \hat{S}_{0})$$

$$= d_{S} (\hat{f}) e_{S} (\hat{J}) f_{S} (\hat{S} + \hat{S}_{0}) g_{S} (\hat{S} - \hat{S}_{0})^{-1}; \qquad (4.12a)$$

$$\hat{f}_{T} = f_{T} (\hat{K} \hat{L}; \hat{T} \hat{T}_{0})$$

$$= d_{T} (\hat{K} - 1 = 2) e_{T} (\hat{L} - 1 = 2) f_{T} (\hat{T}_{0} - \hat{T}) g_{T} (\hat{T}_{0} + \hat{T} - 2) : \qquad (4.12b)$$

Concerning the functions (4.10a) and (4.10b), the inverse should be denable and we set up the conditions shown as follows:

$$d_{X}$$
 (1=2) = d_{X} (0); e_{X} (1=2) = e_{X} (0);
$$f_{X}$$
 (1) = f_{X} (0); g_{X} (1) = g_{X} (0): (X = S; T) (4·13)

Under the above conditions, the states \dot{p}_S i and \dot{p}_T i can be expressed in the form s

In the case where the conditions (4.13) are not set up, we have the forms dierent from those shown in the relations (4.14a) and (4.14b). However, by scaling the parameters $_{\rm R}$, $_{\rm S}$, $_{\rm T}$ and appropriately, we have the same forms as those shown in the relations (4.14a) and (4.14b) with respect to the terms expressed explicitly. Therefore, we treat the functions

(4·10a) and (4·10b) as obeying the conditions (4·13). In the above treatment, we have two types of deform ations. We call the rst type represented in $j_{\mathbb{C}}$ i the S-type and the second in $j_{\mathbb{C}}$ i the T-type deform ation.

We can show that the states \dot{p}_S i and \dot{p}_T i are regarded as the coherent states for the operators $(^S_R; ^S_S; ^S_T; ^S_T; ^S)$ and $(^T_R; ^T_S; ^T_T; ^T_T)$, respectively:

O feourse, we have the relations \sin ilar to those shown in the relation (3.12) for $\binom{0}{0}$; $\binom{$

Finally, we should mention that the case where the functions characterizing the deformation have singularities can also be treated, for example, as was shown in Ref.9), with the help of certain projection operators. In this paper, we do not contact with this problem explicitly.

x5. Construction of the su $(2)_q$ -and the su $(1;1)_q$ -algebras

W ith the aid of the deform ed boson scheme presented in the previous sections, we can construct the su $(2)_q$ -and the su $(1;1)_q$ -algebras. We adopt the same principle as that adopted in (III). The operators \hat{R}^0 , \hat{S}^0 and \hat{T}^0 , the Herm ite conjugate of the generators for the coherent state generating operators \hat{R}^0_+ , \hat{S}^0_+ and \hat{T}^0_+ , are closely related with \hat{R}^0_+ , \hat{S}^0_- and \hat{T}^0_- , respectively, the deformation of which is given in the relations (4·15a) and (4·15b). For \hat{R}^0_- , \hat{S}^0_- and \hat{T}^0_- , we make the same deformation as that in the case of \hat{R}^0_+ , \hat{R}^0_- and \hat{R}^0_- :

$$\hat{K}_{+}^{S} = {}^{1}_{S} \hat{K}_{+}^{0} {}^{S}_{S}; \qquad \hat{K}^{S} = {}^{1}_{S} \hat{K}_{-}^{0} {}^{S}_{S}; \qquad \hat{S}_{+}^{S} = {}^{1}_{S} \hat{S}_{+}^{0} {}^{S}_{S}; \qquad \hat{S}_{-}^{S} = {}^{1}_{S} \hat{S}_{-}^{0} {}^{S}_{S}; \qquad \hat{S}_{-}^{S} = {}^{1}_{S} \hat{S}_{-}^{0} {}^{S}_{S}; \qquad \hat{S}_{-}^{S} = {}^{1}_{S} \hat{S}_{-}^{0} {}^{S}_{S}; \qquad \hat{S}_{-}^{S} = {}^{1}_{S} \hat{T}_{-}^{0} \hat{T}_{-}^{S}; \qquad \hat{S}_{-}^{S} = {}^{1}_{S} \hat{T}_{-}^{0} \hat{T}_{-}^{S}; \qquad \hat{S}_{-}^{S} = {}^{1}_{S} \hat{T}_{-}^{0} \hat{T}_{-}^{S}; \qquad \hat{T}_{-}^{S} = {}^{1}_{S} \hat{T}_{-}^{S}; \qquad \hat{T}_{-}^{S} \hat{T}_{-}^{S}$$

It should be noted that there are two types of the deform ations, which are speci ed by the indices S and T, respectively. Then, the deform ations of $2\hat{R_0}$, $2\hat{S_0}$ and $2\hat{T_0}$ are given as follows:

$$\begin{bmatrix}
 2\hat{K}_{0} \\
 5 \end{bmatrix} = \begin{bmatrix}
 \hat{K}_{+}^{S} \\
 7 \end{bmatrix}; \quad \begin{bmatrix}
 2\hat{S}_{0} \\
 5 \end{bmatrix} = \begin{bmatrix}
 \hat{S}_{+}^{S} \\
 7 \end{bmatrix}; \quad \begin{bmatrix}
 2\hat{T}_{0} \\
 5 \end{bmatrix} = \begin{bmatrix}
 \hat{T}_{-}^{S} \\
 7 \end{bmatrix}; \quad \begin{bmatrix}
 5 \cdot 2a
 \end{bmatrix}$$

$$\begin{bmatrix}
 2\hat{K}_{0} \\
 5 \end{bmatrix} = \begin{bmatrix}
 \hat{K}_{-}^{S} \\
 7 \end{bmatrix}; \quad \begin{bmatrix}
 2\hat{S}_{0} \\
 5 \end{bmatrix} = \begin{bmatrix}
 \hat{S}_{-}^{S} \\
 7 \end{bmatrix}; \quad \begin{bmatrix}
 2\hat{T}_{0} \\
 5 \end{bmatrix} = \begin{bmatrix}
 \hat{T}_{-}^{S} \\
 7 \end{bmatrix}; \quad \begin{bmatrix}
 5 \cdot 2b
 \end{bmatrix}$$

$$\begin{bmatrix}
 5 \cdot 2b
 \end{bmatrix}$$

In order to obtain a possible explicit expression of the form s (5·1) and (5·2), we introduce the operators $\hat{E_X}$ and $\hat{E_X}$ (X = R;S;T) de ned in the form

$$\hat{E}_{X} = \hat{X}^{0}_{+} \qquad q \frac{1}{\hat{X}^{0}\hat{X}^{0}_{+} + 1} \qquad = \qquad q \frac{1}{\hat{X}^{0}_{+}\hat{X}^{0}_{-} + 1} \qquad \hat{X}^{0}_{+};$$

$$\hat{E}_{X} = \hat{X}^{0} \qquad q \frac{1}{\hat{X}^{0}_{+}\hat{X}^{0}_{-} + 1} \qquad = \qquad q \frac{1}{\hat{X}^{0}\hat{X}^{0}_{+} + 1} \qquad \hat{X}^{0}_{-};$$

$$(X = R;S;T) \qquad (5.3)$$

Here, denotes an in nitesimal parameter. The property indispensable for the later discussion is given as

$$\hat{E}_{x}\hat{E}_{x} = 1$$
 $\hat{K}^{0}\hat{X}_{+}^{0} +)^{1}$; $\hat{E}_{x}\hat{E}_{x} = 1$ $\hat{K}_{+}^{0}\hat{X}^{0} +)^{1}$: (5.4)

Then, for the state j i obeying \hat{X}^0_+ j i = 0, we have $\hat{E_x}$ j i = 0 and for j i obeying \hat{X}^0 j i = 0, we also have $\hat{E_x}$ j i = 0. Further, for the state j i with the conditions h j i = 1 and \hat{X}^0_+ j i \(\frac{1}{2} \) 0, the state $\hat{E_x}$ j i is autom atically normalized. With the use of the operators $\hat{E_x}$ and $\hat{E_x}$, we have the following formulae:

$$\hat{X}_{+}^{0} = \hat{X}_{+}^{0} \qquad \overline{\hat{X}^{0}} \hat{X}_{+}^{0} + \qquad \overline{\hat{X}^{0}} \hat{X}_{+}^{0} + \qquad = \hat{E}_{X} \qquad \overline{\hat{X}^{0}} \hat{X}_{+}^{0} \\
= q \qquad \overline{\hat{X}_{+}^{0}} \hat{X}^{0} + \qquad \overline{\hat{X}_{+}^{0}} \hat{X}^{0} + \qquad \hat{X}_{+}^{0} = q \qquad \overline{\hat{X}_{+}^{0}} \hat{X}^{0} \qquad \hat{E}_{X} ;$$

$$\hat{X}^{0} = \hat{E}_{X} \qquad \overline{\hat{X}_{+}^{0}} \hat{X}^{0} = q \qquad \overline{\hat{X}^{0}} \hat{X}_{+}^{0} \qquad \hat{E}_{X} ; \qquad (5.5)$$

Further, the following relations are also useful for our aim:

$$\begin{split} \hat{R}^{0}_{+} \hat{R}^{0} &= (\hat{R} + \hat{R}_{0}) (\hat{R} - \hat{R}_{0} + 1) \\ &= (\hat{S} + \hat{I} - \hat{J}) (\hat{S} - \hat{I} + \hat{J} + 1) = (\hat{I} + \hat{K} - \hat{L}) (\hat{I} - \hat{K} + \hat{L} - 1); \\ \hat{R}^{0} \hat{R}^{0}_{+} &= (\hat{R} - \hat{R}_{0}) (\hat{R} + \hat{R}_{0} + 1) \\ &= (\hat{S} - \hat{I} + \hat{J}) (\hat{S} + \hat{I} - \hat{J} + 1) = (\hat{I} - \hat{K} + \hat{L}) (\hat{I} + \hat{K} - \hat{L} - 1); \\ \hat{S}^{0}_{+} \hat{S}^{0} &= (\hat{S} + \hat{S}_{0}) (\hat{S} - \hat{S}_{0} + 1) = (\hat{I} - \hat{K} - \hat{L}) (\hat{I} + \hat{K} + \hat{L} - 1); \\ \hat{S}^{0} \hat{S}^{0}_{+} &= (\hat{S} - \hat{S}_{0}) (\hat{S} + \hat{S}_{0} + 1) = (\hat{I} - \hat{K} - \hat{L}) (\hat{I} + \hat{K} + \hat{L} - 2); \\ \hat{T}^{0}_{+} \hat{T}^{0} &= (\hat{T}_{0} - \hat{T}) (\hat{T}_{0} + \hat{T} - 1) = (\hat{S} - \hat{I} - \hat{J}) (\hat{S} + \hat{I} + \hat{J} + 1); \\ \hat{T}^{0} \hat{T}^{0}_{+} &= (\hat{T}_{0} + \hat{T}) (\hat{T}_{0} - \hat{T} + 1) = (\hat{S} - \hat{I} - \hat{J}) (\hat{S} + \hat{I} + \hat{J} + 2); \end{split}$$

The use of the above relations gives us the explicit expression of the S-type deform ation shown in the relations (5.1a) and (5.2a):

$$\hat{R}_{+}^{S} = \hat{E}_{q}^{R} \frac{q}{[(\hat{S}+1=2)+(\hat{I}+1)-(\hat{J}+1=2)]_{d_{S}} e_{S}} [(\hat{S}+1=2)-(\hat{I}+1=2)+(\hat{J})]_{d_{S}} e_{S}} = \frac{(\hat{S}+1=2)+(\hat{I}+1=2)-(\hat{I}+1=2)}{[(\hat{S}+1=2)+(\hat{I}+1=2)-(\hat{I}+1=2)-(\hat{I}+1=2)-(\hat{I}+1=2)]_{d_{S}} e_{S}} \hat{E}_{R};$$

$$\hat{R}^{S} = \hat{E}_{R}^{R} \frac{q}{[(\hat{S}+1=2)+(\hat{I}+1=2)-(\hat{I}+1=2)-(\hat{I}+1=2)-(\hat{I}+1=2)+(\hat{J})]_{d_{S}} e_{S}} e_{S};$$

$$= \frac{(\hat{S}+1=2)+(\hat{I}+1=2)-(\hat{I}+1=2)}{[(\hat{S}+1=2)+(\hat{I}+1)-(\hat{I}+1=2)]_{d_{S}} e_{S}} [(\hat{S}+1=2)-(\hat{I}+1=2)+(\hat{J})]_{d_{S}} e_{S}} \hat{E}_{R};$$

$$(5.8)$$

$$\hat{S}_{+}^{S} = \hat{E}_{S} \qquad q \frac{[\hat{S} + \hat{S}_{0} + 1]_{f_{S}} [\hat{S} + \hat{S}_{0}]_{g_{S}}}{[\hat{S} + \hat{S}_{0}]_{f_{S}} [\hat{S} + \hat{S}_{0}]_{f_{S}}} = q \frac{[\hat{S} + \hat{S}_{0}]_{f_{S}} [\hat{S} + \hat{S}_{0}]_{f_{S}}}{[\hat{S} + \hat{S}_{0} + 1]_{f_{S}}} \hat{E}_{S}; \qquad (5.10)$$

$$\hat{S}^{S} = \hat{E}_{S} \qquad [\hat{S} + \hat{S}_{0}]_{f_{S}} [\hat{S} + \hat{S}_{0} + 1]_{f_{S}} = q \frac{[\hat{S} + \hat{S}_{0}]_{f_{S}} [\hat{S} + \hat{S}_{0} + 1]_{g_{S}}}{[\hat{S} + \hat{S}_{0} + 1]_{f_{S}}} \hat{E}_{S}; \qquad (5.11)$$

$$\hat{S}^{S} = \hat{E}_{S} \qquad \hat{S} + \hat{S}_{0} \hat{I}_{f_{S}} \hat{S} \qquad \hat{S}_{0} + 1 \hat{I}_{f_{S}} = \hat{S} + \hat{S}_{0} + 1 \hat{I}_{f_{S}} \hat{S} \qquad \hat{S}_{0} \hat{I}_{f_{S}} \qquad \hat{E}_{S} ; \qquad (5.11)$$

$$2\hat{S}_{0}|_{S} = [\hat{S} + \hat{S}_{0}]_{f_{S}}[\hat{S} + \hat{S}_{0} + 1]_{g_{S}} [\hat{S} + \hat{S}_{0} + 1]_{f_{S}}[\hat{S} + \hat{S}_{0}]_{g_{S}};$$
 (5·12)

$$\begin{split} & \sum_{i} \sum_{j} \sum_{i} \sum_{j} \sum_{j}$$

(5.14)

$$[2\hat{T}_0]_S = [(\hat{S}) + (\hat{I} + 1) + (\hat{J} + 1)]_{d_S e_S} [(\hat{S}) \quad (\hat{I} + 1 = 2) \quad (\hat{J} + 1 = 2)]_{d_S e_S}$$

$$+ [(\hat{S}) + (\hat{I} + 1 = 2) + (\hat{J} + 1 = 2)]_{d_S e_S} [(\hat{S}) \quad (\hat{I}) \quad (\hat{J})]_{d_S e_S} :$$

$$(5.15)$$

Here, [$l_{d_S e_S}$, [l_{f_S} and [l_{g_S} are de ned as

$$[(z) \quad p(x) \quad q(y)]_{d_S \, e_S} \, = \, (z \quad px \quad qy) d_S \, (x \quad 1 = 2)^{2p} e_S \, (y \quad 1 = 2)^{2q} \ ;$$

$$(p;q=1)$$
 (5.16)

$$[x]_{f_s} = xf_S(x)^2 f_S(x - 1)^2;$$
 (5.17)

$$[x]_{g_s} = xg_s(x)^2 g_s(x-1)^2$$
: (5.18)

For the T-type deform ation, we have the explicit form s for the relations (5.1b) and (5.2b):

$$\hat{R}_{+}^{T} = \hat{E}_{R}^{\hat{\Gamma}} \frac{q}{[(\hat{\Gamma} = 1 = 2) + (\hat{K} = 1) + (\hat{K} = 1)]_{d_{T} e_{T}}} (\hat{\Gamma} = 1 = 2) \hat{E}_{R} \frac{q}{[(\hat{\Gamma} = 1 = 2) + (\hat{K} = 1)]_{d_{T} e_{T}}}$$

$$= \frac{q}{[(\hat{\Gamma} = 1 = 2) + (\hat{K} = 1 = 2) + (\hat{K} = 1 = 2) + (\hat{K} = 1 = 2)]_{d_{T} e_{T}}} \hat{E}_{R}^{\hat{\Gamma}} ;$$

$$\begin{split} \hat{\mathbf{R}}^{\mathsf{T}} &= \hat{\mathbf{E}}_{\mathbf{R}}^{\hat{\mathsf{T}}} &= \hat{\mathbf{E}}_$$

$$\hat{S}_{+}^{T} = \hat{E}_{\hat{S}} \frac{q}{[(\hat{\Gamma}) \quad (\hat{K} \quad 1=2) \quad (\hat{L} \quad 1=2)]_{d_{T} e_{T}} [(\hat{\Gamma}) + (\hat{K} \quad 1) + (\hat{L} \quad 1)]_{d_{T} e_{T}}} = \frac{(\hat{S} \cdot 21)}{[(\hat{\Gamma}) \quad (\hat{K} \quad 1=2) \quad (\hat{L} \quad 1=2)]_{d_{T} e_{T}} [(\hat{\Gamma}) + (\hat{K} \quad 1=2) + (\hat{L} \quad 1=2)]_{d_{T} e_{T}}} \hat{E}_{\hat{S}}; \qquad (5 \cdot 22)$$

$$\hat{S}^{T} = \hat{E}_{\hat{S}} \frac{q}{[(\hat{\Gamma}) \quad (\hat{K} \quad 1=2) \quad (\hat{L} \quad 1=2)]_{d_{T} e_{T}} [(\hat{\Gamma}) + (\hat{K} \quad 1=2) + (\hat{L} \quad 1=2)]_{d_{T} e_{T}}} = \frac{(\hat{\Gamma}) \quad (\hat{K} \quad 1=2) \quad (\hat{L} \quad 1=2)]_{d_{T} e_{T}} [(\hat{\Gamma}) + (\hat{K} \quad 1) + (\hat{L} \quad 1)]_{d_{T} e_{T}}} \hat{E}_{\hat{S}}; \qquad (5 \cdot 22)$$

(5.23)

(5.24)

$$\hat{\mathbf{T}}^{T} = \hat{\mathbf{E}}_{T} - \frac{\mathbf{T}}{[\hat{\mathbf{T}}_{0} + \hat{\mathbf{T}}]_{f_{T}}[\hat{\mathbf{T}}_{0} + \hat{\mathbf{T}}]_{g_{T}}} = \frac{\mathbf{T}}{[\hat{\mathbf{T}}_{0} + \hat{\mathbf{T}}]_{f_{T}}[\hat{\mathbf{T}}_{0} + \hat{\mathbf{T}}]_{g_{T}}} \hat{\mathbf{E}}_{T};$$
 (5·26)

$$[2\hat{T}_0]_{\Gamma} = [\hat{\Gamma}_0 \quad \hat{T} + 1]_{f_{\Gamma}} [\hat{T}_0 + \hat{T}]_{g_{\Gamma}} \quad [\hat{T}_0 \quad \hat{T}]_{f_{\Gamma}} [\hat{T}_0 + \hat{T} \quad 1]_{g_{\Gamma}};$$
 (5·27)

Here, [$\ensuremath{ \ensuremath{ |} \ensuremath{ \ensu$

$$[(z) \quad p(x) \quad q(y)]_{d_T e_T} = (z \quad px \quad qy)d_T (x)^{2p}e_T (y)^{2q};$$

$$(p;q=1)$$
 (5.28)

$$[x]_{f_T} = xf_T(x)^2 f_T(x - 1)^2;$$
 (5.29)

$$[x]_{g_T} = xg_T (x - 1)^2 g_T (x - 2)^2$$
: (5.30)

The above expressions are possible expressions of the su (2) - and the su (1;1)-algebras.

x6. An illustrative example of the application

As a naldiscussion in this paper, we dealwith an illustrative example of the applications of the method presented in this work. As a simple many-nucleon model, we know a shell m odelin which many-identical nucleons move in two shell-model orbits with the same degeneracy under the pairing interaction. With the use of two kinds of boson operators, this model

is described in terms of the Holstein-Primako boson representation for the su (2)-algebra and we are able to observe collective motions such as the pairing rotation and vibration. On the other hand, the dam ped-and-amplied oscillation can be treated in the framework of the su (1;1)-algebra in the Schwinger boson representation. In this treatment, the extra boson operator is added to the formalism and it plays a role of the phase space doubling. The above consideration suggests us that an idea for describing thermal and dissipative properties of the collective motion observed in the above mentioned shell model can be formulated by the use of four kinds of boson operators based on the framework developed in this paper.

First, following the idea of Ref.11),12),13), let us derive the H am iltonian in which the phase space doubling is performed form any-body system with two kinds of bosons $(\hat{B_+}; \hat{B_+})$ and $(\hat{B_-}; \hat{B_-})$. We start from the following H am iltonian:

$$\begin{split} \hat{H_{B}^{0}} &= e_{+} \hat{B_{+}} \hat{B_{+}} + e_{+} \hat{B_{-}} \hat{B_{-}} &= g_{+} \hat{B_{+}} + \hat{B_{-}} \hat{B_{-}} + e_{+} \hat{B_{-}} \hat{B_{-}} + \hat{B_{-}} \hat{B_{-}} + \hat{B_{-}} \hat{B_{-}} \hat{B_{-}} + \hat{B_{-}} \hat{B_{-}} \hat{B_{-}} \hat{B_{-}} + \hat{B_{-}} \hat{B_{$$

Here, e , g and Z denote real parameters characterizing the system under investigation and, especially, Z is later treated as su ciently large. Our basic idea is the same as that adopted in the su (1;1)-boson m odel investigated by V itiello et. al., $^{11)}$ and independently by the present authors. For the present aim , we introduce the boson operators $(\hat{A_+};\hat{A_+})$ and $(\hat{A}^-;\hat{A}^-)$ for the phase space doubling. The Ham iltonian for this boson system is obtained by replacing $(\hat{B}^-;\hat{B}^-)$ with $(\hat{A}^-;\hat{A}^-)$ for the Ham iltonian (6·1a):

$$\begin{split} \hat{H_{A}}^{0} &= e_{+} \hat{A_{+}} \hat{A_{+}} + e \hat{A} \hat{A} & g(\hat{A_{+}} + \hat{A})(\hat{A_{+}} + \hat{A}) \\ &+ [g (e_{+} e) = 2] = Z \hat{A_{+}}^{2} \hat{A_{+}}^{2} + [g + (e_{+} e) = 2] = Z \hat{A}^{2} \hat{A}^{2} \\ &+ (g = Z) \hat{A_{+}} \hat{A} \hat{A} \hat{A_{+}} : \end{split}$$
 (6.1b)

Further, \hat{H}_{1}^{0} is introduced for the interaction between both kinds of bosons :

$$\begin{split} \hat{H}_{I}^{0} &= i \left[(\hat{A}_{+} \hat{B}_{+} + \hat{A} \hat{B}_{-}) \quad (\hat{B}_{+} \hat{A}_{+} + \hat{B} \hat{A}_{-}) \right] \\ &+ (e_{+} \quad e_{-}) = \mathbb{Z} \quad \hat{A}_{+} \hat{B}_{+} \hat{A}_{-} \quad \hat{A}_{+} \hat{B}_{-} \hat{B}_{-} \hat{A}_{+}) : \end{split}$$
 (6·1c)

Then, the Ham iltonian in the phase space doubling, \hat{H}^0 , is expressed in the form

$$\hat{H}^{0} = \hat{H}_{R}^{0} + \hat{H}_{\Delta}^{0} + \hat{H}_{T}^{0} :$$
 (6.2)

The Hamiltonian (6.2) can be rewritten as

$$\begin{split} \hat{H}^{0} &= (e_{+} + e_{-}) = 2 \quad \hat{\mathbf{g}}_{0} + (e_{+} \quad e_{-}) = 2 \quad \hat{\mathbf{g}}_{0} \left[\mathbf{1} + (2\hat{\mathbf{T}}_{0} \quad 3) = \mathbf{Z} \right] \\ &+ g \quad \hat{\mathbf{g}}_{0} \left[\mathbf{1} \quad (2\hat{\mathbf{T}}_{0} \quad 3) = \mathbf{Z} \right] \quad g(\hat{\mathbf{g}}_{+}^{0} + \hat{\mathbf{g}}^{0}) + i \cdot (\hat{\mathbf{T}}_{+}^{0} \quad \hat{\mathbf{T}}^{0}) : \end{split}$$
 (6.3)

The above expression is obtained by reading $(\hat{B} : \hat{A})$ as

$$\hat{B}_{+} ! \hat{b}_{+} ; \hat{B} ! \hat{b} ; \hat{A}_{+} ! \hat{a} ; \hat{A} ! \hat{a}_{+} :$$
 (64)

Then, the deform ed H am iltonian \hat{H}^X (X = S;T) is written down in the form

$$\begin{split} \hat{H}^{X} &= (e_{+} + e_{-}) = 2 \quad [\hat{g}_{0}]_{k} + (e_{+} \quad e_{-}) = 2 \quad [\hat{g}_{0}]_{k} [1 + ([\hat{g}\hat{T}_{0}]_{k} \quad 3) = Z] \\ &+ g \quad [\hat{g}_{0}]_{k} [1 \quad ([\hat{g}\hat{T}_{0}]_{k} \quad 3) = Z] \quad g \quad \hat{R}^{X}_{+} + \hat{R}^{X}) + i \quad \hat{T}^{X}_{+} \quad \hat{T}^{X}) : \\ &(X = S; T) \end{split} \tag{6.5}$$

In this paper, we investigate the S-type deform ation concretely under the following condition:

$$f_{S}(\hat{S} + \hat{S}_{0}) = g_{S}(\hat{S} + \hat{S}_{0}) = 1;$$
 $d_{S}(\hat{I} + 1 = 2)^{1} d_{S}(\hat{I}) = 1 + 2\hat{I} = 2;$
 $e_{S}(\hat{J})^{1} e_{S}(\hat{I} + 1 = 2) = 1 + 2\hat{I} = 2;$
(6.6)

We have already investigated the e ects of f_S , g_S , f_T and g_T in (II). In the present paper, the functions d_S , e_S , d_T and e_T are new ly added. Therefore, in order to investigate the e ects coming from d_S , e_S , d_T and e_T , we adopt the condition (6.6). Then, \hat{H}^X for X = S given in the relation (6.5) can be expressed as

$$\hat{H}^{S} = \hat{H}_{B}^{S} \quad \hat{H}_{A}^{S} + \hat{H}_{I}^{S} + \hat{h}_{B}^{S} \quad \hat{h}_{A}^{S} :$$
 (6.7)

Here, each term is given as

$$\begin{split} \hat{H}_{B}^{S} &= e_{+} \hat{B}_{+} \hat{B}_{+} + e_{-} \hat{B}_{-} \hat{B}_{-} \\ (g=Z) & \hat{B}_{+} \frac{1}{Z} \hat{A} \hat{A} \hat{B}_{+} \hat{B}_{+} + \hat{B}_{-} \frac{1}{Z} \hat{A}_{+} \hat{A}_{+} + \hat{B}_{-} \hat{B}_{-} \hat{B}_{-} \\ & \frac{1}{Z} \hat{A}_{+} \hat{A}_{+} + e_{-} \hat{A}_{-} \hat{A}_{-} \hat{A}_{-} \hat{B}_{+} \hat{B}_{+} + \hat{B}_{-} \hat{B}_{$$

$$\hat{h}_{R}^{S} = (q=Z) \hat{B}_{+} \hat{B} \hat{B} \hat{B}_{+};$$
 (6.9a)

$$\hat{h}_{A}^{S} = (g=Z) \quad \mathbf{Z}_{+}^{\hat{A}} \hat{A} \hat{A} \hat{A}_{+} :$$
 (6.9b)

The above is the Ham iltonian deformed from the Ham iltonian (6.2) for the case (6.6). It should be noted that the Ham iltonian (6.7) is derived approximately under the condition that Z is su ciently large. Later, the meaning of this approximation is mentioned.

It is now possible to investigate the relation between the above-derived Ham iltonian and that for the conventional pairing correlation in two-level shell model. For this nucleon system, we are able to write down the Ham iltonian in the form

$$H = (_{+}) + (_{-}) + 2(_{+})_{S_{0}}(+) + 2(_{-})_{S_{0}}()$$

$$G (S_{+}^{0}(+) + S_{+}^{0}(_{-})) (S_{-}^{0}(+) + S_{-}^{0}(_{-})) : (6.10)$$

Here, the indices + and denote the upper and the lower level, the degeneracies of which are the same as each other (= (2j+1)=2 : j= the spin of the j-j coupling shell model state). The parameters , and G mean the single-particle energies, the chemical potential and the strength of the pairing interaction. The sets $(S^0 (+); S_0 (+))$ and $(S^0 (-); S_0 (-))$ form the su (2)-algebras for the levels , respectively. The Holstein-Primako boson representation for $S^0_+ (-), S^0_- (-)$ and $S_0 (-)$ are given by

$$S_{+}^{0}(B) = B_{r}$$
 (B) BB;
 $S_{-}^{0}(B) = (B) BB$;
 $S_{-}^{0}(B) = (B) BB$;
 $S_{-}^{0}(B) = (B) = 2 + BB$; (6.11)

Here, (B ; B) denote boson operators. The quantities (B) play a role of the seniority numbers of the levels and they are independent of the bosons (B ; B). Then, the Ham iltonian (6.10) can be rewritten as

$$H_{B} = (_{+} _{-} _{-})(_{-} (B_{+}) + 2B_{+}B_{+}) + (_{-} _{-} _{-})(_{-} (B_{-}) + 2B_{-}B_{-}) |$$

$$G_{B_{+}} (B_{+}) B_{+}B_{+} + B_{-} (B_{-}) B_{-}B_{-} |$$

$$(B_{+}) B_{+}B_{+}B_{+} + (B_{-}) B_{-}B_{-} |$$

$$(6.12)$$

W ith the aim of the phase space doubling, we, further, introduce the operators S_+^0 (A), S_-^0 (A) and S_-^0 (A) in the form

$$S^{0}(A) = (A) A A A;$$

 $S_{0}(A) = (A) = (A) = 2 + A A;$ (6.13)

The Hamiltonian HA which is necessary for the phase space doubling is given in the form

The m eaning of the notations is self-evident. Under the correspondence (B \hat{B} ; A \hat{A}), we can see that (H $_B$ H $_A$) is reduced to (H $_B$ H $_A$) shown in the relations (6.8a) and (6.8b), if the following relations hold:

$$e_{+} = 2(_{+})$$
 (); $e = 2($) ($_{+}$); $q = G$; $Z = :$ (6.15)

In this case, the seniority numbers (B) and (A) are obtained in the form

$$(B_{+}) = A A ; (B_{-}) = A_{+}A_{+} ; (A_{+}) = B B ; (A_{-}) = B_{+}B_{+} : (6.16)$$

Then, it may be permitted to adopt the following form as the interaction part:

$$H_{I} = (i =) S_{+}^{0} (A_{+}) S_{+}^{0} (B_{+}) + S_{+}^{0} (A_{-}) S_{+}^{0} (B_{-})$$

$$S_{+}^{0} (B_{+}) S_{-}^{0} (A_{+}) S_{+}^{0} (B_{-}) S_{+}^{0} (A_{-}) : (6.17)$$

However, it should be noted that in the two-level model, there does not exist the term such as $(\hat{h}_{R}^{S} - \hat{h}_{A}^{S})$ shown in the relation (6.9).

The above is an illustrative example. For the above treatment, we give two comments. First is related to a possible introduction of the seniority numbers. For example, the form s $(6\cdot11)$, $(6\cdot13)$ and $(6\cdot16)$ tellusthat A A plays a role of the seniority numbers (B_+) for the pairing correlation described by the boson $(B_+;B_+)$. We know that if (B_+) becomes larger, the pairing correlation becomes weaker and, if A A becomes larger, the statistical mixing becomes larger. The above suggests us that if the statistical mixing becomes stronger, the pairing correlation becomes weaker. This may be acceptable as a natural feature. This is the rst comment. Second is related with H I shown in the relation $(6\cdot17)$. In the case of an example of the deformation for two kinds of bosons, we got such a term (S_+^0) (a) S_+^0 (b) S_-^0 (b) S_-^0 (c) S_-^0 (c) S_-^0 (c) S_-^0 (d). The form S_-^0 (f) is its natural generalization. Therefore, it may be interesting to investigate these points concretely, together with the term S_-^0

x7. Concluding remarks

In this paper, we proposed a possible deformed boson scheme for many-body systems consisting of four kinds of boson operators. The basic viewpoint was to focus the interest on the su (2)- and the su (1;1)-algebras. As an example of the application, we investigated two-level shell model under the pairing correlation in the framework of the deformed boson scheme under a special condition. In the present section, as the concluding remarks, we discuss the scheme given in x6 in the language of the su (3) su (3)-algebra in a symmetric representation presented in Ref.16). For this aim, we prepare three kinds of boson operators $\binom{2}{3}$, $\binom{2}{3}$, and $\binom{2}{3}$ and define the following bilinear form:

$$\hat{S}_{+}^{0} = \hat{S}_{+}^{0} = \hat{S}_{0}^{0} = \hat{S}$$

$$^{\circ}_{+} = ^{\circ}_{+} ^{\circ}_{+} ^{\circ}_{+} ^{\circ}_{-} = ^{\circ}_{+} ^{\circ}_{-} ^{\circ}_{-} = (^{\circ}_{-} ^{\circ}_{-} ^{\circ}_{-}) = 2;$$
 (7.1b)

$$^{\circ}_{+}^{\circ} = ^{\circ}_{+}^{\circ}_{+}^{\circ}_{+}^{\circ}_{+}^{\circ}_{+}^{\circ}_{-}^{\circ}_{+}^{\circ}_{+}^{\circ}_{-}^{\circ}_{-}^{\circ}_{+}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_{-}^{\circ}_$$

The sets $(\hat{S}^0; \hat{S}_0)$, $(\hat{S}^0; \hat{S}_0)$, $(\hat{S}^0; \hat{S}_0)$, and $(\hat{S}^0; \hat{S}_0)$ form the su(2)-algebras. However, since $\hat{S}_0 = \hat{S}_0 = \hat{S}_0$, $(\hat{S}^0; \hat{S}_0)$, $(\hat{S}^0; \hat{S}_0)$, and $(\hat{S}^0; \hat{S}_0)$ are independent of each other. These operators from the su(3)-algebra in the symmetric representation and we can call the set the Schwinger boson representation.

We can prove that all the generators commute with the following operator:

Then, the Holstein-Primako boson representation for a xed total boson number can be constructed and the result is as follows:

$$S_{+}^{0} = 2K$$
; $S_{-}^{0} = 2K$; S_{-}^{0}

Here, (;) and (;) denote the boson operators. The quantity K is the eigenvalue of \hat{K} de ned in the relation (72) and K = 0; 1=2; 1; 3=2; . Judging from the form of the relation (73), we can call the set (73) as the Holstein-Primako boson representation for the su (3)-algebra in the symmetric representation.

On the basis of the relations (7.3a) and (7.3b), let us reinvestigate the form s (6.11) and (6.13). In this case, the relation (6.16) is also noted. In associating with the form s (6.11) and (6.13), we do not the operator

$$_{+}^{0}$$
 (+) = B₊A ; $_{-}^{0}$ (+) = A B₊ ; $_{-}^{0}$ () = A₊B : (7-4)

Together with the form (7.4), we can see that the sets $(S^0(B_+); S_0(B_+); S^0(A_-); S_0(A_-); S_0(A_-); S_0(A_+); S_0(A_$

Appendix A

— The proof of the explicit expression of the operator \hat{T} —

First, let us sketch the eigenvalue problem for the su(1;1)-algebra in the systems described in terms of two and four kinds of boson operators. We presuppose the existence of a state jti, which obeys

$$\hat{T}^0$$
 ji = 0; \hat{T}^2 ji = t(t 1) ji; \hat{T}_0 ji = ji: (A·1)

Here, is a function of t, the explicit form of which should be determined in the fram ework of the treatment. In this Appendix, there appear various states in which we om it the normalization constants. Further, for a moment, we om it the extra-quantum numbers additional to t. Since the Casim ir operator can be expressed in the form $\hat{T}^2 = \hat{T}_0 (\hat{T}_0 - 1) - \hat{T}_+^0 \hat{T}^0$, the relation (A·1) gives us

$$(1) = t(t 1) :$$
 (A.2)

As a form alsolution of Eq. (A.2), we have

$$t = i$$
; (A·3)

$$t = 1$$
 : (A-4)

The above means that can be expressed in terms oft.

For the case of two kinds of boson operators, the relation (2.1b) gives us

$$= 1=2 + n=2 : (n = 0;1;2;)$$
 •5(A)

From the relations (A \cdot 3) (A \cdot 5), the following forms are derived:

$$t = 1 = 2 + n = 2$$
; $(t = 1 = 2; 1; 3 = 2;$)

$$t = 1 = 2$$
 $n = 2$; $(t = 1 = 2; 0; 1 = 2;)$

The case n=0 corresponds to the boson vacuum j0i and two expressions (A ·6) and (A ·7) coincide with each other (t = 1=2). Then, t=1=2i can be expressed as

$$t = 1 = 2i = 70i$$
: (A \cdot 8)

In the case n=1;2;3; , we make the relations A and A A or A correspond to the following states, respectively:

$$t = 1 = 2 + n = 2i = (n!)^{1} (6)^{n} (2)^{i}$$
; (A.9)

$$t = 1 = 2$$
 $n = 2i = (n!)^{1} (a)^{n} t$ (A ·10)

The relation (A·6) and (A·7) can be uni ed into the form $n=2=\pm 1=2$ jand, with the aid of the relation (A·5), we have

$$= 1=2+ t 1=2i$$
: (t= 1=2; 1; 3=2;) •1(A)

Successive operations of \hat{T}^0_+ on the state ji gives us the eigenstates of \hat{T}^2_- and $\hat{T_0}_-$ in the form

$$t_{i}t_{0}i = (\hat{T}_{+}^{0})^{t_{0}} \stackrel{(1=2+t_{1}=2)}{\longrightarrow} t_{i}; \tag{A.12}$$

$$t_0 = 1 = 2 +$$
 $t_1 = 2$; $3 = 2 +$ $t_2 = 1 = 2$; $5 = 2 +$ $t_3 = 1 = 2$; $t_4 = 1 = 2$; $t_5 = 2 +$ $t_7 = 1 = 2$; t

Of course, t_0 denotes the eigenvalue of $\hat{T_0}$. The relations (A·6), (A·7), (A·9) and (A·10) suggest us

$$\hat{T}^2 = \hat{T} (\hat{T} - 1); \qquad \hat{T} = 1 = 2 + (\hat{N}_b - \hat{N}_a) = 2:$$
 (A·14)

The expression (A \cdot 14) is nothing but the form introduced in \times 2.

For the case of four kinds of boson operators, we have

$$= 1 + n=2 : (n = 0;1;2;)$$
 1(3)

Judging from the expression (3.1b), the above form may be accepted as self-evident. From the relations (A.3), (A.4) and (A.15), the following forms are derived:

$$t = 1 + n = 2;$$
 $(t = 1; 3 = 2; 2;$)

$$t = n=2;$$
 $(t = 0; 1=2; 1;)$

The case n=0 corresponds to the boson vacuum $\mathcal{D}i$ and two expressions (A·16) and (A·17) do not coincide with each other:

This situation is quite dierent from the case (A-8). This case is attributed to the manner how to specify the vacuum $\mathfrak P$ i in the present quantum number. There exist two manners: the form (A-18) or (A-19). In the present, we adopt the form (A-18). Therefore, as the continuation from the form (A-18), it may be natural to adopt the form (A-16) for n = 1; 2; 3;:

$$j = 1 + n = 2i = (\hat{b})^n \mathcal{D}i (= jni)$$
: (A ·20)

However, in the present case, it is noted that the state which satis es the relation (A·1) is not restricted to the form (A·20). By successive operations of \hat{S}_{+}^{0} and \hat{R}_{+}^{0} on the state t = 1 + n = 2i, we have

In the above, m $_{\rm S}$ and m $_{\rm R}$ denote the extra-quantum numbers additional to t. The state jm $_{\rm S}$ m $_{\rm R}$: ti is the eigenstate of \hat{S}_0 and \hat{R}_0 with the eigenvalues m $_{\rm S}$ + n=2 and m $_{\rm R}$ + n=2, respectively. By successive operation of the operator \hat{T}_+^0 on the state (A 21), we have the eigenstate of \hat{T}_-^2 and \hat{T}_0 :

$$jm_S m_R : t; t_0 i = (\hat{\Gamma}_+^0)^{t_0 t} jm_S m_R : ti;$$
 (A ·22)

$$t_0 = t; t + 1; t + 2;$$
 :

Hereafter, concerning the quantum numbers, we discuss only t and t_0 and, then, we om it the quantum numbers m $_{\rm S}$ and m $_{\rm R}$. The state ${\tt j}$; t_0 i satis es the relations

$$(\hat{\Gamma}^{0})^{r} \mathbf{j}; t_{0} \mathbf{i} = \begin{array}{c} q \\ \hline (t_{0} \ t)!(t_{0} + t \ 1)! \end{array} \qquad q \\ \hline (t_{0} \ t \ r)!(t_{0} + t \ r \ 1)! \end{array} \qquad \begin{array}{c} \mathbf{j}; t_{0} \ r\mathbf{i}; \\ \hline (A \cdot 24) \\ \hline (\hat{\Gamma}^{0}_{+})^{r} \mathbf{j}; t_{0} \mathbf{i} = \end{array} \qquad \begin{array}{c} q \\ \hline (t_{0} \ t)!(t_{0} + t \ 1)! \end{array} \qquad \begin{array}{c} \mathbf{q} \\ \hline (t_{0} \ t + r)!(t_{0} + t + r \ 1)! \mathbf{j}; t_{0} + r\mathbf{i}: \\ \hline (A \cdot 25) \end{array}$$

In the case of two kinds of bosons, we have the operator \hat{T} shown in Eq.(A·14). The aim of this Appendix is to give the explicit form of \hat{T} for the case of four kinds of boson operators.

If the Casim ir operator of the su (1;1)-algebra, \hat{T}^2 , can be expressed in the form $\hat{T}^2 = \hat{T}(\hat{T} + 1)$, \hat{T} satis es

$$\hat{T}$$
 j; t_0 **i** = t **j**; t_0 **i** : (A ·26)

In order to search the explicit form of \hat{T} itself, we introduce the following operator;

$$\hat{g}_{r_0=1} = g_{r_0} (\hat{T}_+^0)^{r_0} (\hat{Z}_{r_0}^0)^{-1} (\hat{T}^0)^{r_0}; \qquad (A \cdot 27)$$

$$\hat{Z}_{r_0} = (2\hat{T}_0)(2\hat{T}_0 + 1)$$
 $\hat{T}_0(2 - 2r_0 - 2)$: (A -28)

Since \hat{T}_0 is positive-de nite and $\hat{\mathcal{L}}_{r_0}$) ¹ is sandwiched between $(\hat{T}_+^0)^{r_0}$ and $(\hat{T}^0)^{r_0}$, the operator \hat{T}_0 can be de ned. The operator \hat{T}_0 is given as

$$D_n = (2n) = (n!)^2$$
: (A ·30)

The coe cients g_{r_0} from $r_0 = 1$ to $r_0 = 10$ are as follows:

$$g_1 = 1$$
; $g_2 = 1$; $g_3 = 2$; $g_4 = 5$; $g_5 = 14$; $g_6 = 42$; $g_7 = 132$; $g_8 = 429$; $g_9 = 1430$; $g_{10} = 4758$: (A·31)

W ith the use of the relations (A 4), (A \cdot 11), (A \cdot 12) and (A \cdot 16) with (A \cdot 17), we can show the following relation:

$$_{0}$$
 t; t_{0} i = $(t_{0}$ t) t; t_{0} i: (A ·32)

The proof is easy. We can prove that, for arbitrary g_r , the state $t;t_0i$ is an eigenstate of \hat{g}_r with the eigenvalue g_r , which is given as

We intend to determ in g_r so as to satisfy the relation

$$_{0} = t_{0} t : (A \cdot 34)$$

The relation (A $\cdot 34$) with (A $\cdot 33$) can be rewritten as

$$g_s r!((2t 1) + 2 (r s))!((2t 1) + r)!$$
 $s=1$

$$[(r s)!((2t 1) + (2r 1))!((2t 1) + (r s))!]^1 = r; \qquad (A \cdot 35)$$
 $t_0 = t + r: \qquad (A \cdot 36)$

The relation (A \cdot 35) with r = 1 gives us

$$g_1 = 1$$
: (A ·37)

The relation (A ·35) can be rewritten as

$$g_{r} = r((2t 1) + 2r)![((2t 1) + 2r)r!((2t 1) + r)!]^{1}$$

$$g_{s}((2t 1) + 2(r s))![(r s)!((2t 1) + (r s))!]^{1} : (A \cdot 38)$$

$$s=1$$

W ith the use of the relation (A \cdot 38), we can determ in e g_r from r=2 to the higher successively. Of course, the relation (A \cdot 37) is used. Then, we can show that g_r does not depend on t for any value of r. Thus, by putting t=1=2 for the form (A \cdot 38), the following recursion formula is derived:

$$g_r = D_r$$
 $g_s D_{r s}$: (A ·39)

Here, D_n is defined in the relation (A·30). The solution of the recursion formula (A·39) is given in the form (A·29). The relation (A·32) is rewritten as

$$\hat{T}_0$$
; \hat{T}_0 ; \hat{T}

Then, we have

$$\hat{T} = \hat{T_0} \quad \hat{}_0 :$$
 (A -41)

Of course, $\hat{0}$ is given in the relation (A·27) with (A·28). It is interesting to see that the operator \hat{T} can be expressed in terms of the generators \hat{T}^0 and \hat{T}_0 .

R eferences

- [1] A. Kuriyama, C. Providência, J. da Providência, Y. Tsue and M. Yamamura, Prog. Theor. Phys. 106 (2001), 751, 765. ibid. 107 (2002), 65.
- [2] K.A.Penson and A.I.Solom on, J.M ath. Phys. 40 (1999), 2354.
- [3] J. Schwinger, in Quantum Theory of Angular Momentum, ed. L. Biedenham and H. Van Dam (Academic Press, New York, 1955), p. 229.
- [4] A.Kuriyam a, J.da Providência, Y.Tsue and M.Yamamura, Prog. Theor. Phys. 98 (1997), 893.
- [5] A.Kuriyam a, J.da Providência, Y.Tsue and M.Yamamura, Prog. Theor. Phys. 98 (1997), 381.
- [6] A. Kuriyam a, J. da Providência, Y. Tsue and M. Yamamura, Prog. Theor. Phys. 100 (1998), 993.

- [7] A. Kuriyam a, J. da Providência, Y. Tsue and M. Yamamura, Prog. Theor. Phys. 100 (1998), 1181.
- [8] A.Kuriyam a, J.da Providência, Y.Tsue and M.Yamamura, Prog. Theor. Phys. Suppl. No. 141 (2001), 113.
- [9] A. Kuriyama, C. Providência, J. da Providência, Y. Tsue and M. Yamamura, Prog. Theor. Phys. 107 (2002), 443.
- [10] A. Kuriyama, C. Providência, J. da Providência, Y. Tsue and M. Yamamura, to appear in Prog. Theor. Phys. 107 (2002), No.6.
- [11] E. Celeghini, M. Rasetti, M. Tarlini and G. Vitiello, Mod. Phys. Lett. B 3 (1989), 1213.
 - E.Celeghini, M.Rasetti and G.Vitiello, Ann. of Phys. 215 (1992), 156.
- [12] Y. Tsue, A. Kuriyam a and M. Yamamura, Prog. Theor. Phys. 91 (1994), 49, 469.
- [13] Y. Tsue, A. Kuriyam a and M. Yamamura, Prog. Theor. Phys. 93 (1995), 541, 1037.
- [14] H. Akaike, Y. Tsue, C. Providência, J. da Providência, A. Kuriyam a and M. Yama-mura, Prog. Theor. Phys. 105 (2001), 717.
- [15] A.Kuriyama, J.da Providência and M.Yamamura, Prog. Theor. Phys. 103 (2000), 305.
- [16] A.Kuriyam a, J.da Providência and M.Yam amura, Prog. Theor. Phys. 103 (2000), 285.