

Estimation of admixture of twelve quark bag state in ${}^4\text{He}$ nucleus

A.M. Mosallem ¹, V.V.Uzhinskii

Joint Institute for Nuclear Research,
Laboratory of Information Technologies

The $p{}^4\text{He}$ elastic scattering at the energy range from 0.695 to 393 GeV is analyzed in the framework of the Glauber theory. The Glauber amplitudes were evaluated using isospin-averaged nucleon-nucleon amplitudes and the ${}^4\text{He}$ wave function as a superposition of the Gaussian functions. The values of the calculated differential cross sections usually exceed the experimental ones. In order to overcome the discrepancy, it is assumed following to the paper by L.G. Dakno and N.N. Nikolaev (Nucl. Phys. **A436** (1985) 653) that the ground state wave function of ${}^4\text{He}$ has an admixture of a twelve quark bag. Neglecting all transition amplitudes, the proton - 12q bag scattering amplitude was chosen in a simple gaussian form. The inclusion of the 12q bag leads to decreasing the $p{}^4\text{He}$ differential cross section and to a shift of the dip position to a large values of t what is needed for a successful description of the experimental data. While fitting the data it is found that the weight of the 12q bag state in the ground state of the ${}^4\text{He}$ nucleus is $\sim 10.5\%$, $\sigma_{p-12q}^{tot} \sim 34$ mb, and the slope parameter of the $p - 12q$ bag elastic scattering is ~ 23 $(\text{GeV}/c)^{-2}$. Inelastic shadowing is not taken into account at the calculations.

arXiv:nucl-th/0205039v1 14 May 2002

¹Math. & Theor. Phys. Dept., NRC, AEA, Cairo, Egypt.

Introduction

The study of structure of light nuclei such as ${}^4\text{He}$, ${}^6\text{He}$, ${}^{11}\text{Li}$ and so on is very popular now. There is a big progress in understanding the structure of light exotic nuclei ${}^6\text{He}$, ${}^{11}\text{Li}$, The deuteron structure is a subject of continuous discussions. Only few hypotheses about the structure of ${}^4\text{He}$ exist now. In paper by L.G. Dakno and N.N. Nikolaev [1] it was assumed and shown that 12% admixture of twelve quark bag configuration in the ground state wave function of the ${}^4\text{He}$ nucleus allows one to understand the irregularities of proton- ${}^4\text{He}$ elastic scattering at high energies. We believe that the hypothesis will permit to describe other reactions – $d + {}^4\text{He}$, ${}^4\text{He} + {}^4\text{He}$, ${}^4\text{He} + C$, ... etc. The matter is the carbon and oxygen nuclei are considered as strong clustering nuclei, consisting of α particles. So, the peculiarities of the ${}^4\text{He}$ nucleus can manifest themselves in the structure of ${}^{12}\text{C}$ and ${}^{16}\text{O}$ nuclei.

To show these, one needs to calculate elastic and inelastic scattering of ${}^4\text{He}$ on different nuclei. The Glauber diffraction theory [2] of multiple scattering processes has been generally accepted as a suitable framework for such calculations. But it was recognized many years ago that the model predictions have been far from being perfect even for the hadron-nucleus scattering process.

Many authors believe that it is due to inelastic screening, and many attempts have been made to take them into account [3, 4, 5, 6, 7, 8]. According to different calculations, the inelastic screening corrections to the total hadron-nucleus cross sections are at the level of 2–5 %. It is not enough to describe the $p^4\text{He}$ scattering. Inclusion of the corrections into calculations of the $p^4\text{He}$ elastic scattering leads to a shift of the first diffraction minimum to low values of the momentum transfer, t . But a good description of the cross sections demands shift of the minimum to large values of t . In order to solve the problem at the first step of our study, we will omit the corrections.

The content of the paper is as follows: Sec. 1 describes calculation of the ${}^4\text{He}$ form factor with different parametrizations of the ground state wave function. Sec. 2 gives calculations of the $p^4\text{He}$ elastic scattering amplitude and differential cross section with these parametrizations. In Sec. 3 we include the twelve quark bag admixture and fit the parameters of the twelve quark bag using experimental data. In the last Sec. we summarize our results.

1 Form-factor of ${}^4\text{He}$

The main characteristic properly of a nucleus is a nuclear form-factor.

$$F(\vec{q}) = \int e^{i\vec{q}\cdot\vec{r}_1} |\psi(\vec{r}_1, \dots, \vec{r}_A)|^2 \prod_{i=1}^A d^3r_i, \quad (1)$$

where ψ is the wave function of a nucleus in the ground state, A – a mass number of the nucleus, $\vec{r}_1, \vec{r}_2, \dots$ – radius vectors of nuclear nucleons, \vec{q} – momentum transfers. It is very often assumed in Glauber calculations that the square module of ψ can be represented as

$$|\psi(\vec{r}_1, \dots, \vec{r}_A)|^2 = (2\pi)^3 \rho_c \delta\left(\sum_{i=1}^A \vec{r}_i\right) \prod_{i=1}^A \varphi(\vec{r}_i). \quad (2)$$

The δ -function is introduced in order to satisfy the obvious condition

$$\left(\sum_{i=1}^A \vec{r}_i \right) = 0. \quad (3)$$

For the ${}^4\text{He}$ nucleus in the paper [1] the following parametrizations of $\varphi(\vec{r})$ were proposed:

$$\begin{aligned} (A) \quad & \varphi(\vec{r}) = \exp[-\vec{r}^2/R_1^2], \\ (B) \quad & \varphi(\vec{r}) = \exp[-\vec{r}^2/R_1^2] - D_1 \exp[-\vec{r}^2/R_2^2], \\ (C) \quad & \varphi(\vec{r}) = (\exp[-\vec{r}^2/2R_1^2] - D_1 \exp[-\vec{r}^2/2R_2^2])^2, \\ (D) \quad & \varphi(\vec{r}) = \exp[-\vec{r}^2/R_1^2] + D_1 \exp[-\vec{r}^2/R_2^2] - (1 + D_1 - D_2^2) \exp[-\vec{r}^2/R_3^2]. \end{aligned}$$

The parameters are given in Table 1.

Table 1: Values of used parameters (from [1])

	R_1^2 (GeV/c) ⁻²	R_2^2 (GeV/c) ⁻²	R_3^2 (GeV/c) ⁻²	D_1	D_2
<i>A</i>	51.01				
<i>B</i>	48.07	3.67		1.0	
<i>C</i>	47.29	1.6		1.6	
<i>D</i>	62.06	19.0	10.13	3.79	0.31

We will use a general form for the function φ as

$$\phi(\vec{r}) = \sum_{i=1}^N C_i e^{\vec{r}^2/R_i^2}. \quad (4)$$

In Eq. (2) ρ_c is the normalization constant determined from the condition

$$\int |\psi(\vec{r}_1, \dots, \vec{r}_A)|^2 \prod_{i=1}^A d^3 r_i = 1. \quad (5)$$

Substituting Eq. (2) in the normalization condition (5), we have

$$\rho_c (2\pi)^3 \int \delta \left(\sum_{i=1}^4 \vec{r}_i \right) \prod_{i=1}^4 \varphi(\vec{r}_i) d^3 r_i = 1. \quad (6)$$

Using the following representation of the δ function

$$\delta \left(\sum_{i=1}^4 \vec{r}_i \right) = \frac{1}{(2\pi)^3} \int d^3 \alpha e^{i\vec{\alpha} \cdot (\sum_{i=1}^4 \vec{r}_i)},$$

the Eq. (6) can be re-written as

$$\rho_c \int (2\pi)^3 \left(\frac{1}{2\pi} \right)^3 d^3 \alpha e^{i\vec{\alpha} \cdot (\sum_{i=1}^4 \vec{r}_i)} \prod_{i=1}^4 \varphi(\vec{r}_i) d^3 r_i = 1 \quad (7)$$

Then

$$\rho_c^{-1} = \int d^3\alpha \prod_{i=1}^4 e^{i\vec{\alpha}\cdot\vec{r}_i} \varphi(\vec{r}_i) d^3r_i = \int d^3\alpha \prod_{i=1}^4 e^{i\vec{\alpha}\cdot\vec{r}_i} \left(\sum_{j=1}^N C_j e^{\vec{r}_i^2/R_j^2} \right) d^3r_i. \quad (8)$$

Integration with respect to r_i gives

$$\begin{aligned} \rho_c^{-1} &= \int d^3\alpha \prod_{i=1}^4 \sum_{j=1}^N C_j (\pi R_j^2)^{3/2} e^{-\alpha^2 R_j^2/4} = \int d^3\alpha \left(\sum_{j=1}^N C_j (\pi R_j^2)^{3/2} e^{-\alpha^2 R_j^2/4} \right)^4 \\ &= \int d^3\alpha \sum_{i_1, i_2, i_3, i_4}^N C_{i_1} C_{i_2} C_{i_3} C_{i_4} \left(\prod_{s=1}^4 (\pi R_{i_s}^2)^{3/2} \right) e^{-\frac{\alpha^2}{4} (\sum_{j=1}^4 R_{i_j}^2)}, \end{aligned} \quad (9)$$

and final integration with respect to α yields

$$\rho_c^{-1} = \sum_{i_1, i_2, i_3, i_4}^N C_{i_1} C_{i_2} C_{i_3} C_{i_4} \left(\prod_{s=1}^4 (\pi R_{i_s}^2)^{3/2} \right) \left(\frac{4\pi}{\sum_{j=1}^4 R_{i_j}^2} \right)^{3/2}. \quad (10)$$

The one-particle density function is determined as

$$\rho(\vec{r}) = \int |\psi(\vec{r}, \vec{r}_2, \vec{r}_3, \vec{r}_4)|^2 d^3r_2 d^3r_3 d^3r_4, \quad (11)$$

and can be calculated in an analogous way. The functions $\rho(\vec{r})$ corresponding to the parametrizations (A – D) of the wave function are shown in Fig. 1. All densities are close to each other at large values of r , and they are different in the nucleus center. So, the parametrizations take various short range NN correlations into account.




Figure 1: The one-particle density of the 4He

Performing nearly the same calculations, we have the following expression for the form factor:

$$\begin{aligned}
F(\vec{q}) &= \int e^{i\vec{q}\cdot\vec{r}_1} (2\pi)^3 \rho_c \delta\left(\sum_{i=1}^4 \vec{r}_i\right) \prod_{i=1}^4 \varphi(\vec{r}_i) d^3 r_i \\
&= (2\pi)^3 \rho_c \int \frac{d^3 \alpha}{(2\pi)^3} e^{i\vec{q}\cdot\vec{r}_1} e^{i\vec{\alpha}\cdot(\sum_{i=1}^4 \vec{r}_i)} \prod_{i=1}^4 \left(\sum_{j=1}^N C_j e^{-\vec{r}_i^2/R_j^2}\right) d^3 r_i \\
&= \rho_c \int d^3 \alpha \left(\sum_{j=1}^N C_j e^{i(\vec{q}+\vec{\alpha})\cdot\vec{r}_1} e^{-\vec{r}_1^2/R_j^2}\right) d^3 r_1 \prod_{i=2}^4 \left(\sum_{j=1}^N C_j e^{-\vec{r}_i^2/R_j^2} e^{i\vec{\alpha}\cdot\vec{r}_i}\right) d^3 r_i. \quad (12)
\end{aligned}$$

Integrating it with respect to r_i we obtain

$$\begin{aligned}
F(\vec{q}) &= \rho_c \int d^3 \alpha \left(\sum_{j=1}^N C_j (\pi R_j^2)^{3/2} e^{\frac{R_j^2}{4}(\vec{q}+\vec{\alpha})^2}\right) \prod_{i=2}^4 \left(\sum_{j=1}^N C_j (\pi R_j^2)^{3/2} e^{-\frac{R_j^2 \alpha^2}{4}}\right) \quad (13) \\
&= \rho_c \int d^3 \alpha \left(\sum_{j=1}^N C_j (\pi R_j^2)^{3/2} e^{\frac{R_j^2}{4}(\vec{q}+\vec{\alpha})^2}\right) \left(\sum_{j=1}^N C_j (\pi R_j^2)^{3/2} e^{-\frac{R_j^2 \alpha^2}{4}}\right)^3 \\
&= \rho_c \sum_{i_1, i_2, i_3, i_4=1}^N C_{i_1} C_{i_2} C_{i_3} C_{i_4} \int d^3 \alpha \left(\prod_{j=1}^4 (\pi R_{i_j}^2)^{3/2}\right) e^{\frac{R_{i_1}^2}{4}(\vec{q}^2+2\vec{\alpha}\cdot\vec{q})} e^{-\frac{\alpha^2}{4}(\sum_{j=1}^4 R_{i_j}^2)} \\
&= \rho_c \sum_{i_1, i_2, i_3, i_4=1}^N C_{i_1} C_{i_2} C_{i_3} C_{i_4} \left(\prod_{i_j=1}^4 (\pi R_{i_j}^2)^{3/2}\right) \left(\frac{4\pi}{\sum_{j=1}^4 R_{i_j}^2}\right)^{3/2} \\
&\quad \cdot \exp\left(-\frac{R_{i_1}^2 \vec{q}^2}{4} \left[\frac{\sum_{j=2}^4 R_{i_j}^2}{\sum_{i_j=1}^4 R_{i_j}^2}\right]\right).
\end{aligned}$$

The charge form factor, $F_{ch}(\vec{q})$, of the ${}^4\text{He}$ is connected with $F(\vec{q})$,

$$F_{ch}(\vec{q}) = F(\vec{q}) G_N(\vec{q}), \quad (14)$$

where $G_N(\vec{q})$ is the nucleon charge form factor, $G_N(\vec{q}) = G_p(\vec{q}) + G_n(\vec{q})$. G_p is the proton form factor chosen in the dipole form [9], $G_p(t) = (1 - t/0.71)^{-2}$. G_n is the neutron form factor, $G_n(\vec{q}) = (1+r_1^2 q^2)^{-2} - (1+r_2^2 q^2)^{-2}$, where $r_1^2 = 1.24 (GeV/c)^{-2}$, $r_2^2 = 1.50 (GeV/c)^{-2}$ [1]. $t = -q^2$ is the four momentum transfer in $(GeV/c)^2$.

In Fig. 2 the charge form factor calculations at the different parametrizations ($B-D$) are compared with the experimental data of R.F. Frosch et. al. [10]. The charge form factor predicted by parametrization A is not presented because it does not reproduce the data at $q^2 > 0.35 (GeV/c)^2$. As seen, at small values of t all parametrizations give the same good description of the data. They are different only at large values of t due to the difference of the corresponding one-particles densities in the center of the nucleus (see Fig. 1). We consider parametrization D as the best one though it gives a dip position at a somewhat smaller value of t than it is needed for a perfect description of the data. We think that the inclusion of the twelve quark bag component of the ground state wave function will not change the results drastically (see consideration in Ref. [1]).




Figure 2: The charge form-factor of the ${}^4\text{He}$ nucleus. The points are the experimental data [10], lines – our calculations.

2 The differential elastic cross section

The Glauber amplitude for hadron-nucleus scattering has a form [2]:

$$F_{1A}(\vec{q}) = \frac{ip}{2\pi} \int d^2b e^{i\vec{q}\cdot\vec{b}} \langle \psi_f | 1 - \prod_{j=1}^A (1 - \gamma(\vec{b} - \vec{s}_j)) | \psi_i \rangle, \quad (15)$$

where \vec{b} is the impact parameter, p is the momentum of the projectile hadron, ψ_i and ψ_f are initial and final states wave functions, respectively. γ is the NN elastic scattering amplitude in the impact parameter representation. The corresponding differential cross section is given as

$$\frac{d\sigma}{d\Omega} = |F_{1A}|^2. \quad (16)$$

In the case of the elastic $p^4\text{He}$ scattering the amplitude F_{1A} given by Eq. (15) can be re-written as

$$F_{14}(\vec{q}) = \frac{ip}{2\pi} \int d^2b e^{i\vec{q}\cdot\vec{b}} \left[1 - \prod_{j=1}^4 (1 - \gamma(\vec{b} - \vec{s}_j)) \right] |\psi(\vec{r}_1, \dots, \vec{r}_A)|^2 \prod_{j=1}^4 d^2r_j. \quad (17)$$

Substituting Eq. (2) in Eq. (17) gives

$$F_{14}(\vec{q}) = \frac{ip}{2\pi} \rho_c \int d^2b d^3\alpha e^{i\vec{q}\cdot\vec{b}} \left[1 - \prod_{j=1}^4 (1 - \gamma(\vec{b} - \vec{s}_j)) \right] e^{i\vec{\alpha}\cdot\sum_{j=1}^4 \vec{r}_j} \prod_{j=1}^4 \phi(\vec{r}_j) d^3r_j, \quad (18)$$

where $\vec{r} = \vec{s} + \vec{z}$, \vec{z} is the component of the position vector \vec{r} in the direction along the projectile momentum \vec{p} . We assume it is the direction of the z-axis. Taking into account

the relation (4), we obtain

$$\begin{aligned}
\frac{2\pi}{i p \rho_c} F_{14}(\vec{q}) &= \int d^2b e^{i\vec{q}\cdot\vec{b}} \left[1 - \prod_{j=1}^4 (1 - \gamma(\vec{b} - \vec{s}_j)) \right] \left(d^3\alpha e^{i\vec{\alpha}\cdot(\sum_{j=1}^4 \vec{r}_j)} \right). \quad (19) \\
&\cdot \prod_{j=1}^4 \sum_{k=1}^N C_k e^{-\vec{r}_j^2/R_k^2} d^3r_j = \int d^2b d^3\alpha e^{i\vec{q}\cdot\vec{b}} \left[1 - \prod_{j=1}^4 (1 - \gamma(\vec{b} - \vec{s}_j)) \right] \\
&\cdot \prod_{j=1}^4 \sum_{k=1}^N C_k e^{i\vec{\alpha}\cdot\vec{r}_j} e^{-\vec{r}_j^2/R_k^2} d^3r_j,
\end{aligned}$$

$\vec{\alpha} = \vec{\alpha}_2 + \vec{\alpha}_1$. As before $\vec{\alpha}_1$ is the component of $\vec{\alpha}$ in the z-axis direction, and $\vec{\alpha}_2$ is two-dimensional vector orthogonal to \vec{p} . Having that

$$\begin{aligned}
1 - \prod_{j=1}^4 (1 - \gamma(\vec{b} - \vec{s}_j)) &= \sum_{k=1}^4 \gamma(\vec{b} - \vec{s}_k) - \sum_{k_1, k_2=1}^4 \gamma(\vec{b} - \vec{s}_{k_1}) \gamma(\vec{b} - \vec{s}_{k_2}) \\
&+ \sum_{k_1, k_2, k_3=1}^4 \gamma(\vec{b} - \vec{s}_{k_1}) \gamma(\vec{b} - \vec{s}_{k_2}) \gamma(\vec{b} - \vec{s}_{k_3}) \\
&- \gamma(\vec{b} - \vec{s}_{k_1}) \gamma(\vec{b} - \vec{s}_{k_2}) \gamma(\vec{b} - \vec{s}_{k_3}) \gamma(\vec{b} - \vec{s}_{k_4}), \quad (20)
\end{aligned}$$

we can write the amplitude as a sum of the multiple scattering terms

$$\frac{2\pi}{i p \rho_c} F_{14}(\vec{q}) = F_{14}^{(1)} - F_{14}^{(2)} + F_{14}^{(3)} - F_{14}^{(4)}. \quad (21)$$

Every term can be calculated separately if γ is chosen as

$$\gamma(\vec{b}) = \beta e^{-\vec{b}^2/2B_{NN}}, \quad (22)$$

where $\beta = (\sigma_{NN}^{tot}(1 - i\alpha_{NN}) / (4\pi B_{NN}))$, σ_{NN}^{tot} is the NN total cross section, B_{NN} – the slope parameter of the NN differential elastic cross section at zero momentum transfer, α_{NN} – the ratio of the real to imaginary parts of the NN elastic scattering amplitude at zero momentum transfer. Then, the first term will be

$$\begin{aligned}
F_{14}^{(1)} &= \int d^2b d^3\alpha \left(\sum_{k_1=1}^4 \gamma(\vec{b} - \vec{s}_{k_1}) \right) e^{i\vec{q}\cdot\vec{b}} \prod_{j=1}^4 \sum_{i=1}^N C_i e^{i\vec{\alpha}\cdot\vec{r}_j} e^{-\vec{r}_j^2/R_i^2} d^3r_j \\
&= \int d^2b d^3\alpha e^{i\vec{q}\cdot\vec{b}} \left(\sum_{k_1=1}^4 \gamma(\vec{b} - \vec{s}_{k_1}) \right) \left(\sum_{i=1}^N C_i e^{i\vec{\alpha}\cdot\vec{r}_{k_1}} e^{-\vec{r}_{k_1}^2/R_i^2} d^3r_{k_1} \right) \\
&\quad \prod_{k_2, k_3, k_4} \left(\sum_{i=1}^N C_i e^{i\vec{\alpha}\cdot\vec{r}_{k_j}} e^{-\vec{r}_{k_j}^2/R_i^2} d^3r_{k_j} \right) \\
&= \sum_{k_1=1}^4 \int d^2b d^3\alpha \gamma(\vec{b} - \vec{s}_{k_1}) e^{i\vec{q}\cdot\vec{b}} \left(\sum_{i=1}^N C_i e^{i\vec{\alpha}\cdot\vec{r}_{k_1}} e^{-\vec{r}_{k_1}^2/R_i^2} d^3r_{k_1} \right) \\
&\quad \prod_{k_2, k_3, k_4} \left(\sum_{i=1}^N C_i e^{i\vec{\alpha}\cdot\vec{r}_{k_j}} e^{-\vec{r}_{k_j}^2/R_i^2} d^3r_{k_j} \right)
\end{aligned}$$

$$\begin{aligned}
&= \sum_{k_1=1}^4 \int d^2b d^3\alpha e^{i\vec{q}\cdot\vec{b}} \left(\sum_{i=1}^N C_i \gamma(b-s_{k_1}) e^{i\vec{\alpha}\cdot\vec{r}_{k_1}} e^{-r_{k_1}^2/R_i^2} d^3r_{k_1} \right) \\
&\quad \prod_{k_2, k_3, k_4} \left(\sum_{i=1}^N C_i e^{i\vec{\alpha}\cdot\vec{r}_{k_j}} e^{-r_{k_j}^2/R_i^2} d^3r_{k_j} \right) \\
&= \sum_{k_1=1}^4 \beta \int d^2b d^3\alpha e^{i\vec{q}\cdot\vec{b}} \left(\sum_{i=1}^N C_i e^{(b-s_{k_1})^2} e^{i\vec{\alpha}\cdot\vec{r}_{k_1}} e^{-r_{k_1}^2/R_i^2} d^3r_{k_1} \right) \\
&\quad \prod_{k_2, k_3, k_4} \left(\sum_{i=1}^N C_i (\pi R_i^2)^{3/2} e^{-\alpha^2 R_i^2/2} \right) \\
&= \sum_{k_1=1}^4 \beta \int d^2b d^3\alpha e^{i\vec{q}\cdot\vec{b}} \left(\sum_{i=1}^N C_i (\pi R_i^2)^{3/2} e^{-\alpha^2 R_i^2/2} \right)^3 \\
&\quad e^{\frac{1}{4} \left(\frac{2BR_i^2}{2B+R_i^2} \right) \left(\frac{b}{B} + i\alpha_2 \right)^2} \left(\sum_{i=1}^N C_i e^{-b^2/2B} e^{-\alpha_1^2 R_i^2/4} \left((\pi R_i^2)^{1/2} \right. \right. \\
&\quad \left. \left. \exp \left[- \left(\frac{1}{R_i^2} + \frac{1}{2B} \right) \left(s_{k_1}^2 - \frac{1}{2} \left(\frac{b}{B} + i\alpha_2 \right) \left(\frac{2BR_i^2}{2B+R_i^2} \right) \right)^2 \right] \right) \right)^3 \\
&= \sum_{k_1=1}^4 \beta \int d^2b d^3\alpha e^{i\vec{q}\cdot\vec{b}} \left(\sum_{i=1}^N C_i (\pi R_i^2)^{3/2} e^{-\alpha^2 R_i^2/2} \right)^3 \\
&\quad \left(\sum_{i=1}^N C_i e^{-b^2/2B} e^{-\alpha_1^2 R_i^2/4} (\pi R_i^2)^{1/2} \left(\frac{2\pi B R_i^2}{2B+R_i^2} \right) \right. \\
&\quad \left. \exp \left[\frac{2B}{4} \left(\frac{R_i^2}{2B+R_i^2} \right) \left(\frac{b}{B} + i\alpha_2 \right)^2 \right] \right) \\
&= 4\beta \sum_{k_{i_j}, j=1}^4 C_{k_1} C_{k_2} C_{k_3} C_{k_4} \left(\prod_{j=1}^4 (\pi R_{k_j}^2)^{3/2} \right) \left(\frac{2B}{2B+R_{k_1}^2} \right) \int d^2b d^3\alpha e^{i\vec{q}\cdot\vec{b}} \\
&\quad e^{-\frac{\alpha_1^2}{4} \sum_{j=1}^4 R_{k_j}^2} e^{-\frac{\alpha_2^2}{4} \left(\sum_{j=2}^4 R_{k_j}^2 \right)} e^{-b/B} e^{i\vec{\alpha}\cdot\vec{b}} \left(\frac{R_{k_1}^2}{R_{k_1}^2+2B} \right) e^{-\frac{b^2}{2B} \left(\frac{R_{k_1}^2}{R_{k_1}^2+2B} \right)} e^{-\frac{B\alpha_2^2}{2} \left(\frac{R_{k_1}^2}{R_{k_1}^2+2B} \right)} \\
&= 4\beta \sum_{k_{i_j}, j=1}^4 C_{k_1} C_{k_2} C_{k_3} C_{k_4} \left(\prod_{j=1}^4 (\pi R_{k_j}^2)^{3/2} \right) \left(\frac{2B}{2B+R_j^2} \right) \left(\frac{4\pi}{\sum_{j=1}^4 R_{k_j}^2} \right)^{1/2} \\
&\quad \int d^2b d^3\alpha e^{-B\alpha_2^2 S_1/2} e^{-\frac{\alpha_2^2}{4} \left(\sum_{j=2}^4 R_{k_i}^2 \right)} e^{-b/B} e^{-(i(\vec{q}+\vec{\alpha}_2 S_1)/2M_1)^2} \\
&\quad \exp \left[-M_1 \left(b^2 - i(\vec{q}+\vec{\alpha}_2 S_1) \cdot \vec{b}/M_1 + (i(\vec{q}+\vec{\alpha}_2 S_1)/2M_1)^2 \right) \right] \tag{23}
\end{aligned}$$

where

$$M_1 = \frac{1}{B} - \frac{1}{B} \frac{R_{k_1}^2}{R_{k_1}^2 + 2B} = \frac{1}{R_{k_1}^2 + 2B}, \quad S_1 = \frac{R_{k_1}^2}{R_{k_1}^2 + 2B}.$$

$$\begin{aligned}
F_{14}^{(1)} &= 4\beta \sum_{k_{i_j}, j=1}^4 C_{k_1} C_{k_2} C_{k_3} C_{k_4} \left(\prod_{j=1}^4 (\pi R_{k_j}^2)^{3/2} \right) \left(\frac{2B}{2B+R_j^2} \right)^{3/2} \left(\frac{4\pi}{\sum_{j=1}^4 R_{k_j}^2} \right)^{1/2} \\
&\quad \left(\frac{\pi}{M_1} \right) \left(\frac{4\pi}{H_1} \right) e^{-q^2/4M_1} \exp \left[q^2 S_1^2 / 4M_1 H_1 \right] \tag{24}
\end{aligned}$$

$$H_1 = R_{k_1}^2 + R_{k_3}^2 + R_{k_4}^2 + 2BS_1 + \frac{S_1^2}{M_1}.$$

The second term also will be,

$$\begin{aligned}
F_{14}^{(2)} &= \int d^2b d^3\alpha \left(\sum_{k_1, k_2=1}^4 \gamma(b - s_{k_1})\gamma(b - s_{k_2}) \right) e^{i\vec{q}\cdot\vec{b}} \prod_{j=1}^4 \sum_{i=1}^N C_j e^{i\vec{\alpha}\cdot\vec{r}_j} e^{-r_j^2/R_i^2} d^3r_i \\
&= \int d^2b d^3\alpha e^{i\vec{q}\cdot\vec{b}} \left(\sum_{k_1, k_2=1}^4 \gamma(b - s_{k_1})\gamma(b - s_{k_2}) \right) \prod_{k_1, k_2} \left(\sum_{i=1}^N C_i e^{i\vec{\alpha}\cdot\vec{r}_{k_j}} e^{-r_{k_j}^2/R_i^2} d^3r_{k_j} \right) \\
&\quad \prod_{k_3, k_4} \left(\sum_{i=1}^N C_i e^{i\vec{\alpha}\cdot\vec{r}_{k_j}} e^{-r_{k_j}^2/R_i^2} d^3r_{k_j} \right) \\
&= \sum_{k_1, k_2=1}^4 \int d^2b d^3\alpha \gamma(b - s_{k_1})\gamma(b - s_{k_2}) e^{i\vec{q}\cdot\vec{b}} \prod_{k_1, k_2} \left(\sum_{i=1}^N C_i e^{i\vec{\alpha}\cdot\vec{r}_{k_j}} e^{-r_{k_j}^2/R_i^2} d^3r_{k_j} \right) \\
&\quad \prod_{k_3, k_4} \left(\sum_{i=1}^N C_i e^{i\vec{\alpha}\cdot\vec{r}_{k_j}} e^{-r_{k_j}^2/R_i^2} d^3r_{k_j} \right) \\
&= \sum_{k_1, k_2=1}^4 \int d^2b d^3\alpha e^{i\vec{q}\cdot\vec{b}} \prod_{k_1, k_2} \left(\sum_{i=1}^N C_i \gamma(b - s_{k_j}) e^{i\vec{\alpha}\cdot\vec{r}_{k_j}} e^{-r_{k_j}^2/R_i^2} d^3r_{k_j} \right) \\
&\quad \prod_{k_3, k_4} \left(\sum_{i=1}^N C_i e^{i\vec{\alpha}\cdot\vec{r}_{k_j}} e^{-r_{k_j}^2/R_i^2} d^3r_{k_j} \right) \\
&= \sum_{k_1, k_2=1}^4 \beta^2 \int d^2b d^3\alpha e^{i\vec{q}\cdot\vec{b}} \prod_{k_1, k_2} \left(\sum_{i=1}^N C_i e^{(b-s_{k_j})^2} e^{i\vec{\alpha}\cdot\vec{r}_{k_j}} e^{-r_{k_j}^2/R_i^2} d^3r_{k_j} \right) \\
&\quad \prod_{k_3, k_4} \left(\sum_{i=1}^N C_i (\pi R_i^2)^{3/2} e^{-\alpha^2 R_i^2/2} \right) \\
&= \sum_{k_1, k_2=1}^4 \beta^2 \int d^2b d^3\alpha e^{i\vec{q}\cdot\vec{b}} \left(\sum_{i=1}^N C_i (\pi R_i^2)^{3/2} e^{-\alpha^2 R_i^2/2} \right)^2 \\
&\quad \prod_{k_1, k_2} \left(\sum_{i=1}^N C_i e^{-b^2/2B} e^{-\alpha_1^2 R_i^2/4} ((\pi R_i^2)^{1/2} e^{\frac{1}{4} \left(\frac{2BR_i^2}{2B+R_i^2} \right) \left(\frac{b}{B} + i\alpha_2 \right)^2} \right. \\
&\quad \left. \exp \left[- \left(\frac{1}{R_i^2} + \frac{1}{2B} \right) \left(s_{k_j}^2 - \frac{1}{2} \left(\frac{b}{B} + i\alpha_2 \right) \left(\frac{2BR_i^2}{2B+R_i^2} \right) \right)^2 \right] \right) \\
&= \sum_{k_1, k_2=1}^4 \beta^2 \int d^2b d^3\alpha e^{i\vec{q}\cdot\vec{b}} \left(\sum_{i=1}^N C_i (\pi R_j^2)^{3/2} e^{-\alpha^2 R_j^2/2} \right)^2 \\
&\quad \prod_{k_1, k_2} \left(\sum_{i=1}^N C_i e^{-b^2/2B} e^{-\alpha_1^2 R_i^2/4} (\pi R_j^2)^{1/2} \left(\frac{2\pi B R_j^2}{2B+R_i^2} \right) \right. \\
&\quad \left. \exp \left[\frac{2B}{4} \left(\frac{R_i^2}{2B+R_i^2} \right) \left(\frac{b}{B} + i\alpha_2 \right)^2 \right] \right) \\
&= 6\beta^2 \sum_{k_i, j=1}^4 C_{k_1} C_{k_2} C_{k_3} C_{k_4} \left(\prod_{j=1}^4 (\pi R_{k_i}^2)^{3/2} \right) \prod_{j=1}^2 \left(\frac{2B}{2B+R_i^2} \right)
\end{aligned}$$

$$\begin{aligned}
& \int d^2b d^3\alpha e^{i\vec{q}\cdot\vec{b}} e^{-\frac{\alpha_1^2}{4}\sum_{j=1}^4 R_{k_j}^2} e^{-\frac{\alpha_2^2}{4}(R_{k_3}^2+R_{k_4}^2)} \\
& e^{-b/B} e^{i\vec{\alpha}\cdot\vec{b}\left(\sum_{j=1}^2 \frac{R_{k_j}^2}{R_{k_j}^2+2B}\right)} e^{-\frac{b^2}{2B}\left(\sum_{j=1}^2 \frac{R_{k_j}^2}{R_{k_j}^2+2B}\right)} e^{-\frac{B\alpha_2^2}{2}\left(\sum_{j=1}^2 \frac{R_{k_j}^2}{R_{k_j}^2+2B}\right)} \\
& = 6\beta^2 \sum_{k_{i_j}, j=1}^4 C_{k_1} C_{k_2} C_{k_3} C_{k_4} \left(\prod_{j=1}^4 (\pi R_{k_j}^2)^{3/2} \right) \prod_{j=1}^2 \left(\frac{2B}{2B+R_j^2} \right) \left(\frac{4\pi}{\sum_{j=1}^4 R_{k_j}^2} \right)^{1/2} \\
& \int d^2b d^3\alpha e^{-B\alpha_2^2 S_2/2} e^{-\frac{\alpha_2^2}{4}(R_{k_3}^2+R_{k_4}^2)} e^{-b/B} e^{-i(\vec{q}+\vec{\alpha}_2 S_2)/2M_2} \\
& \exp \left[-M_2 \left(b^2 - i(\vec{q} + \vec{\alpha}_2 S_2) \cdot \vec{b}/M_2 + (i(\vec{q} + \vec{\alpha}_2 S_2)/2M_2)^2 \right) \right], \tag{25}
\end{aligned}$$

where

$$M_2 = \frac{1}{B} - \frac{1}{2B} \sum_{j=1}^2 \frac{R_{k_j}^2}{R_{k_j}^2 + 2B} = \sum_{j=1}^2 \frac{1}{R_{k_j}^2 + 2B}, \quad S_2 = \sum_{j=1}^2 \frac{R_{k_j}^2}{R_{k_j}^2 + 2B}.$$

$$\begin{aligned}
F_{14}^{(2)} & = 6\beta^2 \sum_{k_{i_j}, j=1}^4 C_{k_1} C_{k_2} C_{k_3} C_{k_4} \left(\prod_{j=1}^4 (\pi R_{k_j}^2)^{3/2} \right) \prod_{j=1}^2 \left(\frac{2B}{2B+R_j^2} \right) \left(\frac{4\pi}{\sum_{j=1}^4 R_{k_j}^2} \right)^{1/2} \\
& \left(\frac{\pi}{M_2} \right) \left(\frac{4\pi}{H_2} \right) e^{-q^2/4M_2} \exp \left[q^2 S_2^2 / 4M_2 H_2 \right] \tag{26}
\end{aligned}$$

$$H_2 = R_{k_3}^2 + R_{k_4}^2 + 2BS_2 + \frac{S_2^2}{M_2}.$$

By the same way the other terms will be

$$\begin{aligned}
F_{14}^{(3)} & = 4\beta^3 \sum_{k_{i_j}, j=1}^4 C_{k_1} C_{k_2} C_{k_3} C_{k_4} \left(\prod_{j=1}^4 (\pi R_{k_j}^2)^{3/2} \right) \prod_{j=1}^2 \left(\frac{2B}{2B+R_j^2} \right) \left(\frac{4\pi}{\sum_{j=1}^4 R_{k_j}^2} \right)^{1/2} \\
& \left(\frac{\pi}{M_3} \right) \left(\frac{4\pi}{H_3} \right) e^{-q^2/4M_3} \exp \left[q^2 S_3^2 / 4M_3 H_3 \right], \tag{27}
\end{aligned}$$

$$M_3 = \frac{3}{2B} - \frac{1}{2B} \sum_{j=1}^3 \frac{R_{k_j}^2}{R_{k_j}^2 + 2B} = \sum_{j=1}^3 \frac{1}{R_{k_j}^2 + 2B}, \quad S_3 = \sum_{j=1}^3 \frac{R_{k_j}^2}{R_{k_j}^2 + 2B},$$

$$H_3 = R_{k_4}^2 + 2BS_3 + \frac{S_3^2}{M_3}$$

$$\begin{aligned}
F_{14}^{(4)} & = \beta^4 \sum_{k_{i_j}, j=1}^4 C_{k_1} C_{k_2} C_{k_3} C_{k_4} \left(\prod_{j=1}^4 (\pi R_{k_j}^2)^{3/2} \right) \prod_{j=1}^2 \left(\frac{2B}{2B+R_j^2} \right) \left(\frac{4\pi}{\sum_{j=1}^4 R_{k_j}^2} \right)^{1/2} \\
& \left(\frac{\pi}{M_4} \right) \left(\frac{4\pi}{H_4} \right) e^{-q^2/4M_4} \exp \left[q^2 S_4^2 / 4M_4 H_4 \right] \tag{28}
\end{aligned}$$

$$M_4 = \frac{2}{B} - \frac{1}{2B} \sum_{j=1}^4 \frac{R_{k_j}^2}{R_{k_j}^2 + 2B} = \sum_{j=1}^4 \frac{1}{R_{k_j}^2 + 2B}, \quad S_4 = \sum_{j=1}^4 \frac{R_{k_j}^2}{R_{k_j}^2 + 2B}, \quad H_4 = 2BS_4 + \frac{S_4^2}{M_4}.$$

In many experimental papers [11, 12, 13] the authors included the Coulomb scattering amplitude in a simple way in order to extract the nuclear total p^4He cross section,

$$\frac{d\sigma}{dt} = |F_c e^{i\phi} + F_N|^2, \quad (29)$$

where

$$F_c(t) = \frac{4\alpha\sqrt{\pi}}{\beta t} G_p(t) G_{He}(t), \quad (30)$$

$\alpha = 1/137$ is the fine structure constant, $\beta = v/c$ is the proton velocity in the laboratory system, $G_p(t), G_{He}(t)$ are the electromagnetic form factor of the proton and He , respectively,

$$G_{He}(t) = \exp\left[\frac{r_{He}^2 t}{6}\right], \quad (31)$$

$$r_{He}^2 = r_e^2 - r_p^2, \quad r_e = 1.67 \text{ fm}, \quad r_p = 0.812 \text{ fm}.$$

$$\phi = 2\alpha\beta^{-1} [\ln(B|t|) + 0.577\dots]. \quad (32)$$

$B = 29 (GeV/c)^{-2}$ and $t = -q^2$. F_n is the nuclear amplitude written at small t in the form

$$F_n = \frac{\sigma^{tot}}{4\hbar\sqrt{\pi}} (i + \alpha) e^{\frac{1}{2}Bt}. \quad (33)$$

We follow the same way replacing F_N by the Glauber scattering amplitude.

The available experimental data on the p^4He elastic scattering have been presented by G.N. Velichko et. al. [11] at the energies of 0.695, 0.793, 0.89, 0.991 GeV ; by A. Bujak et. al. [12] at the energies of 45, 97, 146, 200, 259, 301, 393 GeV , and by J.P. Burq et al. [13] at the energies of 100, 150, 250, 300 GeV . To calculate the Glauber amplitudes at these energies, it is needed to have the values of the nucleon-nucleon amplitude parameters σ_{NN}^{tot} , B_{NN} and α_{NN} . σ_{NN}^{tot} was estimated as an average of the neutron-proton total cross section, σ_{tot}^{np} , and the proton-proton total cross section, σ_{tot}^{pp} , which can be taken from the compilation of the experiential data [14].

More complicated situation is with B_{NN} . There are only few experimental data, and it is not enough for all energies. Thus we have used another way to evaluate B_{NN} from the total and elastic NN cross sections. At chosen form of $\gamma(\vec{b})$ (see Eq. 22) the elastic NN cross section, σ_{NN}^{el} , is given as

$$\begin{aligned} \sigma_{NN}^{el} &= \frac{1}{p^2} \left(\frac{p}{2\pi}\right)^2 \int \gamma(\vec{b}_1) e^{i\vec{q}\cdot\vec{b}_1} \gamma^*(\vec{b}_2) e^{-i\vec{q}\cdot\vec{b}_2} d^2b_1 d^2b_2 d^2q \\ &= \frac{1}{(2\pi)^2} \int e^{i\vec{q}\cdot(\vec{b}_1 - \vec{b}_2)} \gamma(\vec{b}_1) \gamma^*(\vec{b}_2) d^2b_1 d^2b_2 d^2q \\ &= \int \delta(\vec{b}_1 - \vec{b}_2) \gamma(\vec{b}_1) \gamma^*(\vec{b}_2) d^2b_1 d^2b_2 = \int |\gamma(\vec{b})|^2 d^2b \\ &= \left(\frac{\sigma_{NN}^{tot}}{4\pi B_{NN}}\right)^2 (1 + \alpha_{NN}^2) \int e^{-\frac{b^2}{B_{NN}}} d^2b \\ &= \left(\frac{\sigma_{NN}^{tot}}{4\pi B_{NN}}\right)^2 (1 + \alpha_{NN}^2) \pi B_{NN} = \frac{(\sigma_{NN}^{tot})^2}{16\pi B_{NN}} (1 + \alpha_{NN}^2). \end{aligned} \quad (34)$$

Since α_{NN} is very small, we neglect it in our calculations. In this case B_{NN} can be calculated as

$$B_{NN} = \frac{(\sigma_{NN}^{tot})^2}{16\pi\sigma_{NN}^{el}}. \quad (35)$$

σ_{NN}^{el} was taken from the compilation of the experimental data [14] as an average of pn and pp cross sections.

The values of α_{NN} for all mentioned above energies were extracted from the compilation of the experiential data [15].

Table 2: The parameters used at the calculations of the Glauber amplitudes

E_{kin} GeV	σ_{NN}^{el} mb	σ_{pp}^{tot} mb	σ_{pn}^{tot} mb	σ_{NN}^{tot} mb	B_{NN} (GeV/c) ⁻²	α_{NN}
0.695	24.2	42.4	38.38	40.39	4.069	-0.205
0.795	22.5	46.8	38.56	42.68	4.134	-0.1975
0.890	24.4	47.3	38.73	43.01	3.872	-0.19
0.991	24.27	47.6	39.24	43.42	3.967	-0.185
0.992	24.27	47.6	39.24	43.42	3.967	-0.185
45	7.402	38.48	38.32	38.4	10.173	-0.087
97	6.985	37.94	38.89	38.4	10.783	-0.090
100	6.985	37.94	38.89	38.4	10.783	0.1
146	7.03	38.29	39.12	38.71	10.884	-0.049
150	7.03	38.69	39.12	38.91	10.996	0.105
200	6.895	38.98	39.56	39.27	11.422	-0.022
250	6.89	39.34	39.83	39.58	11.614	0.11
259	6.89	39.34	39.83	39.58	11.614	0.024
300	6.888	39.46	39.83	39.65	11.653	0.115
301	6.888	39.46	39.83	39.65	11.653	0.031
393	7.016	40.19	40.01	40.1	11.703	0.067

All the parameter values used for our calculation are presented in Table 2. Typical results of the calculations in comparison with experimental data [11, 12] are shown in Figs. 3, 4. As seen, the model calculations are above the experimental data. The first diffraction minimums are shifted to small t . We can confirm now that the model calculations can not reproduce the data with required accuracy. This pushed us to search for modification of the model.

3 The twelve quark bag admixture

Any nucleus consists of $3A$ -quarks. In the ground state the quarks are forming clusters, bags and nucleons. Following [1] we assume that the 4He wave function is given as

$$|q_1\dots q_{12}\rangle = \alpha|NNNN\rangle + \beta|12q\rangle, \quad |\alpha|^2 + |\beta|^2 = 1, \quad (36)$$

where $|q_1\dots q_{12}\rangle$ is the 12- quark bound state wave function of the 4He . $|12q\rangle$ stands for a component left after projecting $|q_1\dots q_{12}\rangle$ onto the four nucleons state. Since $|NNNN\rangle$ vanishes in the central part of 4He , $|12q\rangle$ must strongly peak in the central part of the

4He . According to the assumption of Ref. [1] $\langle NNNN|12q \rangle = 0$. Thus we neglect $12q - NNNN$ interference terms that vanish at $q = 0$ in the elastic p^4He scattering amplitude and write

$$F_{14} = (1 - w_{12q}) F_{Gl} + w_{12q} F_{12q}, \quad (37)$$

where F_{Gl} is the Glauber amplitude of the $p - 4N$ scattering, and w_{12q} is the weight of the $12q$ bag quark state, $w_{12q} = |\beta|^2$. We take the nucleon - twelve quark bag scattering amplitude in a simple form,

$$F_{12q} = \frac{\sigma_{12q}}{2} e^{tb_{12q}/2}, \quad (38)$$

where σ_{12q} is the $N - 12q$ bag total cross section, and b_{12q} is the slope parameter.

We found the parameters σ_{12q} , b_{12q} and w_{12q} fitting the experimental data [11, 12, 13]. The values are presented in Table 3. As one can see, the parameter uncertainty is very large at low energies ($E_{kin} < 1 \text{ GeV}$). This means that at the energies one does not need to add anything to the Glauber amplitude. At higher energies the values become more stable excepting the results at 146 GeV.

Table 3: The fitting values of w_{12q} , σ_{12q} and b_{12q}

E_{kin} GeV	w_{12q}	σ_{12q} mb	b_{12q} (GeV/c) $^{-2}$	χ^2/NOF
0.695	4.1 ± 66.5	126.0 ± 169.7	32.6 ± 100.5	72/65
0.795	9.8 ± 1.8	117.0 ± 10.7	40 ± 228.9	45/81
0.890	10.6 ± 174.4	169.4 ± 464.4	37.4 ± 164.2	56/95
45	8.34 ± 0.92	32.62 ± 10.23	20.38 ± 3.90	557/127
97	9.23 ± 1.31	30.83 ± 13.37	20.46 ± 5.54	146/84
146	13.51 ± 0.42	65.57 ± 2.54	32.46 ± 0.52	222/84
200	10.51 ± 1.83	28.86 ± 17.24	21.62 ± 7.33	285/84
259	9.72 ± 2.91	29.10 ± 30.04	22.11 ± 12.69	264/86
301	11.08 ± 1.28	25.94 ± 11.78	21.68 ± 5.46	173/86
393	10.80 ± 1.59	25.73 ± 14.95	21.14 ± 6.96	118/85

We have excluded from the fit the data at the energies of 100, 150, 250, and 300 GeV [13]. The data are above the Glauber calculations. Thus at the fitting an unreasonable large weight of 12q-bag ($> 50\%$) and σ_{12q} was obtained. We believe the data are not quite well normalized. To show this, we plot the data at close energies [12] on the same figures 3, 4.

As seen, there is a clear difference between the two groups of experimental data. Maybe, it is due to a normalization error. We do not know a reason of the error. However, one can see that the data by Ref. [13] are falling out from the whole set of the experimental data, and it is not possible to fit them correctly.

The figures show influence of the 12q bag admixture on the differential cross section. The inclusion of the admixture leads to decreasing the Glauber amplitude if σ_{12q} is smaller than σ_{pHe} . In the region of the dip where the Glauber amplitude vanishes, F_{12q} is positive and shifts the dip to a larger values of t . So, the hypothesis really allows one to solve the main part of the problem.




Figure 3: The p^4He differential elastic cross sections. The point are the experimental data [12, 13]. The solid and dashed lines are our calculations with and without 12q admixture, respectively.

Clearly, inclusion of the inelastic screening into calculations will lead to decreasing the cross section in the region of small t , and to increasing in the region of the large t values. To compensate these, one have to increase w_{12q} , σ_{12q} , and b_{12q} . From this point of view we can understand the results of Ref. [1]. According to the Fig. 10 of the Ref. [1], $\sigma_{12q} \sim 140$ mb what is near to the p^4He total cross section, the slope parameter of that F_{12q} is larger than ours. As a result, $w_{12q} \sim 12$ %. We have the average value of $w_{12q} \sim 10.5\%$ So, two values agree quite reasonable with each other. At the same time, our σ_{12q} is too small.

Let us mark that our amplitude F_{12q} is more simple than that of the Ref. [1]. It can be easily used for future calculations.




Figure 4: The p^4He differential elastic cross sections. Points show experimental data [12, 13]. Solid and dashed lines are our calculations with and without 12q admixture, respectively.

Conclusion

The 12q bag admixture to the ground state wave function of the 4He nucleus allows one to describe quite well the elastic p^4He scattering. According to our estimations, the weight of the 12q bag is $\sim 10.5\%$, the proton - 12q bag total cross section is ~ 34 mb, and the slope parameter of the $p - 12q$ bag elastic scattering is ~ 23 $(GeV/c)^{-2}$.

V.V.Uzhinskii thanks RFBR (grand N 00-01-00307) and INTAS (grand N 00-00366) for their financial support. A.M. Mosallem thanks to Profs. A.B.B. Kalila and K.M. Hanna and for support. The authors also wish to thank JINR officials for hospitality.

References

- [1] L.G. Dakno and N.N. Nikolaev, *Nucl. Phys.* **A436** (1985) 653.
- [2] R.J. Glauber, *Lectures in theoretical physics*. W.E. Brittin, L.G. Dunham (eds.). Vol. 1, p.315, New York: Interscience 1959; A.G. Sitenko, *Ukr. Fiz. Zh.* **4** (1959) 152.
- [3] E.S. Abers et al., *Nuovo Cimento* **42A** (1966) 365.
- [4] V.N. Gribov, *Sov. Phys. JETP* **29** (1969) 483.
- [5] J. Pumplin and M. Ross, *Phys. Rev. Lett.* **21** (1968) 1778.
- [6] G. Alberi and L. Bertocchi, *Nuovo Cimento* **61A** (1969) 201.
- [7] D.R. Harrington, *Phys. Rev.* **D1** (1970) 260.
- [8] C. Quigg and L.L. Wang, *Phys. Lett.* **42** (1973) 314.
- [9] R.P. Feynman, *Photon - Hadron Interactions* (1972) W.A.Benjamin, Inc Reading, Massachusetts.
- [10] R.F. Frosch, J.S. McCarthy, R.E. Rand, and M.R. Yearian, *Phys. Rev.* **160** (1967) 874.
- [11] G.N. Velichko et. al., *Yad. Fiz.* **42** (1985) 1325.
- [12] A. Bujak et. al., *Phys. Rev.* **D23** (1981) 1895; JINR Preprint E1-81-289 (1981).
- [13] J.P. Burk et. al., *Nucl. Phys.* **B187** (1981) 205.
- [14] V. Flaminio, W.G. Moorhead, D.R.O. Morrison, N. Rivoire, *Compilation of cross-sections: p and \bar{p} induced reactions* CERN-HERA 84-01 (1984).
- [15] O. Benary, L.R. Price, G. Alexander, *NN and ND interactions above 0.5 GeV/c - a compilation* UCRL-20000NN (1970).