

Final State Interaction Effects in Incoherent Pion Photoproduction on the Deuteron *

E.M. Darwish[†], H. Arenhövel, M. Schwamb

Institut für Kernphysik, Johannes Gutenberg-Universität, 55099 Mainz,
Germany

In the present work (nucl-th/0208030; *Eur. Phys. J. A* (in print)), incoherent photoproduction of pions on the deuteron in the $\Delta(1232)$ resonance region is investigated where besides the impulse approximation (IA) complete two-body rescattering in the nucleon-nucleon (NN) and pion-nucleon (πN) final state subsystems is included (see Fig. 1). The elementary $\gamma N \rightarrow \pi N$ amplitude including Born and $\Delta(1232)$ resonance contributions is taken from previous work of R. Schmidt *et al.* (*Z. Phys. A* **355**, 421 (1996)). The t -matrices for NN - and




Figure 1. Diagrams for $d(\gamma, \pi)NN$: (a) IA, (b) NN rescattering, (c) πN rescattering.




Figure 2. Total cross section for $d(\gamma, \pi)NN$. Dashed: IA; solid: IA+complete rescattering; data: Asai *et al.*, *Phys. Rev. C* **42**, 837 (1990), Benz *et al.*, *Nucl. Phys. A* **65**, 158 (1973) (ABHHM), Chiefari *et al.*, *Lett. Nuovo Cim.* **13**, 129 (1975) (Frascati).

πN -scattering are obtained from the separable interactions of Haidenbauer *et*

*Supported by Deutsche Forschungsgemeinschaft (SFB 443).

[†]Supported by Deutscher Akademischer Austauschdienst. *Present address:* Physics Department, Faculty of Science, South Valley University, Sohag, Egypt.

al. (*Phys. Rev. C* **30**, 1822 (1984)) and Nozawa *et al.* (*Nucl. Phys. A* **513**, 459 (1990)), respectively. Final state interaction effects are found to be significant in total (see Fig. 2) and differential cross sections (see Fig. 3), in particular for π^0 production, and lead to a satisfactory agreement with experimental data. The large effect for π^0 production stems mainly from the fact that in IA quite a




Figure 3. Differential cross sections for $d(\gamma, \pi^0)pn$: Dashed: IA; solid: IA+complete rescattering; Exp.: Krusche *et al.*, *Eur. Phys. J. A* **6**, 309 (1999).




Figure 4. Spin asymmetry of $d(\gamma, \pi^0)pn$: Dotted: IA; dashed: IA+NN rescattering; solid: IA+NN+ πN rescattering. Dash-dotted: results for the nucleon, for π^0 average of proton and neutron (right panel).

fraction of the coherent production is included due to the non-orthogonality of the final plane wave to the bound state. We have also evaluated the contribution to the spin asymmetry of the total cross section which determines the GDH-sum rule. While in IA the spin asymmetry for π^0 production on the deuteron is very similar to the average over neutron and proton asymmetries this is not the case for charged π production due to the Pauli principle. Moreover, final state rescattering is quite important and cannot be neglected (Fig. 4). For charged π production the effect is much larger in the asymmetry than in the total cross section and comparable to the effect in the neutral π production. For the GDH integral up to 550 MeV, we obtain from $d(\gamma, \pi)NN$ in IA $I_{d(\gamma, \pi)NN}^{(IA)} = 187 \mu b$ which is reduced to $I_{d(\gamma, \pi)NN}^{(IA+NN+\pi N)} = 87 \mu b$ by final state rescattering.