

$^{16}\text{O}+^{16}\text{O}$ molecular structure and superdeformation in ^{32}S

Shigeo Ohkubo

Department of Applied Science and Environment, Kochi Women's University,
Kochi 780-8515, Japan*Received 1 October 2002*

Abstract. The molecular structure with the $^{16}\text{O}+^{16}\text{O}$ configuration in ^{32}S and the superdeformed structure in ^{32}S are discussed from the viewpoint of the cluster model.

Keywords: cluster, $^{16}\text{O}+^{16}\text{O}$, ^{32}S , superdeformation

PACS: 21.60.Gx, 21.60.Ev, 27.30.+t,

Almost two hundreds superdeformed rotational bands have been observed in a wide range of mass regions ($A = 60, 80, 130, 150, 190$) [1]. Recently superdeformation has been observed even in the *sd*-shell region such as ^{38}Ar and ^{40}Ca [2, 3]. The superdeformed band in ^{32}S , which is a doubly magic nucleus in the deformed shell model, is very interesting because it gives "the missing link between the known region of superdeformed nuclei around ^{60}Zn and the cluster-like structures in lighter nuclei, like in ^{20}Ne and ^8Be " [1]. Yamagami and Matsuyanagi [4] calculated the superdeformed band structure in ^{32}S in the symmetry-unrestricted cranked Hartree-Fock method using the Skyrme force.

On the other hand, as for the cluster structure in nuclei, the band structure with α -cluster configuration has been clearly established for $4N$ -nuclei such as ^8Be , ^{20}Ne and ^{44}Ti [5]. In order to link the superdeformed structure with the cluster structure in ^{32}S , it is important to make clear the $^{16}\text{O} + ^{16}\text{O}$ clustering aspects in ^{32}S . The $^{16}\text{O} + ^{16}\text{O}$ structure in ^{32}S , which is an analog of the $\alpha+\alpha$ structure in ^8Be , has been most extensively studied from the viewpoint of the molecular structure. However, although many states with the $^{16}\text{O} + ^{16}\text{O}$ molecular structure have been observed, the band structure with the $^{16}\text{O} + ^{16}\text{O}$ configuration in ^{32}S has not been established.

We show that the $^{16}\text{O} + ^{16}\text{O}$ cluster bands exist in ^{32}S from a different point of view, that is, rainbow scattering. From a systematic analysis of rainbow scattering the interaction potential for the $^{16}\text{O} + ^{16}\text{O}$ system can be well determined up to very internal region. The potential with a folding-type diffuse surface can describe the elastic scattering angular distributions in a wide range of energies from $E_L =$

1120 MeV to 75 MeV. It is also shown that this potential can describe the low energy $^{16}\text{O} + ^{16}\text{O}$ elastic scattering between $E_L = 63$ MeV to 25 MeV, which means that the potential is useful for the study of the $^{16}\text{O} + ^{16}\text{O}$ structure of ^{32}S .

In order to reveal the resonant and bound structures of the $^{16}\text{O} + ^{16}\text{O}$ system, we have solved the complex scaling equation with the real part of the optical potential for the $^{16}\text{O} + ^{16}\text{O}$ system. We have obtained the rotational bands with the $^{16}\text{O} + ^{16}\text{O}$ configuration. The first Pauli-allowed band with $N = 24$ appears about 7.5 MeV below the threshold, that is 9 MeV in excitation energy from the ground state of ^{32}S . The second $N = 26$ band starts at about 3 MeV above the threshold. The $N = 28$ band starts at about 10 MeV above the threshold. The centroids of the observed molecular states in this region correspond well to the calculated $N = 28$ band. Many molecular resonances observed in experiment may be fragmented from the $N = 28$ band. There are no experimental candidates for the $N = 26$ band, which is inevitably predicted in-between the $N = 24$ and $N = 28$ bands. Therefore this band states should be searched for in experiment.

The band head of the superdeformed rotational band by Yamagami and Matsuyanagi [4] appears at about 12 MeV with the standard SIII force, which is about 4 MeV higher than the present calculation. The authors point out that the band head energy becomes about 9 MeV if the rotational zero-point corrections are taken into account [4]. Thus the superdeformed band well corresponds to our $N = 24$ cluster band in excitation energy. Also the structure change of the band occurs at $J = 24$ in Ref.[4], which is consistent with our result that due to the Pauli principle the maximum J of the lowest band is 24. The rotational constant k of the superdeformed band estimated from $J = 10 - 22$ in Ref.[4] is about 50-55 keV, which is consistent with our calculated $k = 52$ keV. Thus the superdeformed rotational band in ^{32}S corresponds well to the lowest $N = 24$ cluster band with the $^{16}\text{O} + ^{16}\text{O}$ configuration proposed in Ref.[6]. The calculated $B(E2)$ values of the $N = 24$ band is enhanced very much.

The author thanks Prof. K. Matsuyanagi for useful conversations on superdeformation.

a. E-mail: shigeo@cc.kochi-wu.ac.jp

References

1. J. Dobaczewski *Proc. Int. Conf. on Nuclear Structure '98*, Galtinburg,1998, (Ed. C. Baktash) p.135; nucl-th/981104.
2. C. E. Svensson *et al.*, Phys. Rev. Lett. **85** (2000) 2693.
3. E. Ideguchi *et al.*, Phys. Rev. Lett. **87** (2001) 222501.
4. M. Yamagami and K. Matsuyanagi, Nucl Phys. **A672** (2000) 123; nucl-th/9809038.
5. S. Ohkubo, Prog. Theor. Phys. Suppl. **132** (1998).
6. S. Ohkubo and K. Yamashita, Phys. Rev. C **66** (2002) 021301.