

ABOUT SOME RELATIVISTIC EFFECTS IN THE $A(\vec{d}, p)X$ REACTION WITH THE PROTON EMISSION IN A FORWARD DIRECTION

L.S. Azhgirey¹, N.P. Yudin²,

¹ *JINR, 141980 Dubna, Moscow Region, Russia*

² *Moscow State University, 119899 Moscow, Russia*

Abstract

In a nonrelativistic quantum theory the longitudinal and transversal components of the internal momentum in the deuteron wave function (DWF) are bound together as they have to form the well known superposition of the S- and D-waves. In a relativistic case — at least in light cone dynamics — the dependence of the DWF on the longitudinal and transversal momenta can be considerably changed, because these components turn out in large part to be bound in another way. Previously such a possibility was pointed out by Blankenbecler et al. Later Karmanov et al. have developed a quantitative approach that can be used to describe this situation. Here we present some results of calculations of the tensor analyzing power of (d, p) reaction at relativistic deuteron momenta using Karmanov's DWF. The momentum behaviour of the tensor analyzing power obtained is compared with that calculated with standard DWFs.

In recent years attempts to gain some insight into a manifestation of the structure of the relativistic deuteron in nuclear interactions are concerned mainly with an investigation of spin phenomena in reactions initiated by relativistic polarized deuterons:

$$\vec{d} + p \rightarrow p + d, \quad (1)$$

$$\vec{d} + p \rightarrow p + p + n, \quad (2)$$

$$\vec{d} + p(A) \rightarrow p + X, \quad (3)$$

where detected protons are emitted in forward directions in the lab. syst. The simplest mechanism of these processes can be represented schematically by the diagram shown in Fig. 1. When protons are detected at non-zero angles, they may have a rather high transversal component of the momentum. At usually considered mechanisms these reactions are concerned with the deuteron wave function.

Already upon the first measurements of the tensor analyzing power T_{20} of reaction (3) at 9 GeV/c with the emission of protons at 0° [1-5] a significant discrepancy between the values calculated in the relativistic impulse approximation and experimental ones has come to light. A comparison of currently available experimental data on $A_{yy}(0^\circ)$ (recall, that at 0° $A_{yy} = -T_{20}/\sqrt{2}$) with calculations in the frame of the impulse approximation with the

standard DWFs [6, 7] is made in fig. 2. More recently, on the measurement of the tensor analyzing power A_{yy} of reaction (3) at 9 GeV/c with the emission of protons with large transverse momenta [8], this discrepancy has been compounded.




Fig. 1. The simplest diagram to describe reactions initiated by relativistic deuterons with proton emission in a forward direction. Along with the one-nucleon exchange (N) there also the exchange of baryon resonances (R) can take place [11].

The discrepancy between the theory and experiment in principle can be overcome by the addition of a P-wave into the deuteron wave function (DWF). True enough, the P-wave is conditioned by the different mechanisms in a number of approaches. In the paper [9] the data on T_{20} at 0° have been successfully described taking account of the P-wave arising because of the production of six-quark configuration that gives odd parity resonances at their fragmentation to the baryon channel. At the same time the calculations on the basis of the Bethe-Salpeter equation give the P-wave admixture that is inadequate to eliminate the discrepancy with experiment [10]. In relation to the reaction (1) it is notable that taking account of an admixture of the odd parity baryon resonances to the DWF allows the T_{20} data at 0° to be better described than within the usual one-nucleon exchange approximation [11].

In this report we would like to point out one more factor related to the wave function of a fast moving deuteron. These are the rotational properties of the relativistic DWF in the light cone (LC) dynamics. [12]. Let us consider shortly the calculations in the frame of this approach.

A general expression for the parameter T_{20} has the form

$$T_{20} = \frac{\sum_{M, M'} Sp\{\psi_M t_{20} \psi_{M'}^\dagger\}}{(1/3)Sp\{\psi_M \psi_{M'}^\dagger\}}, \quad (4)$$

where the operator t_{20} is defined by expression

$$\langle m' | t_{20} | m \rangle = (-1)^{j-m'} \langle 1 \ m \ 1 \ -m' | 2 \ 0 \rangle, \quad (5)$$

ψ_M is DWF with spin $J = 1$ and its projections on the quantization axis $M = 0, \pm 1$, ψ^\dagger is the Hermitian conjugate function, and $\langle 1 \ m \ 1 \ -m' | 2 \ 0 \rangle$ is a Clebsh-Gordan coefficient.

In order to calculate T_{20} we should know ψ_M with different projections M . However ψ_M in LC dynamics has several unusual properties. The first one consists in "strange" relations of ψ_M with different M . Let us suppose that we have ψ_M with $M = 0$. Then ψ_M with $M = +1$ should be obtained from ψ_M with $M = 0$ by the action of the standard combination of operators J_x, J_y :

$$\psi_{M=1} \sim (J_x + iJ_y)\psi_{M=0}. \quad (6)$$

But in LC dynamics these operators of angular momentum are dependent on the interaction, and to calculate the sum in the expression for T_{20} the Schrödinger equation for all components of ψ_M has to be solved independently. Of course it is an objectionable problem. The dependence of angular momentum on interaction involves difficulties in forming states with definite angular momentum.

The second complication centres around the dependence of the wave function on variables. Indeed, the wave function of two particles is in general dependent on two variables,

$$\psi = \psi(k_l, k_T), \quad (7)$$

where k_l and k_T are longitudinal and transversal components of momentum, respectively. In the nonrelativistic physics k_l and k_T are rigidly bound together by the necessity to form the usual superposition of S - and D -waves in DWF, with the result that the nonrelativistic DWF is dependent on the only variable k . In LC dynamics this relation between k_l and k_T may be much weaker, and in principle DWF can be dependent on two variables. Attention to such a situation was first called by Schmidt and Blankenbecler [13].

These difficulties of the LC-dynamics were to a great extent overcome by Karmanov et al. [14, 15]. At first sight the approach of Karmanov complicates the situation since there is introduced an additional dynamical variable ω^μ — a four-vector of the normal to the chosen light cone surface. But this new variable permits the necessity dealing with operators of the angular momentum dependent on the interaction to be excluded: it turns out that the rotation of the light cone surface is equivalent to the interaction dependent rotation of the system. Therefore states with the definite angular momentum become formally independent on the interaction.

The dependence on the interaction does not completely disappear, it arises in another context: there are too many states with the same angular momentum and the role of the interaction now is to choose the right combination among these states. But this new problem is much more easier than the problem of solving the Schrödinger equation. The difficulties existing in Karmanov's approach were overcome in [14, 15], where the parameters of the relativistic DWF were given. This function was used in our calculations of the tensor analysing power A_{yy} of the deuteron breakup reaction (3).

The DWF, now the function of nucleon spins, an internal deuteron momentum, and an orientation of the quantization plane, has the form:

$$\Psi_{\sigma_2\sigma_1}^M = w_{\sigma_2}^* \psi^M(\vec{k}, \vec{n}) \sigma_y w_{\sigma_1}, \quad (8)$$

where

$$\begin{aligned} \vec{\psi}(\vec{k}, \vec{n}) = & f_1 \frac{1}{\sqrt{2}} \vec{\sigma} + f_2 \frac{1}{2} \left[\frac{3}{k^2} \vec{k} (\vec{k} \vec{n} - \vec{\sigma}) + f_3 \frac{1}{2} [3\vec{n}(\vec{n} \vec{\sigma}) - \vec{\sigma}] + \right. \\ & \left. f_4 \frac{1}{2k} (3\vec{k}(\vec{n} \vec{\sigma}) + 3\vec{n}(\vec{k} \vec{\sigma}) - 2(\vec{k} \vec{n}) \vec{\sigma}) + f_5 \sqrt{\frac{3}{2}} \frac{i}{k} (\vec{k} \times \vec{n}) + f_6 \frac{\sqrt{3}}{2k} ((\vec{k} \times \vec{n}) \times \vec{\sigma}). \right. \end{aligned} \quad (9)$$

Here \vec{n} is the normal to the light front surface, \vec{k} is the momentum of nucleons in deuteron in their rest frame, $\vec{\sigma}$ are the Pauli matrices, $w_{\sigma_1(\sigma_2)}$ are the spin functions of nonrelativistic nucleons, and $f_{1,\dots,6}$ are the invariant about rotations functions of the kinematical variables, that define the deuteron state.




Fig. 2. Tensor analyzing power A_{yy} of the deuteron breakup with the emission of protons at 0° vs internal momentum k . Experimental data: triangles — [1, 2], squares — [3], diamonds — [4], empty circles — [5] (H target), full circles — [5] (C target). Calculations are made in relativistic impulse approximation with Paris [6] (broken curve), Bonn [7] (dotted curve) and Karmanov's [14] DWFs.

Previously we described $dp \rightarrow pX$ and $dp \rightarrow pd$ reactions [11] in the coordinate system where the z axis was aligned with the beam direction. For the wave function (8), too, let us assume that \vec{n} is directed along z axis, i.e. $\vec{n} = (0, 0, 1)$.

The expression (4) may be written in the form:

$$T_{20} = \sqrt{\frac{3}{2}} \frac{[-3Sp(\psi_z\psi_z^\dagger) + Sp(\vec{\psi}\vec{\psi}^\dagger)]}{Sp(\vec{\psi}\vec{\psi}^\dagger)}, \quad (10)$$

where we went from spherical notations to the vector ones. For the traces we have:

$$\begin{aligned} Sp(\vec{\psi}\vec{\psi}^\dagger) &= 3[f_1^2 + f_2^2 + (3z^2 - 1)f_2f_3 + f_3^2 + 4zf_4(f_2 + f_3) + \\ &\quad f_4^2(z^2 + 3) + (1 - z^2)f_5^2 + (1 - z^2)f_6^2], \\ Sp(\psi_z\psi_z^\dagger) &= f_1^2 + \sqrt{2}[3(kz)^2 - 1]f_1f_2 + 2\sqrt{2}f_1f_3 + 4\sqrt{2}kzf_1f_4 + \\ &\quad \frac{1}{2}[3(kz)^2 + 1]f_2^2 + 2[3(kz)^2 - 1]f_2f_3 + kz[3(kz)^2 + 5]f_2f_4 + \\ &\quad 2f_3^2 + 8kzf_3f_4 + 3\sqrt{3}kz[1 - (kz)^2]f_2f_6 + 3\sqrt{3}[1 - (kz)^2]f_4f_6 + \\ &\quad \left\{ \frac{9}{2}[(kz)^2 + 1] - (kz)^2 \right\} f_4^2 + \frac{3}{2}[1 - (kz)^2]f_6^2, \end{aligned} \quad (11)$$

where $z = \cos\theta$.

The results of calculations of $A_{yy}(0^\circ)$ with the wave function (9) are shown in fig. 2 by the solid curve. It is seen that as opposed to the calculations with the standard nonrelativistic DWFs [6, 7] the solid curve does not cross the horizontal axes and is in reasonable good agreement with experimental data in the region of k from 0.4 to 0.8 GeV/c. This result in our opinion is due to the fact that Karmanov's model establishes a new link between k_l and k_T that is different from those of the S- and D-wave superposition in nonrelativistic DWF. Similar effect was discussed in [15] on the example of Wick-Cutcosky model, where it was shown in the clear form that a S-wave two-particle system becomes dependent on the angle in the LC dynamics. In other words it is a manifestation of the intimate connection between the internal motion and the motion of the system as a whole.

This work was supported in part by the Russian Foundation for Fundamental Research (grant No. 01-02-17299).

References

- [1] C.F.Perdrisat *et al.*, Phys.Rev.Lett. **59**, 2840 (1987).
- [2] V.Punjabi *et al.*, Phys.Rev. **C39**, 608 (1989).
- [3] V.G.Ableev *et al.*, JINR Rapid Comm. No.4[43]-90, 5 (1990).
- [4] T.Aono *et al.*, Phys.Rev.Lett. **74**, 4997 (1995).
- [5] L.S. Azhgirey *et al.*, Phys.Lett. **B387**, 37 (1996).
- [6] M.Lacombe *et al.*, Phys.Rev. **C21**, 861 (1980).
- [7] R.Machleidt *et al.*, Phys.Rep. **149**, 1 (1987).
- [8] S.V.Afanasiev *et al.*, Phys.Lett. **B434**, 21 (1998).
- [9] A.P.Kobushkin, Phys.Lett. **B421**, 53 (1998).

- [10] L.Kaptari *et al.*, Phys.Lett. **B351**, 400 (1995).
- [11] L.S.Azhgirey and N.P.Yudin, Yad.Fiz. **63**, 2280 (2000) (Phys.Atom.Nucl. **63**, 2184 (2000)).
- [12] P.A.M. Dirac, Rev.Mod.Phys. **21**, 392 (1949);
S.Weinberg, Phys.Rev. **150**, 1313 (1966);
L.L.Frankfurt and M.I.Strikman, Phys.Rep. **76**, 215 (1981).
- [13] I.A.Schmidt and R.Blankenbecler, Phys.Rev. **D15**, 3321 (1977);
Ch.-Y.Wong and R.Blankenbecler, Phys.Rev. **C22**, 2431 (1980);
M.Chemtob *et al.*, Nucl.Phys. **A314**, 387 (1979).
- [14] V.A.Karmanov and A.V.Smirnov, Nucl. Phys. A **575**, 520 (1994);
J.Carbonell *et al.*, Phys.Rep. **300**, 215 (1998).
- [15] J.Carbonell and V.A.Karmanov, Nucl. Phys. A **581**, 625 (1994).