Isovector deformation and its link to the neutron shell closure

Dao T. Khoa

Institute for Nuclear Science & Technique, VAEC, P.O. Box 5T-160, Nghia Do, Hanoi, Vietnam e-mail: khoa@vaec.gov.vn

Received: date / Revised version: date

Abstract. DWBA analysis of the inelastic ${}^{30-40}S(p, p')$ and ${}^{18-22}O(p, p')$ scattering data measured in the inverse kinematics has been performed to determine the isoscalar (δ_0) and isovector (δ_1) deformation lengths of the 2_1^+ excitations in the Sulfur and Oxygen isotopes using a compact folding approach. A systematic N-dependence of δ_0 and δ_1 has been established which shows a link between δ_1 and the neutron-shell closure. Strong isovector deformations were found in several cases, e.g., the 2_1^+ state in ${}^{20}O$ where δ_1 is nearly three times larger than δ_0 . These results confirm the relation $\delta_1 > \delta_0$ anticipated from the core polarization by the valence neutrons in the open-shell (neutron rich) nuclei. The effect of neutron shell closure at N = 14 or 16 has been discussed based on the folding model analysis of the inelastic ${}^{22}O+p$ scattering data at 46.6 MeV/u measured recently at GANIL.

PACS. 2 5.40.Ep, 21.10.Re, 24.10.Eq, 24.10.Ht

The neutron and proton contributions to the structure of the lowest 2⁺ excited states are known to be quite different in the neutron-rich nuclei due, in particular, to a strong polarization of the core by valence neutrons [1]. In the distorted-wave Born approximation (DWBA) for the inelastic hadron scattering, the different neutron and proton contributions to the nuclear excitation are explicitly determined by the isospin dependence of the inelastic form factor (FF).

In general, the isospin-dependent nucleon optical potential (OP) can be written in terms of the isoscalar (IS) and isovector (IV) components [2] as

$$U(R) = U_0(R) \pm \varepsilon U_1(R), \ \varepsilon = (N - Z)/A, \tag{1}$$

where the + sign pertains to incident neutron and - sign to incident proton. While the strength of the Lane potential U_1 has been studied since a long time [2] in the (p, p) and (n, n) elastic scattering and (p, n) reaction studies, very few attempts were made to study the isospin dependence of the inelastic FF. Within a collective-model prescription, the inelastic FF for the nucleon-nucleus scattering is obtained by "deforming" the OP (1) with scaling factors δ known as the nuclear deformation lengths

$$F(R) = \delta \frac{dU(R)}{dR} = \delta_0 \frac{dU_0(R)}{dR} \pm \varepsilon \ \delta_1 \frac{dU_1(R)}{dR}.$$
 (2)

The explicit knowledge of the isoscalar (δ_0) and, especially, isovector (δ_1) deformation lengths would give us vital information on the structure of the nuclear excitation under study. There are only two types of experiment that might allow one to determine the IV deformation length δ_1 based on the prescription (2): i) (p, n) reaction leading to the *excited* isobar analog state. It was shown [3], however, that the two-step mechanism usually dominates this process and the calculated DWBA cross sections are not sensitive to δ_1 .

ii) Another way is to extract δ_0 and δ_1 from the (p, p')and (n, n') inelastic scattering measured at the same incident energy and exciting the same target state [3,4]. Such double measurements are presently not feasible with the beams of unstable nuclei.

We have recently suggested a compact folding method [5] to determine $\delta_{0(1)}$ based on the DWBA analysis of the (p, p') data only. In this approach, instead of deforming the OP, we build up the proton and neutron transition densities of a 2^{λ} -pole excitation ($\lambda \geq 2$) by using the Bohr-Mottelson prescription [6] separately for protons and neutrons

$$\rho_{\lambda}^{\tau}(r) = -\delta_{\tau} \frac{d\rho_{\text{g.s.}}^{\tau}(r)}{dr}, \text{ with } \tau = p, n.$$
(3)

Here $\rho_{\text{g.s.}}^{\tau}(r)$ are the proton and neutron ground-state (g.s.) densities and δ_{τ} are the corresponding deformation lengths. Given the explicit proton and neutron transition densities, one can obtain from the folding model [7] the inelastic proton-nucleus FF in terms of the IS and IV parts as

$$F(R) = F_0(R) - \varepsilon F_1(R), \qquad (4)$$

where $F_0(R)$ and $F_1(R)$ are determined [7] from the sum $(\rho_{\lambda}^n + \rho_{\lambda}^p)$ and difference $(\rho_{\lambda}^n - \rho_{\lambda}^p)$ of the neutron and proton transition densities (3), respectively. One can see that $F_1(R)$ is just the prototype of the Lane potential in the inelastic nucleon scattering. It is natural to represent

Ì

the IS and IV parts of the nuclear transition density as

$$\rho_{\lambda}^{0(1)}(r) = \rho_{\lambda}^{n}(r) \pm \rho_{\lambda}^{p}(r).$$
(5)

On the other hand, $\rho_{\lambda}^{0(1)}(r)$ can be obtained using the same Bohr-Mottelson method, by deforming the IS and IV parts of the g.s. density

$$\rho_{\lambda}^{0(1)}(r) = -\delta_{0(1)} \frac{d[\rho_{\text{g.s.}}^{n}(r) \pm \rho_{\text{g.s.}}^{p}(r)]}{dr}.$$
 (6)

It is straightforward to derive in this compact approach [5] a consistent one-to-one correspondence between $\delta_{p(n)}$ and $\delta_{0(1)}$, which can be used to determine $\delta_{0(1)}$ values from $\delta_{p(n)}$ values given by the DWBA analysis. If one assumes that the excitation is purely *isoscalar* and the neutron and proton densities have the same radial shape, scaled by the ratio N/Z, then $\delta_n = \delta_p = \delta_0 = \delta_1$. Therefore, any significant difference between δ_0 and δ_1 would directly indicate a different isospin distribution in the structure of the nuclear excitation under study.

In the present work, we have studied the elastic and inelastic 30,32 S+p scattering data at 53 MeV/u [8] and 34,36,38,40 S+p data [9,10,11,12] at energies of 28 to 39 MeV/u. The IS and IV contributions of the inelastic FF were considered explicitly to find out a systematic behavior of δ_1 along the Sulfur isotopic chain, passing by the magic number N = 20. Then, the folding + DWBA analysis of the elastic and inelastic 18,20 O+p data at 43 MeV/u [13] and 22 O+p data at 46.6 MeV/u [14] has been done to find out the N-dependence of δ_1 in the Oxygen case.

To have the accurate "distorted" waves for the DWBA calculation of inelastic scattering, the optical model (OM) analysis of the elastic data was done using the real folded potential [7] obtained with the density- and isospin dependent CDM3Y6 interaction [15] and nuclear g.s. densities given by the Hartree-Fock-Bogoliubov (HFB) calculation [16]. The imaginary part of the OP was taken in the Woods-Saxon (WS) form from the global systematics CH89 [17]. All the considered elastic data were well reproduced (see Figs. 2 and 4) with the depth of the WS imaginary potential slightly adjusted by the OM fit (keeping the radius and diffuseness unchanged). The experimental reduced electric transition rate $B(E2\uparrow)$ was used in each case to fix δ_p value in the expression (3) for $\rho_2^p(r)$ which is further used in the folding calculation [5,7]. As the only parameter, δ_n was adjusted iteratively in the folding + DWBA calculation to fit the measured inelastic cross section. Since the CDM3Y6 interaction is real, the imaginary nuclear FF was obtained by deforming the imaginary part of the OP with δ_0 and δ_1 values at each iteration step of the folding + DWBA fit to the inelastic data. In each case, the final set of deformation lengths δ_0 and δ_1 was fixed only after the best-fit δ_n has been obtained.

An earlier folding + DWBA analysis [7] of the same inelastic ${}^{30-40}$ S+p scattering data, using inelastic FF given by the microscopic transition densities obtained in the Quasiparticle Random Phase Approximation (QRPA) [8], has shown that the neutron and proton contributions to the 2_1^+ excitation in 30,32,34 S follow approximately the



Fig. 1. The measured $B(E2 \uparrow)$ transition strength (upper part), and the extracted isoscalar and isovector deformation lengths (lower part) of 2_1^+ states in Sulfur isotopes. The lines are to guide the eye.

isoscalar rule [5] which implies $\delta_0 \approx \delta_1$. The present folding model analysis using the collective-model transition densities (3) has shown about the same results (see δ_0 and δ_1 values extracted for 30,32,34 S in Fig. 1). With the neutron shell becomes closed at N = 20, a significant "damping" of the neutron transition strength occurs and suppresses strongly the IV deformation length δ_1 of the 2_1^+ state of ³⁶S. In fact, δ_1 is reaching its minimum as the neutron number N approaches the magic number 20. The N-dependence of the IV deformation length δ_1 is well correlated with the N-dependence of the reduced transition rate $B(E2\uparrow)$ which also reaches its minimum at N=20(see Fig. 1). The shell closure effect is so strong in this case that the proton and neutron QRPA transition densities needed to be scaled down by a factor of 0.63 and 0.88, respectively, for a correct description of the measured 2^+ cross section in our earlier DWBA analysis [7] of the inelastic ${}^{36}S+p$ scattering. The contribution by the two valence neutrons in ${}^{38}S$ to the 2^+_1 excitation is quite strong and δ_n turned out to be larger than δ_p by around 30% (see Fig. 2). This difference between the proton and neutron transition strengths results on the IV deformation length δ_1 larger than δ_0 by about 64% (see Fig. 1). Note that such a strong core polarization by the two valence neutrons in the ${}^{38}S$ case could not be fully accounted for by the QRPA calculation and the QRPA neutron transition density has been scaled by a factor of 1.25 for a good agreement of calculated DWBA cross section with



Fig. 2. Elastic and inelastic ³⁸S+p scattering data at 39 MeV/u [11] in comparison with the DWBA cross sections given by the folded FF. The dashed curve is inelastic cross section obtained in the isoscalar limit ($\delta_n = \delta_p = \delta_0 = \delta_1$).

the data [7]. Since the inelastic ${}^{40}\text{S}+p$ scattering data at 30 MeV/u [12] consist of only two data points with significant error bars, the extracted $\delta_{0(1)}$ values for ${}^{40}\text{S}$ might not be as reliable as those extracted for ${}^{30-38}\text{S}$. The new measurement is, therefore, needed for an accurate estimate of the IV deformation length of 2^+_1 state of ${}^{40}\text{S}$.

The present analysis of the inelastic ${}^{18,20,22}\text{O}+p$ scattering data has shown an interesting N-dependence of δ_1 . Except for the double-magic $(N = Z)^{16}$ O, where one has exact relation $\delta_0 = \delta_1$, the results obtained for ¹⁸O already indicate a rather strong IV deformation for the 2_1^+ state of this nucleus (see Fig. 3). We recall that the IS and IV deformation parameters of the 2^+_1 state of ${}^{18}O$ have been determined long ago by Grabmayr $et \ al. \ [4]$ in a simultaneous DWBA analysis of the (p, p') and (n, n')inelastic scattering data at 24 MeV using the collective form factor (2). It is easy to deduce from the results of Ref. [4] the corresponding deformation lengths $\delta_0 \approx 1.1$ and $\delta_1 \approx 2.6 \pm 1.3$ fm for the 2_1^+ state of ¹⁸O. These values agree reasonably with the results of our folding + DWBA analysis (see Fig. 3) of the inelastic ${}^{18}\text{O}+p$ scattering data at 43 MeV/u [13]. Note that the δ_1 value extracted from our analysis has a much smaller uncertainty compared to that deduced from the results of Ref. [4]. The error bars for $\delta_{0(1)}$ plotted in Figs. 1 and 3 were accumulated from the experimental uncertainties of the measured B(E2) values and (p, p') cross section.



Fig. 3. The measured $B(E2 \uparrow)$ transition strength (upper part) and the extracted IS and IV deformation lengths (lower part) of 2_1^+ states in the Oxygen isotopes. The values for ²⁴O are deduced from the QRPA prediction by Khan *et al.* [19]. The lines are to guide the eye.

As already mentioned above, a double measurement of (p, p') and (n, n') inelastic scattering is not possible with unstable Oxygen isotopes, and our folding method is the only alternative way to determine δ_1 . The present analysis (with a more consistent treatment of the IV part of the imaginary FF) has confirmed again the large IV deformation length of the 2^+_1 state of ²⁰O found earlier in Ref. [5]. With the IV deformation about three times the IS deformation (see Fig. 3), the contribution by the Lane form factor F_1 to the 2^+_1 excitation of ²⁰O amounts up to 40-50% of the total inelastic cross section [5]. If we consider ²⁰O as consisting of the ¹⁶O (or ¹⁸O) core and four (or two) valence neutrons, then a large value of IV deformation length δ_1 indicates a strong core polarization by the valence neutrons in the 2^+_1 state of ²⁰O.

In such a "core + valence neutrons" picture, it is natural to expect that the 2_1^+ state of ²²O should be more collective and have a larger IV deformation length due to the contribution of two more valence neutrons. However, the inelastic ²²O+p scattering data at 46.6 MeV/u measured recently at GANIL [14] show clearly the opposite effect, with the (p, p') cross section about 3 to 4 times smaller than that measured for the 2_1^+ state of ²⁰O at 43 MeV [13] over a wide angular range. The folding + DWBA analysis [14] of these data using the QRPA transition densities for the 2_1^+ state of ²²O has pointed to a much weaker neutron transition strength compared to that of the 2_1^+ state of

²⁰O. Given a significantly higher excitation energy of this state (1.5 MeV higher than that of the 2_1^+ state of ²⁰O), the newly measured inelastic ²²O+*p* scattering data were suggested [14] as an important evidence for the neutron shell closure at N = 14 or 16.



Fig. 4. The same as Fig. 2 but for the elastic and inelastic ${}^{22}\text{O}+p$ scattering data at 46.6 MeV/u [14].

With the proton deformation length δ_p fixed by the adopted $B(E2 \uparrow)$ value for the 2^+_1 state of 22 O [18], the present analysis has given the upper limit for the neutron deformation length as $\delta_n \leq 1.2 \ \delta_p$ (see Fig. 4), which implies $\delta_1 \leq 1.1$ fm. As a result, the extracted δ_1 value is just slightly larger than δ_0 and the N-dependence shown in Fig. 4 indicates that δ_1 reaches its minimum at either N = 14 or 16. Based on the results obtained above for the Sulfur isotopes, we conclude that the deduced Ndependence of δ_1 for the 2^+_1 states of Oxygen isotopes suggests the neutron shell closure at either N = 14 or 16. We note further that the $B(E2 \uparrow)$ value predicted by the QRPA calculation [19] for the 2_1^+ state of ²⁴O is even smaller than that adopted for the 2^+_1 state of ${}^{22}O$, a fact which could favor the shell closure at N = 16. We have also deduced δ_0 and δ_1 of the 2^+_1 state of ^{24}O from these QRPA results and they are nearly the same as those deduced from the (p, p') data in ²²O case. If we take into consideration 4 MeV gap between the $2s_{1/2}$ and $1d_{3/2}$ subshells predicted recently by a consistent HFB calculation [20], then it is also likely that the neutron shell closure occurs at N = 16. In any case, more measurements for

²⁴O are highly desirable for a definitive conclusion on a new magic number N = 16 in the neutron rich nuclei.

In summary, we have shown that the behavior of the dynamic isovector deformation of the 2^+_1 states in neutron rich nuclei is closely correlated with the evolution of the valence neutron shell. This interesting result emphasizes again the importance of (p, p') reactions measured with unstable nuclei in the inverse kinematics.

Acknowledgement

This research project has been supported, in part, by the Natural Science Council of Vietnam, EU Asia-Link Program CN/Asia-Link/008 (94791) and Vietnam Atomic Energy Commission (VAEC).

References

- A.M. Bernstein, V.R. Brown, V.A. Madsen, Comments Nucl. Part. Phys. 11, 203 (1983).
- 2. A.M. Lane, Phys. Rev. Lett. 8, 171 (1962).
- R.W. Finlay, J. Rapaport, V.R. Brown, V.A. Madsen, J.R. Comfort, Phys. Lett. 84B, 169 (1979).
- P. Grabmayr, J. Rapaport, R.W. Finlay, Nucl. Phys. A 350, 167 (1980).
- 5. D.T. Khoa, Phys. Rev. C 68, 011601(R) (2003).
- A. Bohr, B.R. Mottelson, Nuclear Structure (Benjamin, New York, 1975), Vol. 2.
- D.T. Khoa, E. Khan, G. Colò, N. Van Giai, Nucl. Phys. A 706, 61 (2002).
- 8. E. Khan et al., Nucl. Phys. A 694, 103 (2001).
- R. Alarcon, J. Rapaport, R.T. Kovzes, W.H. Moore, B.A. Brown, Phys. Rev. C **31**, 697 (1985).
- A. Hogenbirk, H.P. Blok, M.G.E. Brand, A.G.M. van Hees, J.F.A. van Hienen, F.A. Jansen, Nucl. Phys. A 516, 205 (1990).
- 11. J.H. Kelley et al., Phys. Rev. C 56, R1206 (1997).
- 12. F. Maréchal et al., Phys. Rev. C 60, 034615 (1999).
- 13. E. Khan et al., Phys. Lett. B 490, 45 (2000).
- 14. E. Becheva et al., Phys. Rev. Lett. 96, 012501 (2006).
- D.T. Khoa, G.R. Satchler, W. von Oertzen, Phys. Rev. C 56, 954 (1997).
- M. Grasso, N. Sandulescu, N. Van Giai, R.J. Liotta, Phys. Rev. C 64, 064321 (2001).
- R.L. Varner, W.J. Thompson, T.L. McAbee, E.J. Ludwig, T.B. Clegg, Phys. Rep. 201, 57 (1991).
- S. Raman, C. W. Nestor, Jr., P. Tikkanen, At. Data and Nucl. Data Tables 78, 1 (2001).
- E. Khan, N. Sandulescu, M. Grasso, N. Van Giai, Phys. Rev. C 66, 024309 (2002).
- 20. A. Obertelli, S. Peru, J.-P. Delaroche, A. Gillibert, M. Girod, H. Goutte, Phys. Rev. C 71, 024304 (2005).