

SNUTP 97-033, UTHEP-358

March., 1997

**Comment on: “Meson Masses in Nuclear Matter”**

Tetsuo Hatsuda

*Institute of Physics, University of Tsukuba, Tsukuba, Ibaraki 305, Japan*

Su Hounng Lee

*Department of Physics, Yonsei University, Seoul, 120-749, Korea*

21.65.+f, 12.40.Vv, 14.40.Cs, 24.85.+p

arXiv:nucl-th/9703022v1 11 Mar 1997

Typeset using REVTeX

In a recent letter [1], Eletsky and Ioffe estimated the energy shift of  $\rho$ -meson with *high* momentum ( $\mathbf{q}$ ) in nuclei using the Glauber approximation. They also made remarks on the *low*  $\mathbf{q}$  vector mesons in medium. In this Comment, we will discuss the second point and show that (i) mean-field physics for the mass shift of low- $\mathbf{q}$  mesons is overlooked in [1], and (ii) the *short distant* operator product expansion (OPE) is relevant to analyze the low- $\mathbf{q}$  mesons in QCD sum rules contrary to the claim in [1].

Mesons in nuclei have considerable multiple scattering when  $|\mathbf{q}|^{-1} \gg d$  ( $d$  is the inter-nucleon distance). The relevant quantity here is known to be a polarization function (generalized optical potential)  $\Pi(\omega, \mathbf{q})$  [2], which is written, up to the Lorentz-Lorenz correction, as

$$\Pi(\omega, \mathbf{q}) = -16\pi \int \frac{d^3\mathbf{p}}{(2\pi)^3} F(\omega, \mathbf{q}; \mathbf{p}) n_F(\mathbf{p}), \quad (1)$$

where  $n_F$  is the fermi distribution,  $F$  is an *in-medium* forward scattering amplitude of an incoming vector-meson (V) with a nucleon (N) of momentum  $\mathbf{p}$ . At extremely low density, eq.(1) reduces to  $\Pi(\omega, \mathbf{q}) = -4\pi F(\omega, \mathbf{q}; 0)\rho$ , which does not however have practical use, since  $F$  has not only a smooth  $\mathbf{p}$ -dependence (from e.g. the  $t$ -channel meson exchanges considered in [3]) but also a possible rapid  $\mathbf{p}$ -dependence from  $s$ -channel baryon-resonances coupled to the  $V$ - $N$  system. In other words, the averaging of  $F$  over the fermi sea is essential for the mass shift at low  $\mathbf{q}$ . Note also that low- $\mathbf{q}$  mesons probe large distances and hence the averaged nuclear properties. The mean-field (MF) + RPA treatment of  $\Pi$ , which has been employed in hadronic model calculations (see e.g. [4]), is most suited for studying these features. The importance of this MF description for low- $\mathbf{q}$  mesons is overlooked in [1].

The MF description is also compatible with the finite-density QCD sum rules [5]. Consider the vector-current correlation in medium,

$$D_{\mu\nu}(Q^2, Q^2/q \cdot P) = i \int d^4x e^{iqx} \langle P | T [J_\mu(x) J_\nu(0)] | P \rangle, \quad (2)$$

with  $Q^2 \equiv -q^2$  and  $|P\rangle$  being the nuclear ground state with momentum  $P^\mu$ . At large  $Q^2$ , one can make OPE in two ways [6]: the short-distance (SD) expansion where  $(Q^2, Q^2/(q \cdot P)^2) \rightarrow$

( $\infty$ , finite), and the light-cone (LC) expansion where  $(Q^2, Q^2/q \cdot P) \rightarrow (\infty, \text{finite})$ . Both are at best an asymptotic expansion in  $1/Q^2$  with different composite operators contributing at each order: The SD (LC) expansion is dictated by the dimension (twist) of the operators. Since we are interested in  $\mathbf{q} \simeq 0$  with  $\mathbf{P} = 0$ , OPE in the deep-euclidian region ( $-\omega^2 \rightarrow \infty$ ) is obtained from the SD expansion by re-expanding  $1/Q^2$  and  $(q \cdot P)^2/Q^2$  in terms of  $1/\omega^2$ , which is a natural generalization of the QCD sum rules in the vacuum [5,8]. More importantly, from the analytic properties of  $D_{\mu\nu}$ , one can relate the OPE to the spectral density via the well-known fixed  $\mathbf{q}$  dispersion relation [7]

$$\text{Re } D(\omega^2 < 0, \mathbf{q}) = \int_0^\infty dx^2 \frac{\rho(x, \mathbf{q})}{(x^2 - \omega^2)} + (\text{subtraction}). \quad (3)$$

As in the vacuum sum rules, we keep only the contributions up to dimension 6 operators. The actual expansion parameter at low density is  $\Lambda_{QCD}^l p_F^m |\mathbf{q}|^n / \omega^{l+n+m}$  ( $p_F$  is the fermi momentum). Density dependent contributions from dimension 4 and dimension 6 operators have been shown not to spoil the asymptotic nature of the series [5]. In the work by Drukarev and Levin [9], a kinematics  $Q^2 \rightarrow \infty$  with  $s = (q + P/A)^2$  fixed ( $A$  is the number of nucleons) or its variant are adopted. This leads to the LC expansion as quoted in [1]. However, a dispersion relation for  $Q^2$  with fixed  $s$  is *assumed* in [9]. Such a dispersion relation, unlike eq.(3), cannot be derived from analytic properties of  $D_{\mu\nu}$  alone [8]. Also fixing  $s$  inevitably gives a large  $\mathbf{q}$  limit, which is inappropriate to extract the MF physics we are interested in.

## REFERENCES

- [1] V. L. Eletsky and B. L. Ioffe, Phys. Rev. Lett. **78**, 1010 (1997).
- [2] A. B. Migdal, in *Mesons in Nuclei*, ed. M. Rho and D. Wilkinson (North-Holland, Amsterdam, 1979). T. Ericson and W. Weise, *Pions and Nuclei*, (Oxford Univ. Press, Oxford, 1988).
- [3] B. Friman and M. Soyeur, Nucl. Phys. **A600**, 477 (1996).
- [4] H. C. Jean, J. Piekarewicz and A. G. Williams, Phys. Rev. **C49**, 1981 (1994).
- [5] T. Hatsuda and Su H. Lee, Phys. Rev. **C46**, R34 (1992). T. Hatsuda, Su H. Lee and H. Shiomi, Phys. Rev. **C52**, 3364 (1995).
- [6] T. Muta, *Foundations of Quantum Chromodynamics*, (World-Scientific, Singapore, 1987).
- [7] A. L. Fetter and J. D. Walecka, *Quantum Theory of Many-Particle Systems*, (McGraw-Hill, New York, 1971).
- [8] T. D. Cohen, R.J. Furnstahl, D.K. Griegel and X.M. Jin, Prog. in Part. Nucl. Phys. **35**, 221 (1995), section 3.7. D. K. Griegel, PhD Thesis (Univ. of Maryland) 1991.
- [9] E.G. Drukarev and E.M. Levin, Prog. Part. Nucl. Phys. **27**, 77 (1991).