

Multidimensional Pattern Formation Has an Infinite Number of Constants of Motion

Mark B. Mineev–Weinstein

Courant Institute of Mathematical Sciences
251 Mercer Street
New York University
New York, NY 10012
and
Center for Nonlinear Studies, MS–B258
Los Alamos National Laboratory
Los Alamos, NM 87545

November 14, 2018

Abstract

Extending our previous work on 2D growth for the Laplace equation we study here *multidimensional* growth for *arbitrary elliptic* equations, describing inhomogeneous and anisotropic pattern formations processes. We find that these nonlinear processes are governed by an infinite number of conservation laws. Moreover, in many cases *all dynamics of the interface can be reduced to the linear time-dependence of only one “moment” M_0* which corresponds to the changing volume while *all higher moments, M_1 , are constant in time. These moments have a purely geometrical nature*, and thus carry information about the moving shape. These conserved quantities (eqs. (7) and (8) of this article) are interpreted as coefficients of the multipole expansion of the Newtonian potential created by the mass uniformly occupying the domain enclosing the moving interface. Thus the question of how to recover the moving shape using these conserved quantities is reduced to the classical inverse potential problem of reconstructing the shape of a body from its exterior gravitational potential. Our results also suggest the possibility of controlling a moving interface by appropriate varying the location and strength of sources and sinks.

PACS numbers: 47.15. Hg, 68.10.-m, 68.70.+w, 47.20. Hw.

Many seemingly different pattern formation processes have much in common, both in their mathematical description and in their physical behavior. Among them are the famous Stefan problem (freezing of liquid), flow through porous media, the Rayleigh–Taylor instability, electrodeposition of metals, corrosion, combustion, growth of bacterial colonies, dynamics of earth cracks, diffusion–limited aggregation (DLA), etc. The common feature shared by these processes is the existence of an evolving interface. The problem of the evolution of the interface in these processes attracts considerable attention (see, for example, the book [1]) both because of its great practical importance and because of its connections with such fields as dynamical chaos, nonequilibrium physics, and fractal growth (see DLA [2]).

A general scheme for these processes is as follows. There is a linear PDE (frequently of the second order) for the scalar field determining the process. For example, this is the diffusion equation for the Stefan problem and the Laplace or Helmholtz equations for electrodeposition. This scalar field is temperature in the Stefan problem, pressure in flows through porous media, concentration of the nutrient for bacterial growth, electrostatic potential in electrodeposition, probability of the next jump in DLA, etc. Appropriate boundary conditions are imposed both on the moving part of the boundary (the interface) and non-moving part of the boundary (the outer walls). In addition, a law of interface motion is given in terms of the local behavior of the main scalar field. (Typically the local velocity of the interface is proportional to the gradient of the scalar field near the interface). The main question is: ‘what is the evolution of the interface?’

It is remarkable that some of the problems mentioned above are exactly solvable in two dimensions [3]–[5] despite the nonlinearity of these processes. These problems were solved with the help of time–dependent conformal mapping which cannot be extended to 3D except for a few trivial cases. In 2D, it was found that such processes as two–phase flows in porous media, electrodeposition, and slow solidification in a supercooled liquid or from a supersaturated binary solution are governed by an infinite number of the constants of motion which were obtained explicitly in several special cases [3]–[5]. These constants of motion are related to the conserved moments proposed by Richardson [6]. This invariance is quite subtle and disappears when realistic physical perturbations such as surface tension or random noise are added.

In the 3D case very few exact analytical results are known: a constant-velocity paraboloid [7], and a self-similarly growing ellipsoid [8]. The only known way to obtain these solutions is by using one of the eleven coordinate

system in which the 3D Laplace equation is separable [9]. One then considers level surfaces as moving interfaces. It is clear that this method does not work when the shapes are time-dependent. The traditional attitude is that the main obstacle in obtaining nonperturbative exact results in 3D is the lack of (nontrivial) conformal mappings unlike in the 2D case. But is not this statement too strong?

This article is a natural extension of previous work [3] to more general and realistic *multidimensional growth* problems; and not only for the Laplace equation as was done in [3]–[5], but also for *arbitrary elliptic equations* describing for example, inhomogeneous and anisotropic diffusion in solidification, inhomogeneous dielectric functions and screening in electrodeposition, and inhomogeneous viscosity for flows through porous media. It turns out that these nonlinear processes also possess remarkable properties (an infinite number of conservation laws) similar to the ones mentioned in [3], [6] for the 2D Laplacian case; and these properties do not depend on the dimension of the process considered. Thus we show that, contrary to the traditional attitude, *we do not need a conformal mapping for this invariance. Rather this invariance originates from the more general property: the elliptic nature of the equation for the scalar field.*

Let us state now the following D -dimensional problem:

$$L(u) = \operatorname{div}(p(\mathbf{r}) \operatorname{grad}u(\mathbf{r})) + q(\mathbf{r})u(\mathbf{r}) = 0 \quad (1)$$

for $\mathbf{r} \in B \subset R^D$ where the domain B is bounded by the nonmoving exterior boundary, Σ , and by the moving interior boundary $\Gamma(t)$ (t is time) which is the interface separating the domains B and A . An interior domain A contains the origin and is surrounded by the moving interface $\Gamma(t)$ as shown at the Fig.1. Here p and q are given functions of $\mathbf{r} = (x_1, x_2, \dots, x_D)$. The boundary conditions imposed on u are:

$$\partial_n u|_{\Sigma} = G(\Sigma) \quad , \quad (2)$$

$$u(\Gamma(t)) = 0 \quad . \quad (3)$$

The left-hand side (LHS) of (2) means the normal component of $\operatorname{grad}(u)$ evaluated at Σ .

There can also exist point-like sources and sinks in the domain, B , located at \mathbf{r}_k and having strengths, s_k , ($k = 1, 2, \dots, N$), so that near \mathbf{r}_k , $u(\mathbf{r})$ diverges and satisfies

$$u = s_k/|\mathbf{r} - \mathbf{r}_k|^{D-2} + \text{smooth function} \quad (4)$$

if $D > 2$, or

$$u = s_k \log|\mathbf{r} - \mathbf{r}_k| + \text{smooth function}$$

if $D = 2$.

The law of motion of $\Gamma(t)$ is

$$v_n = -p(\mathbf{r}) \partial_n u|_{\Gamma(t)} \quad , \quad (5)$$

where the LHS is the normal component of the velocity of $\Gamma(t)$.

Equations (1) and (5) together with the boundary conditions (2)–(4) complete the mathematical description of the motion of $\Gamma(t)$. If, for example, $D = 3$, $p(\mathbf{r}) = \text{const}$, and $q(\mathbf{r}) = 0$, this describes slow solidification or two-phase flow in porous media.

In this paper, I show that, in spite of the complexity and nonlinearity of the processes described by eqs. (1)–(5), these processes are governed by an infinite number of conservation laws. Namely, if the outer boundary, Σ , is very far from the origin there is an infinite set of numbers C_l ($l = D - 1, D, D + 1, \dots, \infty$) defined as

$$C_l \equiv dM_l/dt \equiv d/dt \left(\int_B \psi_l d^D \mathbf{r} \right) \quad (6)$$

which are conserved during the evolution of the hypersurface¹, $\Gamma(t)$, and equal

$$C_l = 2 \frac{\pi^{D/2}}{\Gamma(D/2)} \sum_{k=1}^N s_k p(\mathbf{r}_k) \psi_l(\mathbf{r}_k) \quad . \quad (7)$$

Here, $\Gamma(n)$ is the Gamma function (**not** the interface $\Gamma(t)$) and ψ_l is the arbitrary solution of $L(u) = 0$ which decays at infinity no slower than r^{-l} and having singularity only at the origin². If $q(\mathbf{r}) = 0$ we have one more conserved quantity: C_0 , which is the rate of the change of the volume of B when L is Laplacian, and which satisfies

¹We do not consider here the passing of the interface through the singularities.

²The functions ψ_l as well as quantities C_l and M_l are labeled in general by more than one number (which is l here). See for example the Eq.(11) below. But for simplicity and without the loss of generality we drop all labels except the l almost everywhere.

$$\begin{aligned}
C_0 &= dM_0/dt = d/dt \left(\int_B p(\mathbf{r}) d^D \mathbf{r} \right) \\
&= 2 \frac{\pi^{D/2}}{\Gamma(D/2)} \sum_{k=1}^N s_k p(\mathbf{r}_k) + \int_{\Sigma} G(\Sigma) p(\Sigma) d\Sigma \quad , \quad (8)
\end{aligned}$$

Here we took $\psi_0 = 1$ which is a solution of $L(u) = 0$ when $q(\mathbf{r}) = 0$.

We think that the knowledge of C_l 's together with the initial M_l 's defined in (6) (the latters are uniquely determined by the initial shape of the interface) could describe the whole moving shape in many of important physical cases. We consider eqs. (7) and (8) as the main result of this work.

Formula (7) follows immediately from the following considerations:

Since

$$\int_{B(t+dt)} \psi_l d^D \mathbf{r} - \int_{B(t)} \psi_l d^D \mathbf{r} = \int_{\Gamma(t)} \psi_l v_n d\Gamma dt$$

we have

$$d/dt \left(\int_{B(t)} \psi_l d^D \mathbf{r} \right) = \int_{\Gamma(t)} \psi_l v_n d\Gamma$$

Further, because of (5), it equals

$$= \int_{\Gamma(t)} (-p \psi_l \partial_n u) d\Gamma$$

and, finally, in view of (3) this expression is

$$\begin{aligned}
&= \int_{\Gamma(t)} p(u \partial_n \psi_l - \psi_l \partial_n u) d\Gamma \\
&= \int_{\Gamma(t)} (\sqrt{p} u \partial_n (\sqrt{p} \psi_l) - \sqrt{p} \psi_l \partial_n (\sqrt{p} u)) d\Gamma \quad . \quad (9)
\end{aligned}$$

Applying Green's theorem to the functions $\sqrt{p} u$ and $\sqrt{p} \psi_l$ we find that the RHS of eq. (9) is given by

$$\begin{aligned}
&\int_B \operatorname{div}(\sqrt{p} u \operatorname{grad}(\sqrt{p} \psi_l) - \sqrt{p} \psi_l \operatorname{grad}(\sqrt{p} u)) d^D \mathbf{r} \\
&+ \int_{\Sigma} p(u \partial_n \psi_l - \psi_l \partial_n u) d\Sigma \quad (10) \\
&+ \sum_{k=1}^N \int_{\gamma_k} p(u \partial_n \psi_l - \psi_l \partial_n u) d\Sigma \quad .
\end{aligned}$$

Here the summation is over the point-like charges, s_k , and γ_k denotes the surface of the infinitesimal hypersphere around the \mathbf{r}_k .

Considering the RHS of (10) one can see that:

- (i) the volume integral over B vanishes because of (1);
- (ii) the surface integral over Σ also vanishes if the outer boundary, Σ , is far removed from the center and if ψ_l decays at infinity stronger than $1/r^{D-2}$. (When $\psi_l = 1$, this integral is not zero but equals $\int_{\Sigma} G(\Sigma) p(\Sigma) d\Sigma$ as it is in the RHS of (8));
- (iii) the contribution of the first integrand to the surface integral over the γ_k is zero, while the integral from the second term equals

$$-2 \frac{\pi^{D/2}}{\Gamma(D/2)} s_k p(\mathbf{r}_k) \psi_l(\mathbf{r}_k)$$

since $p(\mathbf{r})$ and $\psi_l(\mathbf{r})$ are regular near the \mathbf{r}_k and due to Gauss's theorem.

Thus the RHS of (10) equals the RHS of (7) (or (8) when $q(\mathbf{r}) = 0$ and $l = 0$), so we have obtained the infinite set of the conserved quantities, C_l , if the sources and sinks are nonmoving (i.e. when s_k and \mathbf{r}_k are time-independent). Moreover, if $q(\mathbf{r}) = 0$ and no point-like singularities are in the domain B (i.e. if all $s_k = 0$), then *all the dynamics of the interface $\Gamma(t)$ has been reduced to the linear time-dependence of only one "moment", M_0 , which is the volume of the phase B if L is Laplacian. All higher moments, M_l , are constant in time.* Note also that **all moments M_l have a purely geometrical nature**, and thus carry information about the moving shape.

It is worth mentioning the physical interpretation of the derived invariants, M_l , in the special case when the operator L is Laplacian: $p = 1$, $q = 0^3$. For $D = 2$, if the ψ_l are chosen as $\psi_l = z^{-l}$ where $z = x + iy$, these integrals coincide with those previously found in explicit form [3] via the coefficients of the appropriate conformal map. These are analogs of the Richardson moments, whose invariance was found earlier for the interior Hele-Shaw problem [6].

In the 3D Laplacian case, one can choose the ψ_l to be a set of spherical functions:

$$\psi_l^{(m)} = P_{(l-1)}^{(m)}(\theta) e^{im\phi} / r^l \quad . \quad (11)$$

Here, r , θ , and ϕ are the polar coordinates and $P_l^{(m)}(\theta)$ are the associated Legendre polynomials. In this case, the moments, $M_l^{(m)}$, are the

³ In this case it was noted by Howison [8] that the integral of a harmonic function over the infinite domain enclosed the moving bubble is a constant in time.

coefficients of the multipole expansion of the Newtonian potential at an arbitrary point of the empty interior domain A , if the potential is created by the mass uniformly occupying the domain B . Thus the question of whether one can recover the moving shape using only the numbers, M_l , introduced in (7) is now reduced to the classical inverse potential problem [10] for the reconstruction of the shape of a body of constant density from its Newtonian potential. Our case corresponds to the exterior problem (where the potential is given in the empty hollow of the body: in the phase A). The author believes that the connection between pattern formation studies and the inverse potential problem is especially important and deserves close attention. We do not discuss this problem here. Rather we merely note that in 3D (unlike the 2D case) there is no description of a body (with the exception of the sphere) in terms of a finite number of nonzero moments, M_l . (For a detailed description of these difficulties see [11]⁴).

Note that it is also possible in the general elliptic case (when L is not Laplacian) to preserve the interpretation of the conserved quantities M_l as coefficients of the orthogonal expansion by choosing the Green's function $G(\mathbf{r}, \mathbf{r}_1)$ of the operator, L , to be the integrand in (7), since $G(\mathbf{r}, \mathbf{r}_1)$ satisfies the conditions imposed on the ψ_l if $\mathbf{r} \in B$ and $\mathbf{r}_1 \in A$. Using the Green's function expansion and by choosing the orthogonal set of the eigenfunctions ψ_l of $L(\psi_l) = 0$ bounded at infinity and divergent at the origin and the complementary set of eigenfunctions $\tilde{\psi}_l$ of the same equation but with opposite asymptotics, we have

$$U(\mathbf{r}) = \sum_l \tilde{\psi}_l(\mathbf{r}) d/dt \left(\int_B \psi_l(\mathbf{r}_1) d^D \mathbf{r}_1 \right) .$$

We remark that our main result (7) holds also for time-dependent s_k and \mathbf{r}_k (since we never used the time-independence of s_k and \mathbf{r}_k in obtaining (7)). Although the C_l are no longer conserved, the problem is still integrable, since C_l are known functions of time (if the time-dependence of s_k and \mathbf{r}_k is given). Thus the moments, M_l , are easily controlled parameters, namely they are just primitives of the time-dependent RHS of (7). In this way, one might be able to govern the motion of the interface by the proper choice of the s_k and \mathbf{r}_k .

It should also be mentioned that there could be a few exceptions among the C_l 's (only one in the 3D Laplacian case) for which eq. (7) is not valid and thus they may not be conserved. These nonconserved C_l 's correspond to the

⁴I am grateful to M. Brodsky who brought my attention to this book.

ψ_l 's decaying at infinity but not stronger than $r^{(2-D)}$. For the 3D Laplacian case the only nonconserved quantity is C_1 , which corresponds to $\psi_1 = r^{-1}$. Since we do not know the time-dependence of C_1 one could suppose that the description of the interface is now less complete. However, although C_1 is not conserved, we have the conserved quantity C_0 (see eq. (8)). In other words we think that the set $\{M_1, M_2, M_3, \dots\}$ describes the shape with the same degree of completeness as the set $\{M_0, M_2, M_3, \dots\}$. This question concerning nonconserved quantities among the C_l 's does not arise for the interior problem when the scalar field u is given in the phase enclosed by the moving interface $\Gamma(t)$ (in our case in A instead of B).

In conclusion we pose several questions that arise from these studies and which we believe merit attention:

- 1) Does the relation expressed by (7) really mean complete integrability of the multidimensional growth?
- 2) If yes, to which nonlinear evolutionary PDE's do these constants C_l correspond?
- 3) Is there a Hamiltonian structure for these systems?
- 4) Do finite-time singularities (cusps?) exist here, as in the 2D case? And, if yes, how may surface tension regularise them?
- 5) How can one recover the moving shape from the given set of moments, M_l ?

Acknowledgments

It is a pleasant obligation to thank W. Bruno, D. Leschiner, and J. Pearson for a useful discussion, and I.M. Gelfand for his interest in this activity and for his remarks. This work was begun at the Department of Mathematics at Rutgers University and was completed at the Center for Nonlinear Studies in Los Alamos. I wish to thank both of these institutions and J. Lebowitz and G. Doolen for their generous hospitality.

This work was partially supported by AFOSR Grant 91-0010.

References

- [1] "Dynamics of Curved Fronts" ed. by P. Pelce, Academic Press, Inc. (1988)
- [2] T.A. Witten, L.M. Sander, Phys. Rev. Lett. **47**, 1400 (1981)

- [3] M.B. Mineev, *Physica D* **43**, 288 (1990)
- [4] S.D. Howison, *J. Fluid Mech.* **167**, 439 (1986)
- [5] D. Bensimon, P. Pelce, *Phys. Rev. A* **43**, 4477 (1986)
- [6] S. Richardson, *J. Fluid Mech.* **56**, 609 (1972)
- [7] G.P. Ivantsov, *Doklady Acad. Nauk SSSR*, **58**, No. 4, 567 (1947)
- [8] S.D. Howison, *Proc. Royal Society of Edinburgh*, **102A**, 141 (1986)
- [9] P.M. Morse and H. Feshbach, “*Methods of Theoretical Physics*”, McGraw Hill Book Co., Inc. (1953)
- [10] P.S. Novikoff, *Doklady AN SSSR* **18**, 165 (1938) and V. Strakhov, M. Brodsky, *SIAM J. of Appl. Math.* **46**, n.2, 324 (1986)
- [11] G. Herglotz, “Über die analytische Fortsetzung des Potentials ins Innere der anziehenden Massen”, Teubner-Verlag (1914)