

Propagation and Structure of Planar Streamer Fronts

Ute Ebert and Wim van Saarloos

Instituut-Lorentz, Universiteit Leiden, Postbus 9506, 2300 RA Leiden, the Netherlands,

Christiane Caroli

Université Paris VII, GPS Tour 23, 2 Place Jussieu, 75251 Paris Cedex 05, France

Streamers are a mode of dielectric breakdown of a gas in a strong electric field: A sharp nonlinear ionization wave propagates into a non-ionized gas, leaving a nonequilibrium plasma behind. The ionization avalanche in the tip of the wave is due to free electrons being accelerated in the strong field and ionizing the gas by impact. This chain reaction deeper in the wave is suppressed by the generated free charges screening the field. Simulations of streamers show two widely separated spatial scales: the width of the charged layer where the electron density gradients and the ionization rate are very large ($\mathcal{O}(\mu\text{m})$), and the width of the electrically screened, finger-shaped and ionized region ($\mathcal{O}(\text{mm})$). We thus recently have suggested to analyze first the properties of the charge-ionization-layer on the inner scale on which it is almost planar, and then to understand the streamer shape on the outer scale as the motion of an effective interface as is done in other examples of nonequilibrium pattern formation. The first step thus is the analysis of the inner dynamics of planar streamer fronts. For these, we resolve the long-standing question about what determines the front speed, by applying the modern insights of pattern formation to the streamer equations used in the recent simulations. These include field-driven impact ionization, electron drift and diffusion and the Poisson equation for the electric field. First, in appropriately chosen dimensionless units only one parameter remains to characterize the gas, the dimensionless electron diffusion constant D ; for typical gases under normal conditions $D \approx 0.1$ - 0.3 . Then we determine essentially all relevant properties of planar streamer fronts. Technically, we identify the propagation of streamer fronts as an example of *front propagation into unstable states*. In terms of the marginal stability scenario we then find: the front approached asymptotically starting from any sufficiently localized initial condition (the “selected front”), is the steepest uniformly translating front solution, which is physical and stable. Negatively charged fronts are selected by linear marginal stability, which allows us to derive their velocity analytically. Positively charged fronts can only propagate due to electron diffusion against the electric field; as a result their behavior is singular in the limit of $D \rightarrow 0$. For $D \lesssim 1$, these fronts are selected by nonlinear marginal stability and we have to apply numerical methods for predicting the selected front velocity. For larger D , linear marginal stability applies and the velocity can be determined analytically. Numerical integrations of the temporal evolution of planar fronts out of localized initial conditions confirm all our analytical and numerical predictions for the selection. Finally, our general predictions for the selected front velocity and for the degree of ionization of the plasma are in semi-quantitative agreement with recent numerical solutions of three-dimensional streamer propagation. This gives credence to our suggestion, that the front analysis on the inner (μm) scale yields the moving boundary conditions for a moving “streamer interface”, whose pattern formation is governed by the evolution of the fields on the outer (mm) scale.

47.54.+r, 52.80.Mg, 51.50.+v

I. INTRODUCTION

Discharges are nonequilibrium ionization processes occurring in initially non-ionized matter exposed to strong electric fields. Depending on the spatio-temporal characteristics of the electric field and on the ionization and charge transport properties of the medium, discharges can assume many different modes of appearance. In particular in gases under approximately normal conditions one distinguishes phenomenologically between stationary modes like arc, glow or dark discharges and transient phenomena like leaders, the initial stages of sparks, and streamers [1–6], which occur e.g. in silent discharges [7]. The latter nonstationary discharges often form the initial state of a discharge that later on becomes stationary. We

will focus here on an essential element of many transient discharge phenomena, the initial field-driven ionization wave.

The conceptually simplest problem of this kind has become known as the streamer problem in a non-attaching gas. It treats the dynamics of the free electrons and positive ions in a homogeneous gas at rest taking the following mechanisms into account: *(i)* impact ionization, the process in which a free electron accelerated in a strong local field ionizes a neutral molecule, generating a new free electron and a positive ion; *(ii)* drift and diffusion of charged particles, in particular of the electrons whose mobility is much larger than that of the ions; *(iii)* the coupling of the electric field to the charges through the Poisson equation of electrostatics.

Recent numerical simulations [8,9] of a basic model incorporating these physical ingredients for parameter values appropriate for nitrogen under normal conditions reveal that a streamer consists of a sharp nonlinear ionization front which propagates into a non-ionized gas, leaving a weakly ionized nonequilibrium plasma behind. The underlying mechanism is that in the leading edge of the front the electrons are accelerated by the large imposed electric field; this causes the build-up of an electron avalanche due to impact ionization. The generated free charges eventually screen the field and thus suppress further ionization. It is the nonlinear balance between these two nonequilibrium processes, namely the ionization avalanche and the electric screening, which determines the dynamics of the ionization front and the state of the plasma behind it. In confined geometries, streamers usually have a nontrivial fingerlike shape, as is illustrated by the snapshots in Fig. 1 of streamer dynamics taken from the simulations of Vitello *et al.* [9]. As the sharpness of the electron density profiles in Fig. 1 illustrates, the “passive body” of the finger is separated from the external non-ionized gas by a very narrow region — of width of order microns — in which essentially all the action is occurring. This width has to be compared to the size of the filament, which is of order millimeters. It is in this narrow layer that most of the ionization process is taking place. In this same region, there is a nonzero charge density, and consequently, also a very large electric field gradient. These features indicate that there are two different spatial scales in this process, an “inner” scale associated with the thickness of the zone where the ionization takes place, and an “outer” one where the spatial variations are set by the size of the finger and the external experimental geometry. It is precisely for these reasons that accurate simulations are extremely demanding and that they were accomplished only recently by Dhali and Williams [8] and by Vitello *et al.* [9] (See also [10]).

Such a separation of scales is strongly reminiscent of what occurs in combustion fronts [11,12]. A combustion front is a narrow layer of thickness ℓ_{in} to which the combustion is essentially confined, while outside of it, the temperature field varies on a much longer scale ℓ_{out} . Physically, such sharp combustion fronts occur in the limit when the chemical reaction rates involved in the combustion are very fast once a sufficient temperature is reached. It has been shown that, on the basis of an asymptotic expansion to lowest order in the small parameter $\varepsilon = \ell_{in}/\ell_{out}$ using matched asymptotic expansions [13,14], the problem can be analyzed in terms of the propagation of an “effective interface”. More specifically, one first solves the so-called inner problem of a locally *almost planar* reaction zone. This permits to relate the temperature and chemical composition fields on both sides of the front (at distances L such that $\ell_{in} \ll L \ll \ell_{out}$) and to determine the local front velocity as a function of local curvature and fields. On the scale of the remaining outer problem, these relations then play the role of boundary

conditions and of a kinetic equation for the effective moving interface of zero thickness. Besides in combustion, the technique of asymptotic matching to obtain an effective interface description has also been applied to chemical waves [15], thermal plumes [16] and to phase field models of solidification [17,18].

In spite of some important differences between combustion and streamer fronts as discussed in the appendix, a similar approach appears possible for streamers. As discussed also in [19], building on such a reduced description of streamer dynamics appears very desirable, not only because it might make numerical studies much easier, but also because it will allow us to draw upon the knowledge and methods which have been developed in the last decade in the field of interfacial pattern formation and dynamics [20]. The first step towards this goal is to determine the field dependence of the velocity and the ionization and charge profile of a planar front which propagates into the non-ionized region. We thus analyze in this paper the inner problem for a planar streamer front. This allows us to reduce the problem to effectively one dimension. Our analysis clearly identifies the problem of streamer front propagation as an example of front propagation into unstable states. Physically, the instability of the non-ionized gas against charge fluctuations can be traced back to the fact that any small electron density gets amplified by the impact ionization. As is standard for front propagation into unstable states [21–25], we find that the one-dimensional streamer equations exhibit a one-parameter family of uniformly translating front solutions, parametrized by their velocity. As usual [21–25], the question is then to decide which of these front solutions is the dynamically selected one, i.e., is the one reached at long times after a localized ionized region has been created by some initial ionization event. The existing knowledge of front propagation into unstable states [22,23] provides us with an educated guess for the selected velocity, which we confirm with the help of numerical studies. Taken together, our results provide an essentially complete solution of the inner problem of planar streamer fronts.

In itself, the idea to analyze the planar fronts of a streamer model is not new — we refer to [26–29] for earlier work. Apart from the fact, that the authors from the seventies [26–28] investigate different models, which are more inspired by equilibrium concepts (e.g., the ionization behind the front is determined by thermal ionization, where the electron temperature is raised by application of strong electric fields), our work casts new light on this old problem from two different angles:

First of all, it was empirically noted, that the standard approach to analyze uniformly translating fronts, failed to determine a unique propagation velocity, given the field and the gas parameters. Turcotte and Ong [26] clearly state this failure of their theory (this “great defect” of their theory is recalled in Fowler’s reviews [28]) and suggest, that a unique solution might be determined by a dynamical stability analysis. Albright and Tidman

[27] then perform such a stability analysis, but not in a systematic way, and they draw incorrect conclusions. D'yakonov and Kachorovskii [29] also find the indeterminacy of the speed of uniformly translating planar fronts, now for an approximated version of our model, and propose to solve this by using the tip radius of the streamer finger as an extra length scale, which they, however, cannot determine. We, in contrast, trace the indeterminacy of the velocity from the analysis of uniformly translating streamer front solutions to the fact, that this is an example of front propagation into unstable states. Applying the concepts explained above, we solve the selection problem for planar fronts without additional assumptions or approximations. We argue that a particular front solution out of a whole family of dynamically stable solutions is selected, because it is the only one compatible with the initial condition of a localized ionization seed.

Secondly, this result is the first *ingredient* for studying the formation of patterns, in particular of the tip radius — we do not attempt to model *global features of the pattern formation* with our planar front analysis. Our approach thus is very different in spirit to the earlier investigations: As also stressed in [19], in an effective interface description based on a matched asymptotic expansion, the results of weakly curved, almost planar fronts are essentially used *locally* everywhere in the interface region: They enter the analysis on the outer scale as boundary conditions at the moving interface. It is on this outer scale that pattern formation problems like the size, velocity and shape of the streamer should be analyzed. Once our results on planar fronts will be extended to weakly curved fronts, all the necessary ingredients to tackle these questions appear to be available.

The main results of our present analysis of the streamer equations used in the simulations [8,9] can be summarized as follows:

(a) Dimensional analysis shows that in dimensionless units, a single parameter remains to characterize the gas, the dimensionless electron diffusion coefficient D characteristic of the gas [see Eq. (2.10)]. For gases under normal conditions, D is small, of order 0.1-0.3.

(b) The length scale set by the electron impact ionization coefficient [the coefficient α_0^{-1} in Eq. (2.6)] is on the order of microns for nitrogen. For $D \lesssim 1$ the thickness ℓ_{in} of the charged layer is on the order of this same ionization length for *negatively charged streamer fronts (NSF)* [30]. Given that typical streamer diameters found in the simulations are of the order of 1mm, $\varepsilon = \ell_{in}/\ell_{out}$ is at most of order 10^{-2} ; this justifies an effective interface description of streamer dynamics.

(c) We find that electron diffusion acts as a singular perturbation for *positively charged streamer fronts (PSF)*: without diffusion, such fronts can not propagate, but with any nonzero D , they do. As a result, the behavior is singular in the limit $D \rightarrow 0$: for $D = \mathcal{O}(1)$, the thickness ℓ_{in} is again of order of the ionization length, but for $D \rightarrow 0$ the electron density and its gradients diverge due to the appearance of another smaller length

scale (of order D/α_0).

(d) The electron density generated by the propagating front is again basically set by dimensional analysis for *NSF*. We calculate for $D \lesssim 1.5$ the dependence of the dimensionless electron density σ^- behind the front on the electric field E^+ far ahead of our planar front. Our results compare favorably with those extracted from the simulations [9], according to the prescriptions of the theory of matched asymptotic expansions [13,14]. Namely, E^+ is *not* the field value at the electrode position, but the value obtained by extrapolating the slowly varying outer field to the front position. We also calculate the full D and E^+ dependence of the electron density σ^- behind the front of *PSF* for $D \lesssim 1.5$.

(e) The dynamically relevant (“selected”) front velocity v_f is a unique function of E^+ and D . The analysis confirms the strong asymmetry between *NSF* and *PSF* also found in the simulations [8,9] for fronts propagating into an essentially non-ionized region. The asymmetry is the stronger the smaller D and disappears for $D \gg 1$.

(f) For *NSF*, v_f is given by the so-called linear marginal stability velocity v^* [22] — see Eq. (5.3) below. For parameter values used in the simulations, we find that v_f is typically 30 to 40 % higher than the electron drift velocity just in front of the streamer head, which agrees semi-quantitatively with the findings of Vitello *et al.* [9].

(g) We find that *PSF* propagate for any nonzero value of the dimensionless electron diffusion coefficient D . Due to the singular behavior as $D \rightarrow 0$, we find that fronts propagate with a unique velocity v^\dagger predicted by the so-called nonlinear marginal stability mechanism [23] for small D . For the Townsend expression used in the simulations [8,9], this happens below a well-defined field-dependent value of D of order unity [see Fig. 3]. Above this threshold value, *PSF* propagate with the linear marginal stability value v^* .

In this paper, our main focus will be on those results that are of greatest interest from the point of view of understanding the generation of low temperature plasmas by the streamer mechanism. We note, however, that the equations for planar streamer fronts [Eqs. (3.11) and (3.12) below] appear to be of interest in their own right. As will be discussed briefly in Section V, our streamers have several features in common with the celebrated nonlinear diffusion equation studied in mathematics [31,32] since the early work of Kolmogorov *et al.* [33] and Fisher [34]; at the same time, however, they are sufficiently more complicated that they appear to present new challenges from a mathematical point of view.

This paper is organized as follows. In Section II we introduce the basic equations for streamer formation, and perform a dimensional analysis for the inner problem of streamer fronts. In Section III, we discuss the stability of the basic homogeneous states of interest, the homogeneous non-ionized state and the homogeneous weakly ionized state. We also discuss the physical mechanism of streamer formation and the proper initial and bound-

ary conditions to study these in the case of planar fronts, which allow us to simplify the equations describing planar front dynamics. In Section IV we demonstrate that there exists a one-parameter family of uniformly translating fronts characterized by a continuous range of front velocities v . We also briefly show how in the case $D = 0$, the equations for uniformly translating fronts can be solved analytically. These solutions, which turn out to be useful as a small- D approximation for *NSF*, yield an explicit expression for the electron density σ^- behind the *NSF* in terms of the field E^+ just ahead of it. This is followed by an analysis of the general case $D \neq 0$; then the equations can not be solved analytically, but we demonstrate that there still is a one-parameter family of uniformly translating front solutions. For *PSF*, we show that the limit $D \rightarrow 0$ is singular; we discuss this limit in detail and show that it accounts for the strong asymmetry between *PSF* and *NSF* for realistic values of D . In Section V we then summarize some of the main results [21–25] concerning the so-called selection problem, the question which particular front solution from the family is reached asymptotically for large times for a large class of initial conditions. Application of these concepts allows us to predict the shape and velocity of the dynamically relevant front solution (the selected front) and the value of the electron density generated behind it. This yields the various selection results for *NSF* and for *PSF*, summarized in points (c)-(g) above, and leads us to predict that the behavior of *PSF* in the limit $D \rightarrow 0$ is singular. In Section VI we present numerical simulations of the full partial differential equations for planar streamer dynamics; starting from various initial conditions, we illustrate, that in all cases we have studied the long time dynamics of the system is characterized by a *NSF* and a *PSF* whose behavior is in full agreement with our predictions. In the concluding section we finally reflect on our results and on the future steps to be taken to arrive at an effective interface description of streamer dynamics. In an appendix we discuss differences and similarities between combustion and streamer fronts.

II. MODELING AND DIMENSIONAL ANALYSIS

A. The Minimal Streamer Model

For simulating the dynamical development of streamers out of a macroscopic initial ionization seed in a so-called non-attaching gas like N_2 under normal conditions, Dhali and Williams [8] and Vitello *et al.* [9] use the following set of deterministic continuum equations for the electron density n_e , the ion density n_+ and the electric field \mathcal{E} : balance equations for electrons and ions,

$$\partial_t n_e + \nabla_{\mathbf{R}} \cdot \mathbf{j}_e = source , \quad (2.1)$$

$$\partial_t n_+ + \nabla_{\mathbf{R}} \cdot \mathbf{j}_+ = source , \quad (2.2)$$

where the fact that the two source terms are the same is due to charge conservation in an ionization event; the Poisson equation,

$$\nabla_{\mathbf{R}} \cdot \mathcal{E} = \frac{e}{\epsilon_0} (n_+ - n_e) , \quad (2.3)$$

and the approximate phenomenological expressions

$$\mathbf{j}_e = -n_e \mu_e \mathcal{E} - D_e \nabla_{\mathbf{R}} n_e , \quad (2.4)$$

$$\mathbf{j}_+ = 0 , \quad (2.5)$$

$$source = |n_e \mu_e \mathcal{E}| \alpha_0 e^{-E_0/|\mathcal{E}|} . \quad (2.6)$$

Apart from the fact that we will allow for a slight generalization of Eq. (2.6), these are the equations that we will investigate analytically below.

In these equations, \mathbf{j}_e and \mathbf{j}_+ are the particle current densities of electrons and positive ions, and e is the absolute value of the electron charge. The (dimensional) spatial coordinates are denoted by \mathbf{R} , and $\nabla_{\mathbf{R}}$ is the gradient with respect to these coordinates. The use of only Poisson's law of electrostatics, Eq. (2.3), means that all magnetic fields as well as terms in the Maxwell equations associated with time-dependences of the fields, are neglected [35].

The electron particle current density \mathbf{j}_e is approximated in (2.4) as the sum of a drift and a diffusion term. Note that this diffusion approximation implies that the electron mean free path must be small with respect to the scale of variation ℓ_{in} of the electric field. This condition is just about satisfied for the parameter values taken for N_2 in the simulations, except possibly at the highest field values (see also the discussion in the concluding Section VII). The electron drift velocity is taken to be linear in the field \mathcal{E} , with μ_e the (positive) electron mobility. The electron diffusion coefficient D_e and the mobility μ_e are treated here as independent coefficients, since they effectively depend on the field strength [3] (only in the low-field limit are they related by the Einstein relation). More generally, the diffusion coefficient should be replaced by a diffusion tensor, which is diagonal in a reference frame with one axis along the electric field. Its longitudinal component, the only relevant one for planar fronts perpendicular to E , is somewhat smaller than the transverse one. Since we will see, that N_2 reaches a typical degree of ionization of only 10^{-5} , density fluctuations of the non-ionized gas can be neglected and the mean free path of the electrons and therefore μ_e and D_e can be taken as independent of the degree of ionization.

The ionic current is neglected according to Eq. (2.5), since the mobility of ions is at least two orders of magnitude smaller than that of the electrons [8]. In particular, for the analysis of the inner scale, that we will perform in the present paper, \mathbf{j}_+ is negligible.

The *source* (2.6) finally accounts for the creation of free charges by impact ionization. If the product of electric field \mathcal{E} and electronic mean free path ℓ_{mfp} is large enough, free electrons can gain sufficient kinetic energy to

ionize neutral molecules. Accordingly there is a threshold field $|\mathcal{E}| = E_0 \propto \ell_{mfp}^{-1}$. For $|\mathcal{E}| \gtrsim E_0$ the probability that a scattering event carries at least the ionization energy is large. The effective ionization cross-section $\sigma_{cs}(|\mathcal{E}|)$ then essentially saturates, while for $|\mathcal{E}| \ll E_0$ the ionization rate per scattering event is largely suppressed. The *source*-term is given by the ionization rate, which can be calculated as the product of the drift current of free electrons $|n_e \mu_e \mathcal{E}|$ times the target particle density n_n of the neutral gas times the effective ionization cross-section $\sigma_{cs}(|\mathcal{E}|)$. Commonly, a phenomenological ionization coefficient $\alpha(|\mathcal{E}|) = n_n \sigma_{cs}(|\mathcal{E}|)$ is used, (which clearly has dimension of inverse length,) whose field threshold behavior in the Townsend approximation $\alpha(|\mathcal{E}|) = \alpha_0 \exp(-E_0/|\mathcal{E}|)$ [3] is expressed by Eq. (2.6). As discussed by Raizer [3], in the approximation that every collision is ionizing, if the electron carries an energy larger than the ionization energy I , we have

$$\alpha_0 \approx \ell_{mfp}^{-1}, \quad \text{and} \quad E_0 \approx I/(e \ell_{mfp}). \quad (2.7)$$

Since in much of our analysis the specific form of $\alpha(|\mathcal{E}|)$ is not needed, we will use a slightly more general formulation in Eq. (2.11) below.

In the *source*-term, ionization due to the photons also created in recombination or scattering events, is neglected. This is motivated by the ionization cross-sections due to photons being much smaller than those due to electrons. Note that, if photo-ionization is taken into account, the dynamical equations become nonlocal.

No sink term needs to be included for the analysis of the inner problem, since the recombination length at a degree of ionization of order 10^{-5} that we will derive below is very large as compared with the front width ℓ_{in} . (For this reason, the inner problem is the same for streamers and leaders [3]: the difference between these discharge modes, which consists in the fact that recombination is non-negligible in the plasma body of leaders, would come into play only when solving, at a later stage, the outer problem.) The fact that the degree of ionization remains small is also the reason that saturation effects are neglected in (2.6).

In contrast to the situation in N_2 , that is described by our model equations, in attaching gases like O_2 , a third charged species plays a role, namely negative ions formed by a neutral molecule catching a free electron. For a description of the physics of such attaching gases and simulations thereof, see, e.g., [36].

The equations above are deterministic. Thermal fluctuations in fact can be neglected, since even an unphysically small ionization energy of 3 eV leads to a Boltzmann factor of 10^{-52} at room temperature. Also other stochastic effects are not accounted for in the simulations we compare to. We further discuss possible stochastic effects in the experiments in the conclusion.

Finally, the dynamical system (2.1)-(2.6) must be complemented by:

(i) boundary conditions: as will be discussed in detail in Section III, for the problem of front propagation, these

are specified by the value E^+ of the electric field far ahead of the front, where the total charge density vanishes.

(ii) initial conditions: we ignore the details of the plasma nucleation event (e.g. triggering by radiation from an external source), and assume that at $t = 0$ a small well-localized ionization seed is present. The precise meaning, for our problem, of “well-localized” will be made clear in Section V.

B. Dimensional Analysis

In order to identify the physical scales and intrinsic parameters of our problem, we reduce Eqs. (2.1)-(2.6) to a dimensionless form. The most natural scale of length and electric field are the ionization length $\ell_{ion} = \alpha_0^{-1}$ and the threshold field E_0 of the ionization rate (2.6). The velocity scale is then the electron drift velocity at this field strength, $v_0 = \mu_e E_0$, leading to a time unit $t_0 = (\alpha_0 \mu_e E_0)^{-1}$, and a charge unit $q_0 = \varepsilon_0 \alpha_0 E_0$.

For concreteness, we list here the values of these quantities for N_2 at normal pressure, used in the simulations [8,9]

$$\begin{aligned} \alpha_0^{-1} &\approx 2.3 \mu\text{m}, & v_0 &\approx 7.56 \cdot 10^7 \text{ cm/s}, \\ t_0 &\approx 3 \cdot 10^{-12} \text{ s}, & q_0 &\approx 4.7 \cdot 10^{14} \text{ e/cm}^3 \\ E_0 &\approx 200 \text{ kV/cm} & \mu_e &\approx 380 \text{ cm}^2/\text{Vs}. \end{aligned} \quad (2.8)$$

We now introduce dimensionless quantities by defining

$$\begin{aligned} \mathbf{r} &= \mathbf{R} \alpha_0, & \tau &= t/t_0, \\ q &= (n_+ - n_e) e/q_0, & \sigma &= n_e e/q_0, \\ \mathbf{j} &= -\mathbf{j}_e e/(q_0 v_0), & \mathbf{E} &= \mathcal{E}/E_0. \end{aligned} \quad (2.9)$$

Note that with our definition, \mathbf{j} now plays the role of a dimensionless *charge* current. If we furthermore introduce the dimensionless diffusion coefficient D as

$$D = D_e \alpha_0 / \mu_e E_0, \quad (2.10)$$

we obtain what we call the streamer equations

$$\partial_\tau \sigma - \nabla \cdot \mathbf{j} = \sigma f(|\mathbf{E}|), \quad (2.11)$$

$$\partial_\tau q + \nabla \cdot \mathbf{j} = 0, \quad (2.12)$$

$$q = \nabla \cdot \mathbf{E}, \quad (2.13)$$

$$\mathbf{j} = \sigma \mathbf{E} + D \nabla \sigma, \quad (2.14)$$

where ∇ denotes the gradient with respect to the dimensionless coordinate \mathbf{r} , and where the “ionization function”

$$f(|\mathbf{E}|) = |\mathbf{E}| \alpha(|\mathbf{E}|) / \alpha_0 \quad (2.15)$$

is assumed to vanish at zero field. Townsend’s expression (2.6) yields:

$$f_T(|\mathbf{E}|) = |\mathbf{E}| \exp(-1/|\mathbf{E}|). \quad (2.16)$$

In general, we will treat an ionization function with the properties [37]

$$f(0) = 0 = f'(0), \quad \text{and} \quad f'(|\mathbf{E}|) \geq 0 \quad \text{for all} \quad |\mathbf{E}|. \quad (2.17)$$

The dimensionless equations (2.11)-(2.14) now depend on only one internal parameter, the dimensionless diffusion coefficient D . For the values used in [8,9] for N_2 under normal conditions, $D \approx 0.1$, while according to the data given by Raizer [3], for Ne and Ar, $D \approx 0.3$. We believe that typical values are generally in the range 0.1-0.3, since in the approximation (2.7), $\alpha_0/E_0 \approx I/e$ and since the ratio D_e/μ_e appears to be commonly of the order of volts for large fields, while I is typically of the order of several electron volts.

We are now able, solely on the basis of the dimensional analysis above, to make a first semi-quantitative prediction about streamers. We will in practice be interested in external fields $E^+ = \mathcal{O}(1)$ (for $E^+ \ll 1$ and α_0^{-1} on the order of micrometers, the electron avalanche process becomes much too ineffective for streamer fronts to develop at reasonably small distances; also our scale separation approach discussed in the introduction might break down). We can therefore expect that, for D values $\lesssim 1$, as is the case for N_2 , front widths will be of order α_0^{-1} , and that in addition the reduced electron density σ^- far behind the front on the inner scale will be of order unity as well. This leads one to expect electron densities in the streamer body on the order of 10^{14} cm^{-3} , in agreement with numerical findings.

III. HOMOGENEOUS SOLUTIONS AND THE CONCEPT OF FRONTS

A. Homogeneous States and their Stability

The first task, when studying in general the propagation of a front, is to identify the nature and stability of the states which the front connects. We expect the invading state, here the ionized one created by the front, to be stable [38], while the invaded state can in general either be metastable or unstable. Physically, we of course expect the non-ionized state to be unstable in a non-vanishing field in the present model. (In an attaching gas forming also negative ions, it is conceivable that the non-ionized state is metastable for not too strong fields.)

Equations (2.11)-(2.14) immediately yield, that stationary homogeneous states simply are solutions of

$$\sigma f(|\mathbf{E}|) = 0. \quad (3.1)$$

So, these stationary states decompose into two families:

(i) Non-ionized states, with $\sigma = 0$, \mathbf{E} arbitrary: Since the density of free electrons vanishes, no ionization can occur, whatever the value of \mathbf{E} is. If also the density of ions vanishes, $\nabla \cdot \mathbf{E} = 0$. Since these states correspond to the physical situation far ahead of the front, we label them (+). Moreover, since we will need in particular the

case in which the field ahead of the front is constant, we take $\mathbf{E}^+ = \text{const.}$

(ii) Completely screened states, labeled (-), with $\mathbf{E} = 0$, σ^- arbitrary [39]: Whatever the electron density, for $\mathbf{E} = 0$ impact ionization does not occur and thermal energy is much too small to permit ionization.

Since the steady states we consider as well as the equations of motion are translation invariant in space and time, the eigenstates of the linear perturbations are Fourier modes of the form

$$\begin{pmatrix} \delta\sigma(\mathbf{r}, t) \\ \delta\mathbf{E}(\mathbf{r}, t) \end{pmatrix} = \begin{pmatrix} \sigma_1 \\ \mathbf{E}_1 \end{pmatrix} \exp(i\mathbf{k} \cdot \mathbf{r} + \omega\tau). \quad (3.2)$$

We first investigate the linear stability of the non-ionized state $\sigma^+ = 0$. Upon linearizing the equations about the zeroth order values ($\sigma^+ = 0, \mathbf{E}^+$), we find two branches of modes:

(a) The first, trivial branch is a zero mode ($\omega = 0$), with $\sigma_1 = 0$, expressing that the electron density remains zero. This zero mode accounts for the degeneracy of the non-ionized states, i.e., for the fact that there exists a (+) stationary state for each value of \mathbf{E}^+ . (For $\mathbf{E}^+ \neq \text{const.}$, these zero modes express the degeneracy of all steady states with $q^+ = \nabla \cdot \mathbf{E}^+$ for any ion density q^+ as long as the electron density σ^+ vanishes.)

(b) The second branch of perturbations is associated with fluctuations carrying a finite electron charge; its dispersion relation is

$$\omega^+ = i\mathbf{k} \cdot \mathbf{E}^+ + f(|\mathbf{E}^+|) - Dk^2, \quad (3.3)$$

with $i\omega^+ \mathbf{k} \cdot \mathbf{E}_1 = (f(|\mathbf{E}^+|) - \omega^+) \sigma_1$. The first term on the r.h.s. of (3.3) simply expresses the fact that the electrons drift, to first order, in the electric field \mathbf{E}^+ with velocity ($-\mathbf{E}^+$). The real part $\Re\omega^+$, the sign of which determines whether fluctuations decay or are amplified, contains a destabilizing term, expressing that any small electron density fluctuation is amplified at rate f , and a stabilizing term, due to the diffusive spreading of electron charges. For $k^2 < f(|\mathbf{E}^+|)/D$, $\Re\omega^+ > 0$: non-ionized states are unstable against long-wavelength perturbations.

We note, that the single Fourier eigenmodes (3.2) violate individually the physical constraint that σ be positive everywhere. But Eq. (3.3) also determines the time evolution of physically allowed fluctuations (wavepackets) that are superpositions of these eigenmodes. For example, one easily deduces from it Lozanski's expression [40] for the time-evolution of a Gaussian-shaped small electron density with arbitrary constants $c_1, c_2 > 0$,

$$\delta\sigma(\mathbf{r}, \tau) = c_1 e^{f(|\mathbf{E}^+|)\tau} (c_2 + 4D\tau)^{-3/2} e^{-\mathbf{r} + \mathbf{E}^+\tau)^2 / (c_2 + 4D\tau)}, \quad (3.4)$$

as long as linearization around the non-ionized state holds. As expected, the center of the spreading packet drifts with velocity $-\mathbf{E}^+$, while the total number of electrons it contains is amplified at rate f and the wave-packet stays Gaussian, with time-dependent width

$c_2 + 4D\tau$. Such ionization modes derived by linearizing around the non-ionized state are known as electron or ionization avalanches in the gas discharge literature.

We now perform the same linear stability analysis for the completely screened states ($\sigma^- = \text{const.}, \mathbf{E}^- = 0$). The fact that $f'(0) = 0$ from Eq. (2.17) assures that the linear perturbations are not affected by ionization; the dynamics thus evolves with conserved particle densities.

Again, due to the existence of a continuous family of screened stationary states, parametrized by σ^- , the spectrum contains a branch of $\omega = 0$ modes. For the nontrivial branch, the dispersion relation is given by

$$\omega^- = -\sigma^- - Dk^2, \quad (3.5)$$

while the eigendirection of such a perturbation is given by

$$\sigma_1 + i\mathbf{k} \cdot \mathbf{E}_1 = \sigma_1 + q_1 = 0. \quad (3.6)$$

Since $(\sigma_1 + q_1)$ is the dimensionless ion density of the linear mode, Eq. (3.6) simply expresses the fact that ions are completely immobile in our model.

Equation (3.5) expresses the fact that the completely screened ($-$) states are stable, the decay of perturbations being due to the added stabilizing effects of overdamped plasmons ($-\sigma^-$) and electron diffusion. The $\mathbf{k} \rightarrow 0$ limit of the plasmon mode leads to dielectric screening [41].

B. The Mechanism of Front Creation

Let us now investigate the dynamical evolution of an initial state in which the electron and ion densities vanish everywhere except in a small localized region. An example of such localized initial conditions is an initially Gaussian electron density, as in the simulations [8,9] — under what circumstances initial conditions are sufficiently localized will become clear later. As long as the electron and ion densities are small enough, we can neglect in linear approximation the changes in the field as we did above when linearizing about the non-ionized state. As a result, both densities will grow due to impact ionization. If this were the only mechanism, the space charge would remain unchanged and the ionization would continue indefinitely. However, the electrons are mobile, and at the same time they start to drift in the direction opposite to the electric field \mathbf{E} . If we neglect for the moment the diffusion, this drift has two effects: First of all, the electrons start to drift in the direction of the anode. Impact ionization then starts in previously non-ionized regions as well, so the ionized region expands towards the anode. Secondly, as the electrons drift while the ions stay put (on the fast time scale), a charge separation occurs which tends to suppress the field strength in the ionized region. When the size of the initial perturbation and/or the time during which the avalanche has built up are large enough, the screening of the field becomes almost complete in the

ionized region so that ionization stops there. The behavior in this region can be described by linearizing around the screened state as done above. After an electrically screened body of the ionized region has developed, the initial ionization avalanche is said to have developed into a streamer. Thus streamer fronts are strongly nonlinear and determined by two competing mechanisms, which dynamically balance each other: the ionization process which is strongest at the leading edge and the screening of the field due to the free charges which increases towards the rear end of the front. This balance also explains our finding that the ionization length and the screening length in the plasma behind the *NSF* are of the same order of magnitude for field values that are not too small. Technically speaking, the challenge in constructing the full front is to connect the two regimes linearized about the homogeneous states in an appropriate way through the nonlinear regime of the front.

In the above discussion, we have neglected electron diffusion. In this case the *NSF* propagates towards the anode with a velocity that is at least the drift velocity of the electrons in the local electric field. The *PSF*, in contrast, is moving in the direction *opposite* to the drift of the only mobile species, the electrons. Its space charge is formed by the ions staying put, while the electrons are drawn into the ionized body. Propagation of a *PSF* is therefore only possible if the electron diffusion current overcompensates the drift current. This in turn implies that if the diffusion coefficient D is small, electron density gradients must be extremely steep. From this discussion it already becomes evident — and we will derive this below — that for an *NSF*, diffusion is a small correction for $D \ll 1$, since drift and diffusion currents are acting in parallel directions. In *PSF*, however, diffusion has to overcome the drift, and as a result in this case the limit of vanishing diffusion is very singular. We will see in Section IV that this manifests itself through the emergence of a new inner length scale $D/\alpha_0 = D_e/(\mu_e E_0)$, the diffusion length associated with the electron drift velocity.

Of course, a charged front only screens the normal component of the electric field. This is why electric screening is efficient in the head of the streamer, while the field penetrates in the body of a single streamer in the simulations [8,9]. Our planar front analysis thus serves as a first approximation for the mechanisms in the moving tip of the streamer finger.

C. The one-dimensional Streamer Equations

Let us now restrict our analysis to the case of plane fronts perpendicular to a constant electric field. Of course, in practice planar streamer fronts will be unstable to deformations along the front (very much like in the Mullins-Sekerka instability in crystal growth [20]), but as explained in the introduction, the planar front analysis is a first step towards understanding the dynamics on the

inner length scale α_0^{-1} and time scale t_0 . As such, it is the first basic ingredient for deriving an effective interface model on scales $\gg \alpha_0^{-1}$.

We choose the x axis as parallel to the field and perpendicular to the planar front so that $\mathbf{E} = E\hat{\mathbf{x}}$ and $\nabla = \hat{\mathbf{x}} \partial_x$. From the point of view of matched asymptotic expansions, the electric field in the non-ionized region before the front will vary adiabatically slowly on the “inner” time scale τ of the front, the timescale on which the front propagates over a distance comparable to its width, because the length scales of the outer problem determining the changes of E are assumed to be much larger than the inner scale α_0^{-1} . For our study of the inner problem, we thus take the asymptotic field value \mathbf{E}^+ in the unionized region constant in time. Furthermore, we will use the convention that the unionized initial state into which the front propagates is at the right towards large positive values of x , so that there

$$\sigma \rightarrow \sigma^+ = 0, \quad q \rightarrow q^+ = 0, \quad E \rightarrow E^+, \quad \partial_\tau E^+ = 0, \\ \text{for } x \rightarrow +\infty, \quad (3.7)$$

which motivates now the use of the superscript $+$. We emphasize again that “ $x \rightarrow +\infty$ ” should be interpreted on the length scale α_0^{-1} of the inner problem in the sense of matched asymptotic expansions [13,14]. Far behind the front, i.e., for $x \rightarrow -\infty$, the discussion of Section II leads us to expect a homogeneous stable state

$$\sigma \rightarrow \sigma^- \neq 0, \quad q \rightarrow q^- = 0, \quad E \rightarrow 0, \\ \text{for } x \rightarrow -\infty. \quad (3.8)$$

Which value σ^- will be dynamically selected and what the corresponding front velocity and profile are, for a given fixed value of the electric field E^+ before the front, is the selection problem, we aim to solve.

The boundary condition (3.7) allows an important simplification of the equations in one dimension: If we insert (2.13) into (2.12), we obtain

$$\partial_\tau E + j = h(\tau), \quad (3.9)$$

where $h(\tau)$ is an arbitrary function of time which is constant in space. In view of the boundary condition (3.7), $h(\tau)$ vanishes at $x \rightarrow \infty$ and thus everywhere. For planar fronts, the model Eqs. (2.11)-(2.14) then reduce to

$$\partial_\tau \sigma = \partial_x (\sigma E) + D \partial_x^2 \sigma + \sigma f(|E|), \quad (3.10)$$

$$\partial_\tau E = -\sigma E - D \partial_x \sigma, \quad (3.11)$$

with space charge and electric current given by

$$q = \partial_x E \quad \text{and} \quad j = \sigma E + D \partial_x \sigma. \quad (3.12)$$

We will refer to this set of equations as to *the one-dimensional streamer equations*. They are the basic equations of this paper, on which the rest of our analysis will be based.

Eq. (3.11) implies that the field decays behind the front, if no strong density gradients act against it. As we shall see later when we will discuss our simulation results in Section VI, such strong density gradients often occur during the transient regime before a *PSF* emerges. Once, however, a front has approached an approximately uniformly translating state, the electron density σ^- behind the front is almost homogeneous and the field behind the front then decays to zero on a time scale $1/\sigma^-$ according to (3.11). Note, that the local decay of the field for any nonzero electron density is due to electro-dynamics of conserved quantities that continues also after the impact ionization has been suppressed.

We finally note that in the limit where the diffusion is small ($D \ll 1$), it is easy to identify the crossover time from the linear avalanche regime to that of streamer propagation in the case that the initial electron density is small and nonzero only in a very narrow localized region. As explained in the beginning of this section, in the avalanche regime we can neglect the changes in the background field E^+ due to the build-up of the charges. The evolution of the electron density is then described by the linearized version of (3.10), a linear equation with drift, diffusion and growth. Hence, if the initial electron density is, e.g., Gaussian, the electron density will according to (3.4) remain a Gaussian profile, whose maximum drifts with a velocity $|\mathbf{E}|$ in the direction opposite to the field and whose amplitude grows exponentially as $\exp(f(E^+)\tau)$. In other words, if the total initial electron charge is $N_e(0) = \int dx \sigma(x,0)$, then the total number of electrons in this avalanche regime grows as $N_e(\tau) = N_e(0) \exp(f(E^+)\tau)$. Likewise, the total ion charge grows exponentially, but if both the diffusion constant and the width and amplitude of the initial perturbation are small, the electron drift will separate the negative electron charge and the positive ion charge almost completely. The crossover to the nonlinear streamer regime will therefore occur when the total charge in the positively and negatively charged regions is big enough that screening of the field becomes appreciable, i.e., at a time τ_c when

$$N_e(\tau_c) \approx |E^+|, \quad \Rightarrow \quad \tau_c \approx \frac{1}{f(E^+)} \ln[|E^+|/N_e(0)]. \quad (3.13)$$

IV. UNIFORMLY TRANSLATING FRONT SOLUTIONS

Above we already have introduced the idea, that fronts asymptotically approach some shape, which is independent of the initial conditions. This is based on our experience [21–25] with other examples of front propagation into unstable states that the front will acquire some asymptotic shape and velocity in the long time limit, which will be the same (“universal”) for a large class

of “sufficiently localized” initial conditions that comprise most physically relevant initial states. This property is often referred to as the front selection problem. Our subsequent analysis will therefore follow the usual strategy in examples of this type: We will first show in this section that there generally is a one-parameter family of front solutions. In Section V we then summarize our present understanding of the front selection problem, and on the basis of this predict the properties of the selected streamer front. The numerical simulations that confirm our predictions are presented in Section VI.

Uniformly translating fronts with velocity v are stationary in a coordinate system moving with velocity v . If we denote this comoving coordinate by $\xi = x - v\tau$, the partial differential equations (*pde*'s) (3.11) and (3.12) in this coordinate system become

$$\begin{aligned}\partial_\tau \sigma|_\xi &= v \partial_\xi \sigma + \partial_\xi (\sigma E) + D \partial_\xi^2 \sigma + \sigma f(|E|) , \\ \partial_\tau E|_\xi &= v \partial_\xi E - \sigma E - D \partial_\xi \sigma .\end{aligned}\tag{4.1}$$

A front translating uniformly with velocity v in the *fixed* frame x is stationary in this comoving frame, $\partial_\tau \sigma|_\xi = 0 = \partial_\tau E|_\xi$. As a result, the corresponding front profiles are solutions of the ordinary differential equations (*ode*'s). (We continue to use partial differential signs ∂_ξ even though the uniformly translating solutions are functions of the variable ξ only.)

$$\begin{aligned}D \partial_\xi^2 \sigma + (v + E) \partial_\xi \sigma + \sigma \partial_\xi E + \sigma f(|E|) &= 0 , \\ D \partial_\xi \sigma - v \partial_\xi E + \sigma E &= 0 ,\end{aligned}\tag{4.2}$$

These equations are analyzed below. Both for $D = 0$ and for $D \neq 0$, they admit solutions for a range of values of the velocity, so we are indeed faced with the question of front selection.

It is important to realize that not all the exact uniformly translating front solutions of these *ode*'s correspond to physically relevant solutions. In particular, any physical electron density σ needs to be non-negative ($\sigma \geq 0$), but as we shall see the set (4.2) admits *PSF* solutions where σ goes negative. We expect these solutions to be unstable (in accord with the “nonlinear marginal stability” scenario [23]), and also not to be approachable from an initial condition with $\sigma \geq 0$. Hence they are neither dynamically nor physically relevant. Furthermore, note that in our model the ion density q_i ($= \rho + \sigma$) can only increase due to impact ionization [Eqs. (2.11) and (2.12) imply $\partial_\tau q_i = \sigma f(E) \geq 0$]. With our convention that the non-ionized state is on the right, this implies that uniformly receding front solutions with $v \leq 0$ are unphysical. We will therefore call a uniformly translating front solution physical if

$$v > 0 \quad \text{and} \quad \sigma(\xi) \geq 0 \quad \text{for all } \xi .\tag{4.3}$$

A. $D = 0$ Front Solutions

In contrast to the case $D \neq 0$, where we can derive properties of uniformly translating fronts only either qualitatively by discussing flows in phase space or quantitatively by numerical integration, Eqs. (4.2) for $D = 0$ can be integrated explicitly. Doing so, we derive a simple explicit expression for the electron density σ^- behind the front in terms of the field E^+ before the front; this analysis generalizes an earlier result of D'yakonov and Kachorovskii [29], and explicitly illustrates the existence of a family of uniformly translating solutions. For *NSF*, these results extend smoothly to the case $D \neq 0$: The electron density $\sigma^-(E^+)$ derived for $D = 0$ will turn out to be a good approximation for $D \lesssim 1$, and the small overshoot of σ at the rear end of the front visible in the three-dimensional simulations in Fig. 1(c), is also recovered for $D = 0$. For *PSF*, on the other hand, we will see that D acts as a singular perturbation, so that the class of $D = 0$ *PSF*-solutions that we derive here is not relevant for the *PSF* selection problem for $D \lesssim 1$.

The *ode*'s describing uniformly translating fronts for vanishing diffusion are found by putting $D = 0$ in (4.2). These equations then become

$$\partial_\xi \left[(v + E) \sigma \right] = - \sigma f(|E|) ,\tag{4.4}$$

$$v \partial_\xi \ln |E| = \sigma .\tag{4.5}$$

Upon insertion of the l.h.s. of Eq. (4.5) for σ in the r.h.s. of Eq. (4.4), this equation can then be expressed as a complete derivative by writing

$$\partial_\xi \left[(v + E) \sigma + v \int_c^{|E|} dx \frac{f(|x|)}{x} \right] = 0 .\tag{4.6}$$

For physical fronts with $v > 0$ and $\sigma \geq 0$ [see (4.3)], we see from (4.5), that E is a monotonic function of ξ ,

$$\begin{aligned}\text{sgn} \left(\partial_\xi E(\xi) \right) &= \text{sgn} q(\xi) = \text{sgn} E(\xi) = \text{sgn} E^+ \\ &\text{for all } \xi .\end{aligned}\tag{4.7}$$

This allows us to use E as a coordinate instead of ξ . According to (3.7), before the front at $\xi \rightarrow \infty$ the electron density vanishes, so $\sigma^+ = \sigma[E^+] = 0$. Eqs. (4.6) and (4.7) together then determine σ as a function of E as

$$\sigma[E] = \frac{v}{v + E} \rho_{E^+}[E] ,\tag{4.8}$$

with the function

$$\rho_{E^+}[E] = \int_{|E|}^{|E^+|} dx \frac{f(|x|)}{x} = \rho_{|E^+|}[|E|] \quad (\geq 0) .\tag{4.9}$$

The function $\rho_{E^+}[E]$ is nothing but the ion density, as can be deduced by inserting $q = \partial_\xi E$ into (4.5) and equating the charge density q with $\rho - \sigma$. The ion density

ρ for $D = 0$ turns out to be a function of E and E^+ only, and to be independent of the particular front shape parametrized by v .

The fields σ , ρ and E as a function of ξ can be found by solving the implicit equation for $E = E(\xi)$

$$\partial_\xi \ln |E| = \frac{\rho_{|E^+|}(|E|)}{v + E}, \quad (4.10)$$

which can be derived from Eqs. (4.5) and (4.9).

Eq. (4.8) immediately shows that physically allowable solutions with $\sigma \geq 0$ and $v > 0$ must have $v + E \geq 0$ for all field values. Because of the monotonicity of E as a function of ξ , this is automatically satisfied for *PSF* with $E^+ > 0$, but for *NSF* this implies in particular that $v + E^+ \geq 0$; together with $v > 0$ we thus have for physical fronts

$$v \geq \max \left[0, -E^+ \right]. \quad (4.11)$$

In physical terms, the condition $v \geq -E^+$ expresses, that the velocity of uniformly translating fronts must be at least the drift velocity $-E^+$ of free electrons in the leading edge of the front, where the field is strongest. (Remember, that (4.7) implies, that the field is monotonic in space.)

For all values of v obeying the inequality (4.11) the solutions of (4.8) and (4.10) are proper, physically allowable solutions for fixed E^+ ; within the context of the present model, this illustrates a general feature of front propagation into unstable states, namely that there exists a family of front solutions parametrized by the velocity [42].

In Fig. 2(a), we plot the solution (4.8) for σ as a function of E for the fixed value of the velocity $v = 2$ in the case that the impact ionization function $f(E)$ is given by the Townsend expression $f_T(E)$ of (2.16) as in the numerical simulations [8,9]. Note that in this representation, the state behind the front at $\xi = -\infty$ corresponds to a point on the σ axis, and that the front solution $\sigma(\xi)$, $E(\xi)$ is represented in this diagram by the flow along one of the trajectories towards either the positive E -axis for *PSF* or the negative E -axis for *NSF* for $\xi \rightarrow \infty$. Note furthermore that σ overshoots the value $\sigma^- (= \sigma(\xi \rightarrow -\infty))$ in the case of *NSF*. This property as well as the monotonicity of $\sigma[E]$ and accordingly of $\sigma(\xi)$ for positive fronts, follows immediately from Eq. (4.8). For *NSF*, it can also be observed in the three-dimensional simulations of Vitello *et al.* [9], shown in Fig. 1(c).

The smallest E^+ for which a front solution with $v = 2$ is shown in Fig. 2(a), is $E^+ = -1.999$. For this value of E^+ , $\sigma[E]$ continues to increase till $E \approx E^+$ and then suddenly decays to zero. A short analytical investigation of (4.8) shows, that this behavior develops into a discontinuity of $\sigma[E]$ at the point $E = E^+$ for $v = -E^+$. $\sigma[E]$ then increases monotonically up to $f(E^+)$ for $E \downarrow E^+$ and then jumps to zero discontinuously at E^+ . This shock-like behavior stays unchanged under a parameter change

to $\sigma(\xi)$. It is further discussed and motivated in Section V.

An immediate consequence of Eqs. (4.8) and (4.9) for the electron and ion density is that the value σ^- behind the front (where $E \rightarrow 0$) is a simple function of the value E^+ of the field ahead of the streamer profile:

$$\sigma^-(E^+) = \rho_{|E^+|}[0] = \int_0^{|E^+|} dx \frac{f(|x|)}{x} \quad (4.12)$$

The virtue of this expression for the electron density σ^- far behind the front as well as of the expression (4.9) for the ion density ρ throughout the whole front, is that it is independent of the velocity v , hence independent of whichever front profile is selected, provided that the limit $D \rightarrow 0$ is smooth. We shall see later that this $D = 0$ result remains relatively accurate for *NSF* fronts with $D \lesssim 1$, and compare it to the results of the simulations [8,9] in Section VI. For *PSF*, on the other hand, the above result will turn out to be less relevant due to the non-perturbative nature of the limit $D \rightarrow 0$ in this case.

For the Townsend function $f_T(|x|)$ [Eq. (2.16)] the function $\sigma^-(E^+)$ can be expressed as

$$\begin{aligned} \sigma^-(E^+)_T &= |E^+| E_2(|E^+|^{-1}) \\ &= f_T(|E^+|) - E_1(|E^+|^{-1}), \end{aligned} \quad (4.13)$$

where $E_n(z)$ is the exponential integral [43].

We finally note that the second form of (4.13) shows that σ^- approaches f_T for large fields, since $f_T \gg E_1$ for $E^+ \gg 1$. For E^+ of order unity, σ and f_T are still of the same order, and this shows (for small D) that the growth rate (3.3) of long wavelength unstable modes in the unionized state is comparable to the damping rate (3.5) of stable modes in the plasma behind a *NSF*. For small fields, the strict bounds on E_2 [43] show that $\sigma^- \approx E^+ f_T(E^+)$, so that the approximate equivalence of these two time scales does not hold for $E^+ \ll 1$, but in the small field range our starting model is not very realistic anyway, because of the neglect of stabilizing recombination terms.

B. $D \neq 0$ Front Solutions

For $D \neq 0$, we can not obtain the uniformly translating solutions analytically. Moreover, perturbation theory around the $D = 0$ case is not simply possible, as D appears in front of the highest derivative in Eq. (4.2), so the diffusion term acts as a singular perturbation. As a consequence, Eqs. (4.2) reduce to a set of two coupled first order *ode*'s for $D = 0$, while three are required for $D \neq 0$. However, we can still easily demonstrate the existence of a one-parameter family of uniformly translating front solutions for $D \neq 0$ through standard counting arguments for *ode*'s. Building on the results of such an analysis,

the solutions can then be constructed by integrating numerically in a stable direction, using so-called “shooting methods” [44].

To perform the analysis, it is convenient to write the equations as a set of three coupled first order *ode*'s. There is some freedom for the choice of the third variable: The standard choice would be $\sigma' = \partial_\xi \sigma$, but for the discussion of the singular limit as well as for numerical stability, the charge density q has turned out to be the most convenient choice. The *ode*'s (4.2) then become

$$\begin{aligned}\partial_\xi \sigma &= -\frac{\sigma E - vq}{D}, \\ \partial_\xi E &= q, \\ \partial_\xi q &= -\frac{\sigma f(|E|)}{v} + \frac{\sigma E - vq}{D}.\end{aligned}\tag{4.14}$$

Just as we thought of the profiles for $D = 0$ as describing flow in a two-dimensional (σ, E) phase space, we can now think of Eqs. (4.14) as describing a flow in a three-dimensional (σ, E, q) phase space. The velocity v just plays the role of a parameter in the flow equations, while ξ again plays the role of a time-like variable — see the sketch in Fig. 2(b).

The steady states of the full *pde*'s discussed in Section III correspond to fixed points of the flow: the points $(\sigma, 0, 0)$ on the σ -axis are fixed points of the flow (4.14) and correspond to homogeneously ionized plasma states, while the E -axis is a line of fixed points $(0, E, 0)$ each of which corresponds to a non-ionized state with $\sigma = \sigma' = 0$ and $E \neq 0$.

A uniformly translating front solution now corresponds to the existence of a trajectory in this phase space that starts at “time” $\xi = -\infty$ on the σ -axis and flows to the E -axis for $\xi \rightarrow \infty$. The multiplicity of such solutions (i.e., whether they exist as discrete sets, or, e.g., as a one- or two-parameter family) can be determined as follows. If we linearize the flow near an arbitrary point $(\sigma^-, 0, 0)$ on the σ -axis by writing $(\sigma, 0, 0) = (\sigma^-, 0, 0) + A \exp(-\Lambda^- \xi)$, we find the eigenvalue equation

$$\Lambda^- \left(\Lambda^{-2} - \Lambda^- \frac{v}{D} - \frac{\sigma^-}{D} \right) = 0.\tag{4.15}$$

The fact that there is a zero eigenmode is a consequence of the fact that the σ -axis is a line of fixed points. For the two nontrivial eigenvalues (which correspond to the linearized modes (3.5) about the ionized and screened region by equating $i\mathbf{k} \cdot \hat{\mathbf{x}} = -\Lambda^-$ and $\omega^- = \Lambda^- v$) we have

$$\Lambda_\pm^- = \frac{v \pm \sqrt{v^2 + 4D\sigma^-}}{2D}.\tag{4.16}$$

The eigenvalue Λ_+^- is positive, and hence gives a decaying exponential; thus points along the corresponding eigendirection flow into the σ -axis as ξ increases. The eigenvalue Λ_-^- , on the other hand, is negative and hence corresponds to an unstable eigendirection, with flow away from the axis. This implies that at each point $(\sigma^-, 0, 0)$

on the σ -axis, there is, for fixed v , a unique eigendirection $(-\Lambda_-^-, 1, \Lambda_-^-)E_1$ along which the flow is away from the axis. This flow can be followed in two anti-parallel directions, determined by the sign of E_1 . The one flowing towards positive values of E is the beginning of a *PSF* front profile, the one flowing towards negative E is the beginning of an *NSF* profile. From these eigendirections, one derives that for *PSF* with field perturbations $E_1 > 0$, the electron density decreases close to the σ axis, while for *NSF* it increases. Accordingly, before reaching $\sigma = 0$ for $\xi \rightarrow \infty$, a *NSF* profile has at least one maximum of σ , while a negative one can be (and is) monotonic. This generalizes our result for $D = 0$, and is consistent with the findings of Vitello *et al.* [9] shown in Fig. 1(c). The physical origin of the maximum of σ in the rear end of the *NSF* profile is the screening of the field: Due to the low ionization rate in an already fairly suppressed field the ion density has already almost acquired its final value, so the electron density has to overshoot its asymptotic value σ^- so as to make $\partial_\xi E < 0$. The screening behind a *PSF* happens by suppressing the electron density faster than the ion density for increasing ξ , and so there σ is monotonic.

Let us now investigate the stability of the flow near a point $(0, E^+, 0)$ on the E -axis. Upon linearizing the flow equations (4.14) and writing the ξ dependence of the perturbations in the form $\exp(-\Lambda^+ \xi)$, we find the eigenvalue equation

$$\Lambda^+ \left(\Lambda^{+2} - \Lambda^+ \frac{v + E^+}{D} + \frac{f(|E^+|)}{D} \right) = 0.\tag{4.17}$$

Again, there is a zero eigenvalue due to the fact that the whole E -axis is a line of fixed points. The two nontrivial eigenvalues are

$$\Lambda_\pm^+ = \frac{v + E^+ \pm \sqrt{(v + E^+)^2 - 4Df(|E^+|)}}{2D}.\tag{4.18}$$

These eigenvalues can be related to (3.3) in the same way as (4.15) could be related to (3.5). For $v + E^+ > 0$, the real parts of these eigenvalues are always positive, so that both eigendirections are stable. In other words, for $E^+ > -v$, all points near the E -axis flow towards this axis — in slightly more technical terms: there is a two-dimensional stable manifold flowing into each of these points on the E -axis. For $E^+ < -v$, the flow is away from the E -axis, and fronts with $v + E^+ < 0$ can not be constructed. This generalizes (4.11) to $D \neq 0$.

The existence of a one-parameter family of fronts with velocity $v > -E^+$ can now simply be understood as follows. As we saw before, there is one unique *PSF* and one unique *NSF* trajectory flowing out of each point on the σ -axis for fixed v and D . Since the flow defined by Eqs (4.14) is continuous, we can expect each trajectory to extend smoothly [45]. Once the flow gets near the E -axis, we know from the above analysis that the trajectory will be attracted completely to the axis, provided

v is large enough. Thus, for each σ^- and v , there will exist two unique trajectories, i.e., a unique *PSF* solution and a unique *NSF* solution. Since each of these trajectories flows into a unique point on the E -axis, the flow equations implicitly define a unique relation of the form $\sigma^- = \sigma^-(v, E^+)$ for each of the two types of fronts. For a given value of E^+ , we thus have a one-parameter family of front solutions, parametrized by v .

There are two important properties of the front solutions associated with their asymptotic large ξ behavior. First of all, we note that according to (4.18) the eigenvalues Λ_{\pm}^{\pm} are only real for

$$v \geq v^* \equiv -E^+ + 2\sqrt{Df(|E^+|)}. \quad (4.19)$$

This implies that the corresponding front profiles can certainly not approach the asymptotic state $\sigma = 0$ ahead of the front in a monotonic way for $v < v^*$: When the eigenvalues are complex, the front profiles have an oscillatory tail of the form $\exp[-(\Re\Lambda^+)\xi] \cos[(\Im\Lambda^+)\xi]$. Clearly, this violates the physical condition that the electron density σ should remain positive, so solutions with $-E^+ < v < v^*$ are physically excluded: v^* denotes, in the present case, the smallest velocity of physically allowable uniformly translating front solutions.

The identification of v^* as a bound on the velocity of physically allowed front profiles depends only on the structure of the eigenvalues Λ^+ associated with the *linear* flow near unstable states. There is a second, *nonlinear*, way in which the range of physically allowed values of v can be bounded. To understand this, note that for any $v \geq v^*$, the asymptotic decay of $\sigma(\xi)$ for $\xi \rightarrow \infty$ for a uniformly translating profile will be

$$\sigma(\xi) = A_- e^{-\Lambda_+^+ \xi} + A_+ e^{-\Lambda_+^+ \xi} + \text{h.o.t.} \quad (4.20)$$

with real coefficients A_- and A_+ . Here, h.o.t. stands for higher powers of the two exponentials generated when expanding the equations to higher than linear order in the variables. Clearly, the smallest eigenvalue Λ_+^+ governs the asymptotic decay of the profile provided $A_- \neq 0$. That A_- will generically be nonzero for an arbitrary velocity v follows again from the counting argument above for the flow in phase space: Each *PSF* and *NSF* trajectory flowing out of a point on the σ -axis is unique, and hence there is no freedom to impose an additional condition $A_- = 0$ close to the E -axis. Furthermore, the coefficients A_- and A_+ depend on the full global nonlinear behavior of the flow, and hence they depend implicitly on v .

There might exist, however, particular velocities $v^{part} > v^*$, for which

$$A_-(v^{part}) = 0. \quad (4.21)$$

For discussing these we invoke again a continuity argument for the front properties as a function of v . We expect a very slowly decaying, nearly homogeneous uniformly translating front solution to have a non-negative

density everywhere, and to have a very large velocity, since the velocity of a profile is essentially inversely proportional to its slope in the limit of small slopes. (So indeed the roots Λ_-^- given by (4.16) and Λ_+^+ given by (4.18) vanish in the limit that v becomes large.) So for large v we expect to find physical solutions. These are characterized by $A_- > 0$ in the leading edge of the front. Decreasing v continuously, we either reach $v = v^*$ smoothly with still $A_- > 0$, or we reach the first particular velocity, v_1^{part} , where A_- vanishes. In the latter case, we expect by continuity $A_-(v) < 0$ for $v < v_1^{part}$. This implies that then σ approaches zero from below, i.e., that the front solution is unphysical. Below the next v_2^{part} , we expect the electron density to develop two zero's and so forth. The largest v^{part} , if it exists, thus plays the role of the *nonlinear front velocity* v^\dagger [24],

$$v^\dagger = \max \{ v^{part} \mid A_-(v^{part}) = 0 \} \quad (4.22)$$

for a given E^+ . (Note that if $\Lambda_+^+ < 0.5\Lambda_+^+$, the higher order terms in (4.20) of order $\exp(-2\Lambda_+^+ \xi)$ are actually larger than the second term $\exp(-\Lambda_+^+ \xi)$. This does not change our argument, though, as the prefactor of this second order term will vanish if A_- vanishes.)

At the velocity $v = v^\dagger$ or at any $v = v^{part}$, the flow in phase space approaches the E -axis along the eigenvector where the flow is most rapidly contracting. The trajectory corresponding to the nonlinear front solution is therefore more appropriately referred to as a strongly heteroclinic orbit, where heteroclinic indicates that it is a trajectory from one fixed point to another one. The existence and properties of strongly heteroclinic orbits have recently been under active investigation [46].

Such a velocity v^\dagger , if it exists, bounds the continuum of velocities of physical uniformly translating solutions from below, and thus replaces the earlier bound v^* derived from linearizing the equations in the leading edge of the front.

C. Nonlinear Front Solutions for PSF

For *NSF*, the bounding velocity v^* given by (4.19) is always positive. Moreover, by integrating the flow equations (4.14) numerically and searching for particular solutions for which, according to (4.21), $A_-(v^{part}) = 0$, we have convinced ourselves that there are no such solutions for any $D \neq 0$ and $E^+ < 0$. Hence, the smallest velocity of physical *NSF* solutions is always v^* , for any value of the parameters.

For *PSF*, on the other hand, the situation is very different, since $v^* < 0$ for $(E^+)^2 > 4Df(|E^+|)$ — for the Townsend function (2.16), this happens for $D \leq 0.25E^+e^{1/E^+}$, hence for any E^+ for $D \leq 0.68$. In particular for *PSF* at small D the question therefore arises whether there are nonlinear front solutions defined by

(4.21) and (4.22) with $v^\dagger > 0$. The results of a numerical search for such solutions are shown in Fig. 3, as a function of D and E^+ . Below the full line in this diagram, there exists indeed a nonlinear front $v^\dagger > v^*$, whereas above this line v^* denotes the smallest velocity of physical front solutions. While these results have been obtained numerically, the existence of a single (unique) particular solution with $A_-(v^{part}) = 0$ in the limit $D \rightarrow 0$ can be demonstrated analytically. Since a full discussion of these results will be given elsewhere [47], we confine ourselves here to a brief outline of the arguments that also demonstrate the singular nature of these solutions for $D \rightarrow 0$.

If we take the limit $D \rightarrow 0$ with v fixed, assuming no nontrivial scaling of the variables σ , E and q and of the spatial coordinate ξ , Eqs. (4.14) can easily be shown to reduce to those studied in Section IV.A for $D = 0$. Hence, we can recover in this way the family of front solutions obtained there. *Any* particular solution, on the other hand, for which $A_-(v^{part}) = 0$, decays according to (4.20) as $\exp(-\Lambda_+^+ \xi)$ as $\xi \rightarrow \infty$. Since $\Lambda_+^+ \propto D^{-1}$ for $D \rightarrow 0$, such a particular front solution becomes extremely steep as $D \rightarrow 0$: its gradients diverge as $1/D$ and so that the diffusion term can still overcome the drift term as $D \rightarrow 0$. That the velocity of such a solution must also have a nontrivial scaling in this limit can be seen from the third equation of (4.14), written in the form

$$\partial_\xi q = \sigma \left(-\frac{f(|E|)}{v} + \frac{E}{D} \right) - \frac{v}{D} q. \quad (4.23)$$

Any *nontrivial* scaling of this equation in the limit $D \rightarrow 0$ can only occur if the first term between brackets remains of the same order as the other two, which diverge as $1/D$. This is only possible if v scales as D . In this limit, the third term can then be neglected, and since $\partial_\xi q$ has to change sign in the front region (as the charge density q vanishes as $\xi \rightarrow \pm\infty$), there must be an intermediate value $\hat{E} < E^+$ of the field for which $v = Df(|\hat{E}|)/\hat{E}$.

Now that we know the scaling of the spatial gradient of the velocity of such particular front profiles for $D \rightarrow 0$, one easily convinces oneself that the electron and charge density of these solutions must *diverge* as $1/D$ in this limit. To study the existence of such possible solutions, it is therefore convenient to introduce new variables and coordinates according to

$$x = D\tilde{x}, \quad v = D\tilde{v}, \quad \xi = D\tilde{\xi}, \quad \sigma = \tilde{\sigma}/D, \quad q = \tilde{q}/D, \quad (4.24)$$

with E and τ unchanged. In these new variables, the flow equations (4.14) become

$$\begin{aligned} \partial_{\tilde{\xi}} \tilde{\sigma} &= -\tilde{\sigma} E + D \tilde{v} \tilde{q}, \\ \partial_{\tilde{\xi}} E &= \tilde{q}, \\ \partial_{\tilde{\xi}} \tilde{q} &= \tilde{\sigma} \left(E - \frac{f(|E|)}{\tilde{v}} \right) - D \tilde{v} \tilde{q}. \end{aligned} \quad (4.25)$$

The limit $D \rightarrow 0$ can now be taken simply by leaving out the term $D\tilde{v}\tilde{q}$ in the first and last equation. We will show elsewhere [47] that the resulting equations have *one unique* physical front solution thus fixing one particular value of the scaled velocity \tilde{v}_1 and in view of the scaling (4.24) and the scaling of the eigenvalues Λ_\pm^+ , this solution must have $A_-(\tilde{v}_1) = 0$. This solution is therefore precisely the $D \rightarrow 0$ limit of the nonlinear front solution with velocity $v^\dagger = \tilde{v}_1 D$. Furthermore, since the limit $D \rightarrow 0$ is smooth for Eqs. (4.25), this shows that there exists a nonlinear front solution with $v^\dagger > 0$ for *any* E^+ and nonzero but small D . Due to the singular scaling (4.24), the corresponding front solutions are determined by *ode's* that have a different structure from those studied for $D = 0$ in Section IV.A, and therefore these nonlinear front solutions can not be obtained perturbatively from the latter class of solutions — of course, the latter class of solutions still exists for $D \neq 0$, in agreement with the counting arguments given earlier, but these now correspond to a singular limit of Eqs. (4.25)! The significance of these nonlinear front solutions lies in the fact that they will turn out to be the selected fronts that dominate the dynamics of *PSF* in the physically important range $0.1 \lesssim D \lesssim 0.3$.

The nonlinear front solution can be constructed numerically very easily by integrating Eqs. (4.25) using standard numerical “double shooting” routines [44]. Fig. 4 shows our numerical results for the smallest physical velocity, $\max(v^\dagger, v^*)$ in the case that the ionization function is given by the Townsend expression. The scaled velocities v^\dagger/D and v^*/D are plotted; in agreement with our arguments above the scaled velocity v^\dagger/D of the nonlinear front solution approaches a finite limit as $D \rightarrow 0$. Furthermore, the ratio v^\dagger/D hardly varies with D in the physical range $0.1 \lesssim D \lesssim 0.3$, and for small fields E^+ , the scaled velocity v^\dagger/D becomes exponentially small, in agreement with the bound $v^\dagger/D < E^+ \exp(-1/E^+)$ that follows from the observations discussed after Eq. (4.23) above.

We finally note that our numerical routines have not only allowed us to obtain the results show in Figs. 3 and 4, but have also enabled us to verify numerically all the statements made above about the multiplicity of solutions, the parameter ranges for physical fronts, the monotony properties, the singular behavior of the small D *PSF* limit, and the persistence of the family of front solutions for $D \rightarrow 0$.

V. SELECTION OF THE ASYMPTOTIC FRONT

A. Front Propagation into Unstable States

We have seen that the non-ionized state into which the streamer fronts propagate, is an unstable state, that the homogeneous weakly ionized plasma is a stable state, and

that there is a family of uniformly translating front solutions connecting the two. The existence of a family of front solutions is a generic feature of front propagation into unstable states. We, therefore, briefly recall what is known in the literature for analogous problems and then translate this experience to the streamer problem. The prototype equation for studies of this type of front propagation is

$$\partial_t u = \partial_x^2 u + g(u) , \quad (5.1)$$

where $g(u)$ is some nonlinear function which satisfies

$$g(0) = 0 , \quad g(1) = 0 , \quad g'(0) > 0 , \quad g'(1) < 0 . \quad (5.2)$$

Note that these relations imply that the “state” $u = 0$ is unstable, and that the “state” $u = 1$ is stable. The study of the propagation of fronts into the unstable state $u = 0$ in this equation dates back to the early work of Kolmogorov *et al.* [33] and Fisher [34] in the context of population dynamics. Later Gel’fand [48] studied a particular example of a function $g(u)$ motivated by combustion. The mathematical research on this equation culminated in the work by Aronson and Weinberger [31], who rigorously solved the front propagation problem for (5.1). In particular, they proved that any initial perturbation, that is nonzero only in a finite part of space, approaches a unique uniformly translating front solution with velocity v_f in the long time limit. If $g''(u) < 0$ for all u , v_f equals $v^* = 2\sqrt{g'(0)}$ (derived from linearizing in the tip of the front), while for general $g(u)$, v_f approaches either v^* or some $v^\dagger > v^*$. We refer to the literature for a detailed discussion of this work [31,32].

The velocities v^* and v^\dagger of the above problem directly correspond to our v^* (4.19) and v^\dagger (4.22), since they are also the smallest velocities, which still allow for uniformly translating fronts with $u \geq 0$ everywhere. So if u is interpreted as a population density or a chemical concentration, the selected front for every sufficiently localized initial state is the slowest physical uniformly translating front. In other interpretations no physical constraints bind u to positive values. Nevertheless the selected velocity stays the same. In this case, one can prove that every front with smaller velocity is dynamically unstable [21], i.e., that the selected front is marginally stable. The slowest physical or stable solution, which is selected, coincides with the steepest physical or stable one.

In the last decade, it has been recognized that several aspects of the front selection problem encountered for the nonlinear diffusion equation (5.1), seem to have more general validity. Certain scenarios, justified by heuristic arguments but lacking a detailed mathematical proof, were formulated and numerically tested on more complicated *pde*’s that were often of higher order in the spatial derivatives [49,21–25]. Some of the equations studied lead to non-uniformly translating fronts that leave a nontrivial spatially periodic state behind [49,21,23,24,50]. A particular scenario is the one distinguishing between the so-called *linear marginal stability*

regime where $v_f = v^*$ and the *nonlinear marginal stability* regime where $v_f = v^\dagger$ [21–24]. These names stem from the fact that in this formulation, the two regimes of front selection are related to the stability properties of the front solutions — in both cases, the selected front separates stable front solutions from unstable ones. Applied to (5.1), this scenario just provides an intuitive explanation of all the well-known mathematical results. For plasma physicists, it is worth mentioning that dynamics in the linear marginal stability regime is related to that determined by the “pinch point analysis” which was developed in plasma physics in the late fifties [51,52,23].

B. Predictions for Streamer Fronts

By extending the arguments in the appendix of [23], one may show that in the streamer case just like in the case of the above problem (5.1), all physical solutions, i.e., all solutions with $u \geq 0$ resp. $\sigma \geq 0$ everywhere, are stable. For a detailed discussion, we refer to [47]. It can be argued [22,23], and proven for (5.1) [21], that a sufficiently localized initial condition will approach the physical uniformly translating front, which is closest in “phase space”, i.e., the steepest one. Both for (5.1) and for the streamer equations, the steepest uniformly translating physical front is uniquely defined. It is also the slowest one.

We can immediately prove this when initially $\sigma(x, \tau = 0) = 0$ for $x > x_c$ for streamer fronts with $D = 0$: In general, there is a front solution for every $v \geq \max[0, -E^+]$, but now the only way in which the electrons can enter the range $x > x_c$ is through electron drift with velocity $-E^+$. Clearly, therefore, the asymptotic front speed of a *NSF* can only be $-E^+$, while a *PSF* can not propagate at all. If the initial electron density, however, decays exponentially, the local electron density grows by drift and ionization, and the front can move quicker than $-E^+ > 0$ for a *NSF*.

For $D \neq 0$, we will here only conjecture the analogous statements, and we will test them numerically in Section VI:

1. *Selected front velocity.* If the initial conditions are sufficiently localized, *the selected front is the slowest physically acceptable front solution*, i.e., the slowest front profile for which $\sigma(\xi) \geq 0$ for all ξ . In view of the discussion of Section IV, this means that the selected front velocity v_f is predicted to be

$$v_f = v^* = -E^+ + 2\sqrt{Df(|E^+|)} , \quad (5.3)$$

except when there exists a nonlinear front solution satisfying (4.22): In that case

$$v_f = v^\dagger . \quad (5.4)$$

Note that the result (5.3) (v^* is the *Linear Marginal Stability* value in the terminology of [22,23]) is an explicit expression for v_f in terms of parameters associated with the linear instability of the unstable state only. On the other hand, the existence of a nonlinear front and the value of v^\dagger (the *Nonlinear Marginal Stability* value) depends on the whole nonlinear behavior of the flow equations (4.14).

2. *Localized initial conditions.* Initial conditions are sufficiently localized if their spatial decay is faster than the asymptotic decay associated with the smallest eigenvalue of the selected profile, i.e., if

$$\sigma(x, \tau = 0) < C e^{-\Lambda_-^+(v^*)x}, \quad \text{or} \quad (5.5)$$

$$\sigma(x, \tau = 0) < C e^{-\Lambda_-^+(v^\dagger)x}, \quad (5.6)$$

for $x \rightarrow \infty$,

depending on whether the selected front is v^* or v^\dagger . Here C is an arbitrary constant, and $\Lambda_\pm^+(v^*)$ ($= \Lambda_\pm^+(v^*)$) and $\Lambda_\pm^+(v^\dagger)$ are given by (4.18).

3. *Non-localized initial conditions.* If an initial condition does not obey (5.5) or (5.6), faster front speeds are possible. In particular, if initially $\sigma(x, \tau = 0) \sim \exp(-\Lambda x)$, with $\Lambda < \Lambda_-^+(v^*)$ or $\Lambda < \Lambda_-^+(v^\dagger)$, whichever regime applies, then the front speed is given by

$$v = -E^+ + D\Lambda + \frac{f(|E^+|)}{\Lambda}, \quad (5.7)$$

which is obtained by solving (4.17) for v in terms of Λ .

We now combine the analytic and numeric findings from Section IV with the selection rules above to quantitative predictions for asymptotic fronts, which evolve from sufficiently localized initial conditions, in the case that the impact ionization is given by the Townsend expression (2.16):

NSF: For *NSF*, we numerically have not found any nonlinear fronts for any D and E^+ , so our simple yet powerful prediction is that for *NSF* $v_f = v^*$ with v^* given by (5.3). In principle it is possible that for other ionization functions $f(E)$ than the Townsend function (2.16), there can be nonlinear front solutions also in the *NSF* regime. In practice, we expect, however, that this will not be the case for physically reasonable functions $f(E)$, i.e., for functions consistent with (2.17).

Once the predicted velocities are known, the value σ^- of the electron density behind the streamer head is obtained from the numerical integration of the flow equations. The results of these calculations are shown in Fig. 5(a). Since for *NSF*, the limit

$D \rightarrow 0$ is smooth, also σ^- depends only weakly on D for $D \lesssim 1$, so that the $D = 0$ prediction (4.13) is quite accurate for realistic values of the diffusion coefficient.

At the predicted values of the selected front velocity, the width of the front region can be obtained directly from our numerical solutions of the flow equations. We have somewhat arbitrarily defined the width w as the distance between the points where σ is 90% and 10% of the value σ^- . As Fig. 6 shows for *NSF* fronts with $D = 0.1$, this front width is typically of order 3 for field values of order unity. This confirms again that in the small D limit the impact ionization length α_0^{-1} sets the inner scale of streamer fronts. Furthermore, we find that our numerical data are well fitted by the expression $w \approx 6/\Lambda_\pm^+(v^*)$, which shows that the front width simply scales with the spatial decay rate $\Lambda_\pm^+(v^*) = \Lambda_\pm^+(v^*)$ of the streamer profile in the leading edge. *NSF* fronts always have a maximum of the electron density within the front.

PSF: As we saw in Section IV, for *PSF* with $D \lesssim 0.9$, there always is a nonlinear front solution with velocity $v^\dagger > v^*$. The prediction is that in this range the selected front solution is the nonlinear front solution, i.e., $v_f = v^\dagger$. Values of v^\dagger as a function of D and for several values of E^+ were already given in Fig. 4. We also saw before that these nonlinear front solutions are singular in the limit $D \ll 1$, where $v^\dagger \approx D\tilde{v}^\dagger(D=0)$ and $\sigma^- \approx \tilde{\sigma}^-(D=0)/D$. The resulting predictions for σ^- are shown in Fig. 5(b).

The fact that the dimensionless inner decay length of these nonlinear fronts scales as D , implies that the physical decay length of such solutions is $D/\alpha_0 = D_e/(\mu_e E_0)$, i.e., is given by the electron diffusion length. However, since simultaneously the electron density σ^- diverges as $1/D$, the total front width w defined above still approaches a finite limit as $D \rightarrow 0$ in units of the ionization length α_0^{-1} .

We finally note that the front propagation problem posed by the one-dimensional streamer equations has a number of interesting differences and similarities with the Aronson Weinberger front propagation problem (5.1). In particular, it can be hoped, that techniques of strict bounds developed for the time development of these fronts [31] as well as for the nonlinear front velocity v^\dagger [46] might be also applicable to planar streamer fronts.

VI. NUMERICAL TESTS OF THE PREDICTIONS

We have tested the predictions listed in Section V by numerically integrating the *pde*'s (3.10), (3.11) forward in time. Our computer program is a finite difference code

with a time integration which is based on a semi-implicit method.

We have performed an extensive search through parameter space, varying D between 0.02 and 3, and $|E^+|$ between 0.3 and 10. All our numerical studies of the dynamics fully confirm our predictions for fronts, and therefore we only present a sample of our results that illustrate the important features.

All the simulations of the initial value problem we present in the remaining figures, have initially a field $E = -1$ constant in space. We keep the field constant in time in the non-ionized region. The simulations of Fig. 7 – 10 start with the same localized initial ionization seed, a Gaussian profile for the electron density, $\sigma(x, t = 0) = 0.01 \exp -(x - x_0)^2$. Fig. 7 shows a run for $D = 1$ and times $t = 0 - 130$ in time steps $\Delta t = 2$. As can be seen, the small ionization seed near $x_0 = 50$ initially grows while drifting to the right in accord with Eq. (3.4). At time $t = \mathcal{O}(20)$, the ionization is strong enough that field saturation sets in and two asymmetrically propagating fronts emerge. The one propagating to the right develops into a uniformly propagating *NSF* with velocity $v^* = 2.21$ [53] and degree of ionization behind the front $\sigma^- = 0.130$. The maximum value of σ in the rear part of the front is $\sigma_{max} = 0.150$. At the same time, a structure develops on the left, which at time $t = 130$ not yet has reached a stationary form, and which eventually will develop into a *PSF*. (Note that propagation to the left into a negative field $-E^+$ corresponds to a *PSF* front moving to the right towards $x \rightarrow \infty$ in a field $+E^+$). How the *PSF* actually reaches a uniformly translating profile, is shown in Fig. 8, where the development for $x_0 = 150$ and otherwise identical initial and boundary conditions is followed in time steps of $\Delta t = 10$ during the time $t = 0 - 500$. An asymptotic velocity of $v^\dagger = 0.22$ and a degree of ionization $\sigma^- = 0.43$ is reached. Note the huge difference in the degree of ionization and in the front velocity already for the unrealistically large value of the diffusion constant $D = 1$.

The predictions from Section V for the selected uniformly translating fronts for $D = 1$ and $E^+ = \pm 1$ yield for the *NSF* $v^* = 2.213$ and $\sigma^- = 0.129$, and for the *PSF* $v^\dagger = 0.2199$ and $\sigma^- = 0.432$. They thus correctly predict the simulations of the initial value problem shown in Figs. 7 and 8 within the accuracy given. Note that for the velocity v^\dagger of the *PSF* and for the degrees of ionization σ^- both behind the *PSF* and the *NSF* this fact also shows the relative accuracy of the two very different numerical methods used, while for the velocity v^* of the *NSF* the numerical integration of the initial value problem exactly reproduces the analytic result.

As D decreases, both the structures within the fronts and the asymmetry between *NSF* and *PSF* become more pronounced. We illustrate this in Figs. 9 and 10 with the temporal development starting from the same initial perturbation as before, but now for $D = 0.1$, the value corresponding to the parameter values of the earlier three-dimensional simulations [8,9]. The time

ranges in each plot are chosen appropriately for seeing the *NSF* and the *PSF* evolve into a uniformly translating state. Fig. 9 shows a perturbation (initially localized at $x_0 = 50$) evolving during time $t = 0 - 130$ in time steps $\Delta t = 2$. Except for the smaller diffusion constant and the stretched x axis, the situation is thus identical with that of Fig. 7. The *NSF* on the right propagates with a somewhat smaller velocity $v^* = 1.39$, leaving a slightly higher ionization $\sigma^- = 0.147$ behind. The maximum $\sigma_{max} = 0.199$ is relatively higher, since diffusional smoothing of structures is less pronounced. On the time scales of Fig. 9, the left front does not propagate, but retracts into an apparently immobile structure. The electrons drift with the field into the ionized region, leaving a layer of screening ions behind. Thus the electrons and the field are almost separated such that ionization on this side almost cannot occur. Eventually few electrons will reach the nonzero field region by diffusion and slowly build up a higher ionization and ultimately a propagating *PSF*. That a *PSF* actually emerges, is shown in Fig. 10. Only times $t = 4000 - 8000$ in time steps of $\Delta t = 100$ after the initial perturbation at $t = 0$ and $x_0 = 60$ are shown. The front propagates with velocity $v^\dagger = 0.0149$ leaving behind an ionization $\sigma^- = 6.32$. The numerical values predicted in Section V are $v^* = 1.384$ and $\sigma^- = 0.144$ for *NSF*, and $v^\dagger = 0.0146$ and $\sigma^- = 6.234$ for *PSF*. The remaining numerical discrepancy of maximally 2% could be resolved by choosing a still smaller gridsize in Figs. 9 and 10. Comparison of the *PSF* for $D = 1$ and $D = 0.1$ indicates that the time it takes such a front to build up, rapidly increases with decreasing D , but we have not pursued the scaling of the transient time with D .

We finally show in Fig. 11 the evolution of streamer fronts starting from non-localized initial conditions, i.e., not obeying the bounds (5.5) or (5.6) for $D = 0.1$. We used an initial electron density profile $\sigma(x, t = 0) = 0.01/(2 \cosh \Lambda(x - 200))$ with $\Lambda = 0.25$ and an initial field $E = -1$. At these values, for the *NSF*, $\Lambda^+(v^*) = 1.918$ and for the *PSF*, $\Lambda^+(v^\dagger) = 0.3766$. In this case, the bounds (5.5) or (5.6) are indeed violated for both fronts, and Eq. (5.7) predicts a *PSF* with velocity $v = 0.497 > v^\dagger = 0.0146$ and an *NSF* with velocity $v = 2.497 > v^* = 1.384$. The simulations find the fronts propagating with velocities 0.50 and 2.50, respectively. The ionization behind the *NSF* is $\sigma^- = 0.149$ and behind the *PSF* $\sigma^- = 0.158$, so that now both are comparable to each other and to $\sigma^-(D = 0) = 0.1485$ found from Fig. 5(a). Note that the diffusion constant is identical with that of Figs. 9 and 10, the only difference being the extended initial perturbation.

The simulations confirm that streamer front propagation is indeed correctly described by the marginal stability scenario, which in the present case amounts to the statement that the slowest physical velocity is selected, whenever one starts from sufficiently localized initial conditions, just as for the simpler case (5.1).

VII. CONCLUSIONS AND OUTLOOK

The analysis in this paper fully supports the validity of an effective interface approach suggested by the results of the full three-dimensional simulations of Dhali and Williams and of Vitello *et al.* [8,9]. This emerges from our detailed study of the associated one-dimensional problem, which yields the following results:

- (a) After a very brief stage of transient exponential amplification of the initial ionized seed, the growing streamer evolves into an electrically screened plasma body separating two narrow fronts which propagate into the non-ionized outer region. We show that these two fronts correspond, for all practical purposes, to translating profiles which propagate independently. This entails that the separation of spatial scales between an inner front and an outer one, set by the global geometry, is indeed justified.
- (b) This enables us to draw upon the existing knowledge about front propagation into unstable states and thus to provide definite predictions about:
 - the relationship $v_f(E^+)$ between the velocity of a planar streamer front and the value of the electric field ahead of it, and
 - the value of the degree of ionization of the plasma created by the front, $\sigma^-(E^+)$.

These predictions, although only valid as such in the absence of front curvature, still compare very favorably with the numerical results of Ref. [9]. The two values of $\sigma^-(E^+)$ on the axis of Fig. 1(a) and 1(b) behind the curved fronts of the 3D simulations [9] (with the convention that E^+ should be understood as the electric field value extrapolated from the external non-ionized region to the front position) are plotted in Fig. 5(a). Without adjustable parameters our one-dimensional predictions for $\sigma^-(E^+)$ are well within a factor of 2 from the 3D simulations. Likewise, the velocity values for $v_f(E^+)$ even agree to about 20%.

Moreover, our analysis shows that *NSF* and *PSF* propagate in this model and for realistic values of the reduced diffusion coefficient D , in a very asymmetric manner:

- *NSF* rapidly reach a regime of uniform propagation — typically, on the scale of several tens of time units, i.e., in less than 10^{-10} s. Their velocity is slightly larger than the electron drift velocity in the field E^+ .
- This is to be contrasted with the dynamics of *PSF*: For realistic D -values, of order 0.1-0.3, they approach uniform translation considerably more slowly than *NSF* — typically on the time scale of 10^{-8} s. Moreover, their asymptotic velocity is also much smaller than v_f^{NSF} . It obeys the inequality $v_f^{PSF} < DE^+ \exp(-1/E^+)$ [47]. Finally, while the widths of *PSF* and *NSF* are comparable,

the degree of ionization behind *PSF* is much larger (up to a hundred times for $D = 0.1$) than that behind *NSF*.

These results answer the question of whether *PSF* do or do not propagate, while explaining why the simulations of Vitello *et al.* [9] could not yield a definite answer — most probably, because, although their total width is of order α_0^{-1} , their true inner length scale, as defined by the steepness of the profile, was too small to be resolved by their grid size. (Note that the apparent symmetry between *PSF* and *NSF* found in earlier simulations [8] is to be related to the fact that there propagation into a pre-ionized medium (with initial electron density of $10^8/\text{cm}^3$) is studied, and possibly also due to the use of a poorly resolving grid.)

It was observed empirically by Dhali and Williams [8] that in the three-dimensional simulations, the dielectric relaxation time in the plasma behind the front was of the same order as the intrinsic time scale set by the front motion. Our analysis shows, that this was no accident: It is a manifestation of the fact that the balance of the growth mechanism (impact ionization) and the stabilization mechanism (screening) leads to a single time scale $t_0 = (\alpha_0 \mu_e E_0)^{-1}$ for a *NSF* and for the relaxation behind it for fields of order E_0 . Since our dimensionless value of σ^- is the inverse dielectric relaxation time, it is of order unity (or slightly smaller) for fields $E^+ \approx -1$.

Of course, the above results should only be considered as a first step towards a realistic treatment of streamer propagation. They will have to be developed and extended along two different directions:

(i) *Predictions of patterns within the present model and comparison with the simulations:* Within the frame of the present continuous and fully deterministic model, we here have only considered the restricted case of a one-dimensional geometry. This enabled us to demonstrate that the concept of effective interfaces does apply to streamers. This approach will now have to be extended to the description of curved fronts. As also discussed in [19], one will then be equipped with a reduced formulation, valid on the outer scale, which will permit us to study real three-dimensional streamers as pattern-forming systems, as was done, e.g., for viscous fingers and dendritic solidification fronts [20]. This should provide a direct approach to the question of dielectric patterns, alternative to the phenomenological *DLA*-inspired dielectric breakdown models [54].

(ii) *Possibly, extensions of the model will be necessary to predict real experiments:* We have based our analysis on the minimal model as defined in Section II. It contains several restricting simplifications. A first step in the improvement of the model would be to include the field dependence of the transport coefficients D_e and μ_e . It is clear that this will not modify our qualitative analysis, as, e.g., the counting argument for the existence of front solutions in Section IV depends only on the linearization about the stable and unstable states. Moreover, the qualitative asymmetry between the *NSF* and *PSF* will

persist as these result from the asymmetry of the electron drift. Quantitatively, the value of v^* , the selected value of NSF , will simply be given by (5.3) with the transport coefficient and ionization rate evaluated at the field value E^+ . The slow transient build-up and small speed of PSF could be affected quantitatively by ionic motion, but from this effect, we expect no major qualitative differences.

Finally, it should be kept in mind that our continuum equations are only valid on length scales larger than the mean free path ℓ_{mfp} . On the other hand, we find for the strongest field values appearing in the simulations (which are much larger than the values of the field across the gap, due to the enhancement near the streamer tip), that the front width decreases down to about $3\alpha_0^{-1} \approx 3\ell_{mfp}$ in the approximation (2.7). In such limits, nonlocality of the transport and ionization effects begin to play a role. In addition, under these conditions, a typical volume of size ℓ_{mfp}^3 contains only of the order of 1000 electrons for the parameter values (2.8) used in the simulations. Fluctuations are then likely to become non-negligible. In principle, treating these effects calls for a full kinetic description. This is probably out of reach for the moment, but one might want to mimic the main features of these effects by introducing stochastic terms in the equations. These also could mimic photo-ionization somewhat before the front due to photons released in the impact ionization events, or the natural homogeneous background ionization due to radioactivity and cosmic radiation. Investigation of their relevance for branching of dielectric breakdown patterns might help to understand the asymmetry between the macroscopic patterns of discharges propagating into a positive or a negative field [55].

In conclusion, our analysis opens the way to a microscopically based interface approach to discharges that seems promising for building a coherent framework for the analysis of breakdown patterns of various degrees of complexity.

ACKNOWLEDGMENTS

WvS gratefully acknowledges hospitality of the Université Paris VII, where this work was initiated. UE thanks F. Döbele and A. Stampa for valuable discussions about streamer experiments and plasma physics. Her work was made possible by the Priority Program Non-Linear Systems of the Dutch Science Foundation NWO. We also gratefully acknowledge financial support by NWO and the Lorentz Fund for visits of UE and WvS to the Université Paris VII. Finally, we like to thank P.A. Vitello for making copies of figures from [9] available, which appear here as Fig. 1.

APPENDIX: DIFFERENCES AND SIMILARITIES BETWEEN COMBUSTION AND STREAMER FRONTS

In the introduction, we draw on the similarity between the streamer problem and other problems like combustion, chemical waves, thermal plumes, phase field models, *etc.*, to motivate the development of an effective interface approach. Of these problems, streamer propagation is most closely analogous to combustion, in that the strong nonlinearity of the reaction rates (the combustion rate and the ionization rate) is an important factor in giving rise to front development in flames and streamers, respectively. There are important differences as well, however, and since several interface techniques were originally developed in the context of combustion [11,12,17], we highlight some of the differences and similarities here:

(a) In combustion the reaction rate depends strongly on the temperature, whose outer dynamics is governed by a diffusion equation of the form $\partial_t T = \nabla^2 T$, while for streamers the ionization rate depends strongly on the field $|\mathbf{E}|$, with \mathbf{E} the *gradient* of the potential Φ that obeys the Laplace equation $\nabla^2 \Phi \approx 0$ in the outer region where the total charge density vanishes. This field strength \mathbf{E} varies strongly in the streamer front, since the increased screening resulting from the rising electron density suppresses \mathbf{E} — and hence the ionization rate — to zero. In combustion, on the other hand, the temperature hardly varies throughout the combustion zone.

(b) Combustion fronts are essentially fronts propagating into a metastable state, because the front has to supply the heat that increases the temperature and hence the reaction rate, while streamer front propagation is an example of front propagation into unstable states, where the reaction starts for any nonvanishing electron density.

(c) In a flame front typically the temperature remains high enough that all the reactions proceed to saturation: all the combustible material burns. The temperature difference between the flame front and the background is then essentially determined by conservation (conversion) of energy. In typical streamer fronts, on the other hand, the field \mathbf{E} is suppressed long before saturation effects start to play a role, and hence the ionization level behind the front is set by the internal dynamics of the front rather than by conservation laws (i.e., the gas density).

(d) The electron drift $-\mu_e \mathbf{E}$ has no clear analogue in combustion.

(e) Finally, the relevant asymptotic expansion for streamers is not quite like the “activation energy asymptotics” of combustion [11,12], since we consider here fields strengths that are comparable to the characteristic field scale E_0 of the ionization rate given in Eq. (2.6) before the front, whereas in combustion activation energy asymptotics is often appropriate since the flame temperature remains much smaller than the chemical activation energy. For streamers, an analysis like activation

energy asymptotics is appropriate in the limit of small fields $|\mathbf{E}| \ll E_0$. Of course, in streamers the rapid variation of the field \mathbf{E} in the front region, and hence the rapid suppression of the ionization rate, looks, at first sight, similar to the suppression of the chemical reaction rate with decreasing temperature in flames. However, in flames this is due to the strongly nonlinear dependence of the reaction rate on temperature before the front, (so that a slight suppression of the temperature reduces the reaction rate dramatically), while in streamers in large external fields of order E_0 this is due to the fact that the field itself is reduced significantly behind the streamer front, as a result of screening.

-
- [1] I. Gallimberti, J. Phys. (France) **40**, Colloque C7, 193 (1979).
- [2] A. Bondiou and I. Gallimberti, J. Phys. D.: Appl. Phys. **27** 1252 (1994).
- [3] Y.P. Raizer, *Gas Discharge Physics* (Springer, Berlin, 1991).
- [4] A.N. Lagarkov and I.M. Rutkevich, *Ionization Waves in Electrical Breakdown of Gases* (Springer, New York, 1994)
- [5] L.M. Vasilyak, S.V. Kostyuchenko, N.N. Kudryavtsev and I.V. Filyugin, Physics–Uspekhi **37**, 247 (1994).
- [6] The distinction between the various discharge phenomena seems to vary between authors.
- [7] See, e.g., D. Braun, U. Kuchler and G. Pietsch, J. Phys. D: Appl. Phys. **24**, 564 (1991); W.R. Rutgers, in: *Proc. XXI Int. Conf. on Phenomena in Ionized Gases*, ed.: AG Plasmaphysik, Bochum, Düsseldorf, Essen, Jülich (RUB, Bochum, 1993); U. Kogelschatz, in: *Proc. X Int. Conf. on Gas Discharges and their Applications*, ed. W.T. Williams (Swansea, UK: Univ. College Swansea, 1992), p. 972, vol. 2.
- [8] S.K. Dhali and P.F. Williams, Phys. Rev. A **31**, 1219 (1985) and J. Appl. Phys. **62**, 4696 (1987).
- [9] P.A. Vitello, B.M. Penetrante, and J.N. Bardsley, Phys. Rev. E **49**, 5574 (1994). See also P.A. Vitello, B.M. Penetrante, and J.N. Bardsley, in *Non-Thermal Plasma Techniques for Pollution Control*, ed. B.M. Penetrante and S.E. Schultheis (Springer, Heidelberg, 1993).
- [10] M C. Wang and E.E. Kunhardt, Phys. Rev. A **42**, 2366 (1990). See references in this article to earlier numerical studies by Kunhardt and co-workers.
- [11] F.A. Williams, *Combustion Theory*, (Benjamin/Cummings, Menlo Park, 1985).
- [12] J.D. Buckmaster and G.S.S. Ludford, *Theory of Laminar Flames*, (Cambridge University Press, Cambridge, 1982), or: J.D. Buckmaster and G.S.S. Ludford, *Lectures on Mathematical Combustion*, (SIAM, Philadelphia, 1983); a nice elementary account of several of the essential ingredients of flame fronts and their instabilities can be found in J.D. Buckmaster, Physica **12D**, 173 (1984).
- [13] M. van Dyke, *Perturbation Methods in Fluid Dynamics* (Parabolic, Stanford, 1975).
- [14] C.M. Bender and S.A. Orszag, *Advanced Mathematical Methods for Scientists and Engineers* (McGraw-Hill, New York, 1978).
- [15] E. Meron, Phys. Rep. **218**, 1 (1992).
- [16] G. Zocchi, P. Tabeling and M. Ben-Amar, Phys. Rev. Lett. **69**, 601 (1992), and M. Ben-Amar, Phys. Fluids **A4**, 2641 (1992).
- [17] P. Fife, *Dynamics of Internal Layers and Diffusive Interfaces* (SIAM, Philadelphia, 1988).
- [18] A. Karma and W.J. Rappel, Phys. Rev. E **53**, R3107 (1996) and references therein.
- [19] U. Ebert, W. van Saarloos, and C. Caroli [submitted].
- [20] Some important reviews of recent advances are: J.S. Langer in *Chance and Matter*, ed. J. Souletie (North-Holland, Amsterdam, 1987); D.A. Kessler, J. Koplik and H. Levine, Adv. Physics **37**, 255 (1988); P. Pelcé, *Dynamics of Curved Fronts* (Academic, Boston, 1988); and Y. Pomeau and M. Ben Amar, in: *Solids far from Equilibrium*, C. Godreche, Ed. (Cambridge University Press, Cambridge, 1992).
- [21] E. Ben-Jacob, H.R. Brandt, G. Dee, L. Kramer, and J.S. Langer, Physica **14D**, 348 (1985).
- [22] W. van Saarloos, Phys. Rev. A **37**, 211 (1988).
- [23] W. van Saarloos, Phys. Rev. A **39**, 6367 (1989).
- [24] W. van Saarloos and P.C. Hohenberg, Physica D **56**, 303 (1992); errata: Physica D **69**, 209 (1993).
- [25] G.C. Paquette and Y. Oono, Phys. Rev. E **49**, 2368 (1994); G.C. Paquette, L.-Y. Chen, N. Goldenfeld and Y. Oono, Phys. Rev. Lett. **72**, 76 (1994).
- [26] D.L. Turcotte, and R.S.B. Ong, J. Plasma Phys. **2**, 145 (1968).
- [27] N.W. Albright, and D.A. Tidman, Phys. Fluids **15**, 86 (1972).
- [28] R.G. Fowler, Advances in Electronics and Electron Physics **35**, 1 (1974); and **41**, 1 (1976).
- [29] M.I. D'yakonov and V.Y. Kachorovskii, Sov. Phys. JETP **68**, 1070 (1989). We have been informed (V. Shumeiko, private communication with WvS), that D'yakonov has in a talk also pointed out the strong similarity between streamers and other problems of pattern formation such as dendrites.
- [30] Instead of distinguishing the two different types of streamers with the acronyms *ADS* (*anode directed streamer*) and *CDS* (*cathode directed streamer*) as is sometimes done, we prefer to use *NSF* (*negative streamer front*) and *PSF* (*positive streamer front*), as this wording is more appropriate for an effective interface approach.
- [31] D.G. Aronson and H.F. Weinberger, in *Partial Differential Equations and Related Topics*, ed.: J.A. Goldstein (Springer, Heidelberg, 1975); Adv. Math. **30**, 33 (1978).
- [32] P. Collet and J.-P. Eckmann, *Instabilities and Fronts in Extended Systems* (Princeton University Press, Princeton, 1990).
- [33] A. Kolmogorov, I. Petrovsky, and N. Piskunov, Bull. Univ. Moskou, Ser. Internat., Sec. A **1**, 1 (1937).
- [34] R.A. Fisher, Ann. Eugenics **7**, 355 (1937).
- [35] A short calculation shows, that a homogeneous current $|\mathbf{j}| = 1$ generates magnetic fields, whose Lorentz force

on the free charges moving with the drift velocity $\mu_e E_0$ becomes comparable to the force of an electric field of order E_0 , when the current extends over a region of width $\ell_B = \alpha_0^{-1} c^2 / (\mu_e E_0)^2 = 1.6 \cdot 10^5 \alpha_0^{-1} \approx 0.4$ m for the parameters of [8,9]. In practice, streamer dimensions are much smaller than ℓ_B , and hence the magnetic fields can generally be neglected.

- [36] S.K. Dhali and A.P. Pal, J. Appl. Phys. **63**, 1355 (1988).
- [37] It is useful to realize that our model is inaccurate for small fields, since any nonzero electron density is unstable in our model for any field strength, due in particular to the neglect of recombination terms. A similar situation occurs in combustion; there, the exponential reaction rate is sometimes modified by introducing a so-called switch-on-temperature, below which the combustion rate is zero. As discussed in [12], the results do not strongly depend on the switch-on-temperature provided it is low enough, and an asymptotic expansion in fact nicely circumvents the introduction of this temperature. Similar observations hold for the streamer problem: A crude way to circumvent the introduction of recombination terms would be to introduce a threshold-field E_{thr} below which $f(|\mathbf{E}|)$ vanishes. Essentially all of our analysis applies to such cases as well, and the results do not depend sensitively on the value of this field.
- [38] One should be aware that it is possible that the state generated by the front is actually unstable or metastable. In such cases the front is usually followed by a second front invading the unstable or metastable state. See, e.g., K. Nozaki and N. Bekki, Phys. Rev. Lett. **51** 2171 (1983), and F.J. Elmer, J. Burns and H. Suhl, Europhys. Lett. **22** 399 (1993).
- [39] The discussion is easily generalized to the case in which the ionization function $f(|\mathbf{E}|) = 0$ in an interval $E \in [0, E_{thr}]$ — compare Ref. [37]. In this case, ionization is absent for any field $E < E_{thr}$, so that there is a larger multiplicity of stationary states of the model, and so that for fields below this threshold the dynamics is due to conventional electrodynamics with conserved particle densities. This generalization, however, does not affect our conclusions about front propagation qualitatively: It is easy to convince oneself that the state generated after long times by a propagating front is still the one with $\mathbf{E} = 0$, since as long as the field behind the front is nonzero but of the same sign as E^+ , charge is pumped into the streamer head from behind in such a way that the field jump across the streamer head increases in time. Since E^+ is fixed, this implies that the field behind the streamer head decays.
- [40] E.D. Lozanskii, Zh. Tekh. Fiz. **38**, 1563 (1968) [Sov. Phys. Tech. Phys. **13**, 1269 (1969)].
- [41] W.K.H. Panofsky and M. Phillips, *Classical Electricity and Magnetism*, (Addison-Wesley, Reading, Massachusetts, 1962), p. 122.
- [42] This is argued in appendix A of [22]: For equations like ours that admit uniformly translating front solutions, one generally expects a *one-parameter* family of front solutions, a feature well known for the case of the nonlinear diffusion equation discussed briefly in Section V. For pattern forming fronts propagating into an unstable state, one typically has a *two parameter* family of fronts — see, e.g., [24] or P. Collet and J.-P. Eckmann, Commun. Math. Phys. **107**, 39 (1986), although a general proof is lacking.
- [43] M. Abramowitz and I.A. Stegun, *Handbook of Mathematical Functions* (Dover, New York, 1972).
- [44] W.H. Press, B.R. Flannery, S.A. Teukolsky and W.T. Vetterling, *Numerical Recipes* (Cambridge University Press, Cambridge, 1986).
- [45] Of course, in principle in a three-dimensional dynamical system, one can have limit cycles and other (strange) attractors. Limit cycles would in our case correspond to spatially periodic ionization waves, but the occurrence of such solutions is clearly excluded physically by the fact the ion density must be a monotonically increasing function of ξ . Other types of nontrivial behavior of the flow are excluded for the same reason.
- [46] See, e.g., R.D. Benguria and M.C. Depassier, Phys. Rev. Lett. **73**, 2272 (1994); and A. Goriely, Phys. Rev. Lett. **75**, 2047 (1995).
- [47] U. Ebert [in preparation].
- [48] I.M. Gel'fand, Usp. Mat. Nauk **14**, No. 2 (86) 87 (1959) [Am. Math. Soc. Transl. Ser. 2 **29** 295 (1963)].
- [49] G.T. Dee and J.S. Langer, Phys. Rev. Lett. **50**, 383 (1983).
- [50] G.T. Dee and W. van Saarloos, Phys. Rev. Lett. **60**, 264 (1988).
- [51] For a review, see A. Bers, in: *Handbook of Plasma Physics*, ed.: M.N. Rosenbluth and R.Z. Sagdeev (North-Holland, Amsterdam, 1983), Vol. 1.
- [52] E.M. Lifshitz and L.P. Pitaevskii, *Physical Kinetics*, Course of Theoretical Physics Vol. 10 (Pergamon, New York, 1981), Chap. VI.
- [53] By tracking the position $x_f(\tau)$ of the point where $\sigma(x_f(\tau), \tau) = 0.1\sigma^-$, we followed the convergence of the local velocity $v_f(\tau) = dx(\tau)/d\tau$ towards v^* , and found it to be consistent with the formula $v_f(\tau) = v^* - 1.5D/(\sqrt{f|E^+|\tau}) + \dots$ derived in [23].
- [54] L. Niemeyer, L. Pietronero and H.J. Wiessmann, Phys. Rev. Lett. **52**, 1033 (1984).
- [55] J.C. Maan, B. Willing, and P. Uhlig, Physica B **204**, 311 (1995).




FIG. 1. Results of the numerical simulations of the full three-dimensional streamer equations (2.1)-(2.6) of Vitello *et al.*, reprinted from Fig. 1 and 10 in [9]. (a) Negative streamer propagating downwards towards the anode. Electrodes are planar and located at $z = 0$ and 0.5 cm; the voltage between the electrodes is 25 kV, which in the absence of the streamer amounts to a constant electric field $|\mathcal{E}| = E_0/4$. The system continues sideways sufficiently far to make the lateral boundaries irrelevant. The streamer is assumed to be cylindrically symmetric. The dimensionless diffusion constant is $D = 0.1$. Each line indicates an increase of n_e by a factor 10; densities of $10^3 - 10^{14} \text{ cm}^{-3}$ can be seen (initial background ionization: 1 cm^{-3}). Shape at time 4.75 ns after an initial ionization seed was placed near the upper electrode. (b) Shape at time 5.5 ns. (c) Logarithmic electron n_e and total charge n_s density along the symmetry axis of (b). Solid line: n_e ; dot-dashed: $|n_s|$ for $n_s > 0$; dotted: $|n_s|$ for $n_s < 0$. Note the exponential increase of the densities on μm scale within the front as well as maximum of both densities in the rear part of the front. Courtesy of P.A. Vitello.




FIG. 2. (a) Uniformly translating fronts for $D = 0$ and $v = 2$ shown as flows in the two-dimensional (E, σ) phase space. Out of each point σ^- on the σ axis, there is a *PSF* flowing to the right and a *NSF* to the left. Both reach the same value $|E^+|$ on the horizontal axis, which also is independent of v . Note, that *NSF* have a maximum of σ within the front, while *PSF* have monotonic σ . Note also, that no physical fronts (i.e., with $\sigma \geq 0$ everywhere) reach a value $E^+ < -v = -2$, in agreement with Eq. (4.11). (b) Sketch of a uniformly translating *PSF* and *NSF* for $D \neq 0$ as a flow in three-dimensional (E, σ, q) phase space. The thick curves indicate the trajectories, while the thin ones show their projection into the $\sigma = 0$ and $q = 0$ planes. For fixed v , there is at each point of the σ axis still only one outgoing vector, which can be followed in two antiparallel directions. The E axis is fully attractive and will always be reached.




FIG. 3. Phase diagram for *PSF* as a function of D and E^+ . Above the solid line the lowest speed of physical front solutions is given by v^* , below the line v^\dagger corresponds to the smallest speed of physical front solutions. Accordingly, the selected front speed is v^* above the solid line (*linear marginal stability regime*), and v^\dagger below the solid line (*nonlinear marginal stability regime*). The dotted curve indicates $v^* = 0$ and is a lower bound for the cross-over to v^\dagger behaviour of the selected fronts.




FIG. 4. $\tilde{v}^\dagger = v^\dagger/D$ (solid) and $\tilde{v}^* = v^*/D$ (dashed lines) as a function of D for $E^+ = 0.3 - 1.0$ in steps of 0.1, and for $E^+ = 1.0 - 2.0$ in steps of 0.2. \tilde{v}^\dagger depends only weakly on D , i.e., the physical front velocity v^\dagger is approximately proportional to D . At $\tilde{v}^\dagger(E^+, D_{cr}(E^+)) = \tilde{v}^*(E^+, D_{cr}(E^+))$, the selected front crosses over from v^\dagger to v^* ; the v^\dagger solutions disappear. Plotting $D_{cr}(E^+)$ in the (E^+, D) plane yields the solid curve in the phase diagram of Fig. 3, while the zeros of v^* determine the dotted curve in Fig. 3.




FIG. 5. Electron density σ^- behind the planar selected front as a function of the field E^+ before the front for several D ; dotted: v^* fronts; solid: v^\dagger fronts. (a) *NSF*: For v^* fronts, σ^- depends but weakly on D . Results for $D = 0, 1, 3$ are shown. Crosses: Extrapolation of $\sigma^-(E^+)$ for $D = 0.1$ for the curved fronts of the 3D simulations [9]. (b) *PSF* results for $D = 0.05, 0.1, 0.2, 0.3, 0.5, 0.7, 1, 2, 3$. For v^\dagger fronts, $\tilde{\sigma}^- = \text{const.} + \mathcal{O}(D)$, i.e., $\sigma^-(E^+)$ is approximately proportional to $1/D$.




FIG. 6. Width w of the front profile (measured between points with 0.1 σ^- and 0.9 σ^-) as a function of E^+ for the selected *NSF* fronts with $D = 0.1$. The dashed line is given by $w = 6/\Lambda_+^-(v^*)$.




FIG. 7. Numerical integration of the time evolution given by Eqs. (3.10) and (3.11) for $D = 1.0$ in a constant background field $E = -1$ (numerical gridsize $\Delta x = 0.1$ and timestep $\Delta\tau = 0.05$, initial perturbation at $x_0 = 50$). Initial condition at $t = 0$: small charge-neutral, ionized region of Gaussian shape depicted by the lowest line. Each new line corresponds to a time step $\Delta t = 2$ and the upper line to $t = 130$. (a) The electron density $\sigma(x, t)$ initially grows and then, after field screening in the middle sets in, develops into a *NSF* propagating to the right and a *PSF* propagating to the left. (b) The electric field $E(x, t)$ stays $E = -1$ in the non-ionized region and becomes dynamically screened to zero in the ionized region.




FIG. 8. Emergence of the uniformly translating *PSF* on the left in the system of Fig. 7. Conditions identical to Fig. 7 except for $x_0 = 150$ and different numerical gridsize ($\Delta x = 0.05$ and $\Delta\tau = 0.01$). $\sigma(x, t)$ is shown in time range $t = 0 - 500$ in time steps $\Delta t = 10$. Note the difference in the duration of the transient regimes, in the propagation velocities of *PSF* and *NSF*, and in the degrees of ionization behind these.




FIG. 9. Identical with Fig. 7(a), except that here $D = 0.1$. Time range also $t = 0 - 130$ in steps of $\Delta t = 2$. The *NSF* has sharper contours and propagates slower than for $D = 1$, the *PSF* appears not to develop.




FIG. 10. Emergence of the uniformly translating *PSF* on the left for $D = 0.1$. Initial conditions identical with Fig. 9. The time range $t = 4000 - 8000$ after an initial perturbation at $t = 0$ and $x_0 = 60$ is shown in time steps of $\Delta t = 100$. (Numerical gridsize $\Delta x = 0.01$ and $\Delta\tau = 0.5$.)




FIG. 11. A non-localized initial condition with $\Lambda = 0.25$ as described in the text; otherwise, the situation is like in Fig. 9, and $D = 0.1$.