Solitonic Gluons

Elena A. Ostrovskaya and Yuri S. Kivshar Optical Sciences Centre, Research School of Physical Sciences and Engineering Australian National University, Canberra ACT 0200, Australia

Zhigang Chen[†] and Mordechai Segev[‡]

Department of Electrical Engineering and Center for Photonics and Optoelectronic Materials Princeton University, Princeton, New Jersey 08544

We describe a physical mechanism for creating multi-soliton bound states where solitons are glued together by attraction between the beams they guide, *solitonic gluons*. We verify the concept of the solitonic gluons experimentally, observing a suppression of the repulsion between dark solitons due to an attractive force between bright guided beams.

It has been established for many models of nonlinear physics that solitons behave as effective particles when interacting with external fields or other solitons [1]. In the case of bright solitons described by the nonlinear Schrödinger (NLS) equation, the coherent interaction of solitons depends on the relative phase between them, so that identical solitons with opposite phases repel each other, whereas in-phase solitons attract each other [2]. Interaction of dark solitons is unconditionally repulsive, in all types of models described by the generalized NLS equation [3].

If a nonlinear system supports propagation of two, or more, waves of different frequencies or polarization, vector solitons consisting of more than one components can be formed. Indeed, when two vector solitons are closely separated, they may form a bound state if the sum of all forces acting between different soliton components is zero. As an example, let us consider interaction of two vector solitons consisting of dark and bright components in a defocusing optical medium. Two dark solitons always repel each other, and as a result, they can not form a bound state [3]. However, if we introduce out-of-phase bright components guided by each of the dark solitons, their attractive interaction creates a proper balance of forces, which results in a stationary two-soliton bound state (see Fig. 1).

The mechanism leading to the formation of bound states of the compound solitons described above, makes it somewhat tempting to draw an analogy between the soliton interaction and the fundamental theory of quarks and gluons – quantum chromodynamics. As is well known [4], gluons mediate the strong force and are responsible for binding the quarks into protons and neutrons, for example. Unless forming a bound state with quarks, gluons do not exist as separate particles. Taking these properties of gluons into consideration, we believe that it would be plausible to call bright components guided by dark solitons in a defocusing medium 'solitonic gluons'. On the one hand, they do not 'live' in a defocusing medium by themselves, i.e. they diffract when separated from the dark solitons; and on the other hand, they 'glue' the dark solitons together in a bound state which otherwise can not be formed.





To demonstrate this general concept on a particular example and verify it later experimentally, we consider two incoherently interacting linearly polarized beams in a photorefractive medium. The corresponding physical model is described by the equations for the normalized envelopes U and W [5]:

$$i\frac{\partial U}{\partial z} + \frac{1}{2}\frac{\partial^2 U}{\partial x^2} + \frac{\beta(1+\rho)U}{1+|U|^2+|W|^2} = 0,$$

$$i\frac{\partial W}{\partial z} + \frac{1}{2}\frac{\partial^2 W}{\partial x^2} + \frac{\beta(1+\rho)W}{1+|U|^2+|W|^2} = 0.$$
(1)

Here $\rho = I_{\infty}/I_d$, $\beta = (k_0 x_0)^2 n_e^4 r_{33} E_0/2$, where I_{∞} stands for the total power density in the limit $x \to \infty$, I_d is the so-called dark irradiance, k_0 is the propagation constant, x_0 is the spatial width of the beam, and $n_e^4 r_{33} E_0$ is a correction to the refractive index due to the external field applied to a crystal along the x-axis [5].

Solitonic solutions of Eqs. (1) can be sought for in the form $U = \sqrt{s} u \exp[i\sigma(1-s)z]$ and $W = \sqrt{s} w \exp[i\sigma(1-s\lambda)z]$, where s < 1 and $\lambda < 1$ are two dimensionless parameters and $\sigma \equiv \operatorname{sgn}\beta$ defines the type of nonlinearity, namely defocusing ($\sigma = +1$) or focusing ($\sigma = -1$). Amplitudes u and w satisfy the normalized equations:

$$i\frac{\partial u}{\partial z} + \frac{1}{2}\frac{\partial^2 u}{\partial x^2} - \frac{\sigma(|u|^2 + |w|^2)u}{1 + s(|u|^2 + |w|^2)} + u = 0,$$

$$i\frac{\partial w}{\partial z} + \frac{1}{2}\frac{\partial^2 w}{\partial x^2} - \frac{\sigma(|u|^2 + |w|^2)w}{1 + s(|u|^2 + |w|^2)} + \lambda w = 0,$$
(2)

where the spatial coordinates z and x are measured in the units of $s/|\beta|(1+\rho)$ and $\{s/[|\beta|(1+\rho)]\}^{1/2}$, respectively.

In Eq. (2), the parameter s characterizes nonlinearity saturation.

First, assuming $\sigma = +1$, we look for stationary, zindependent solutions of the system (2) in the form of dark-bright solitons, where u(x) has nonvanishing asymptotics but w(x) vanishes for $|x| \to \infty$. Then, a compound dark-bright solitary wave (u, w), emerges from the one-component dark solution (u, 0) at a certain bifurcation point $\lambda = \lambda_0(s)$. Near the bifurcation point, we can treat the dark component as an effective waveguide which guides the spatially localized mode w. Such compound solitons form a continuous family in λ , for a fixed s. The existence of this type of vector dark-bright solitons in the photorefractive media has been earlier established theoretically [5] and verified experimentally [6]. The families of two-component dark-bright solitons have been numerically found in Ref. [7].

Interaction between two weakly overlapping darkbright solitons can be described by the effective interaction energy as a function of the soliton separation $x_0, V_{\text{eff}}(x_0) = V_d(x_0) + a^2 V_b(x_0) \cos \phi$, where the first and second terms stand for the interaction between dark and bright soliton components, respectively, with ϕ being a relative phase between the bright components of the equal amplitude a. In the case of small s, the effective interaction energies $V_d(x_0)$ and $V_d(x_0)$ can be found analytically [7], $V_d(x_0) \approx 2 \exp(-4x_0)$ and $V_b(x_0) \approx$ $4x_0 \exp(-2x_0)$. Thus, interaction between two darkbright solitons is crucially affected by the forces acting between the bright components guided by the closely separated dark solitons. If the bright components are out of phase ($\phi = \pi$), an attractive force acting between them can balance (or even overbalance) the repulsion between the dark solitons, thus creating a bound state. This observation has been confirmed for Eqs. (2) by numerically identifying the families of two-soliton bound states [7].

To demonstrate the concept of force balance, we have investigated numerically the interaction of a pair of dark solitons generated by a localized input in the shape of a box-like notch on a cw background. Figures 2(a) and 2(b) show the results of our simulations for two distinct cases. First, the box-like input is launched without a bright component. It generates a pair of scalar (one-component) dark solitons which move away from eachother as they propagate [see Fig. 2(a)] due to their mutual repulsion [3]. However, the propagation dynamics is drastically modified by introducing an additional bright beam. Figure 2(b) shows the results for the input field as in Fig. 2(a) but with the added w-component in the form of the first (one-node) mode of the symmetric box-shaped waveguide. Such an amplitude profile of the bright component has been chosen to mimic a solitonic gluon binding two closely separated dark solitons. As is seen from Fig. 2(b), the attraction between out-of-phase bright guided beams compensates for the repulsive force between dark solitons, so that two dark-bright solitons propagate almost in parallel.



FIG. 2. Generation of a pair of dark solitons from a localized box-like input: (a) no bright component introduced, (b) in the presence of the bright component. Bottom row: corresponding input beam profiles.



FIG. 3. Bound states of bright solitons in the model (2) at $\sigma = -1$. (a),(b) Examples of compound solitons with the first and second guided modes, respectively. (c),(d) Interaction of two bright solitons from (a), without and with the gluon component, respectively.

The concept of solitonic gluons can be readily extended to describe multi-soliton bound states composed of various types of bright vector solitons [8,9]. Bright solitons of Eqs. (2) are possible for a self-focusing nonlinearity, $\sigma = -1$. Continuous in λ , families of such solutions originate at the points of bifurcation of the scalar (u-component only) bright soliton families. Similar to the case of dark-bright solitons, the w- field near the bifurcation points can be described as a linear mode of an effective waveguide induced by the u-soliton, the latter can support more than one mode, depending on the value of s.

A rich variety of the soliton structures, consisting of a fundamental bright soliton and a n-node mode of the soliton-induced waveguide, results in complex shapes of solitonic gluons of the focusing nonlinearity. Interaction between the bright components depends on their relative phase, and therefore the bound states of bright vector solitons should be constructed from out-of-phase (mutually repelling) and in-phase (mutually attracting) components. Some vector solitary waves of this type have already been described [9]. In Figs. 3(a, b) we present two examples of such soliton bound states, found numerically for Eqs. (2), where the repulsion of the out-of-phase fundamental (no nodes) solitons of the u- component is balanced by an attractive force acting between two types of solitonic gluons, formed by the in-phase first or second modes of effective soliton-induced waveguides.

Numerical simulations of the beam propagation, with the input of the type presented in Fig. 3(a), revealed a drastic change in the soliton interaction as shown in Figs. 3(c,d) [cf. Fig. 2(a,b)]. The binding force of the solitonic gluons makes the otherwise mutually repelling out-of-phase solitons propagate in parallel [Fig. 3(d)].

To verify the concept of solitonic gluons experimentally, we have studied the interaction of dark solitons in a photorefractive medium, using the experimental setup for generating dark-bright soliton pairs earlier reported in Refs. [6,10]. A cw argon-ion laser beam (488 nm) is collimated and split by a polarizing beam splitter. The ordinary polarized beam is used as uniform background illumination to mimic the dark irradiance, and the extraordinary polarized beam is split into two soliton-forming beams. One of these two beams is used to generate a dark notch on reflection, by illuminating a wire. The dark notch is then imaged onto the input face of a SBN:61 crystal with the background beam covering the entire input face. One half of the other beam (the bright component) goes through a tilted thin glass that introduces a π phase jump at its centre, thereby creating a field profile resembling the first mode of a box-like waveguide, i.e. a bright beam with a node [Fig. 4(a)]. We make the dark and bright input beams mutually incoherent by having their optical path difference greatly exceed the coherence length of the laser. 53 µm



FIG. 4. Photographs showing the effect of the solitonic gluons via coupling of a bright (lower row) and dark (upper row) incoherently interacting beams: (a) an input at z = 0 mm, and three outputs at z = 11.7 mm for (b) normal diffraction of coupled beams, (c) stationary two-soliton bound state, (d) decoupled beam propagation.

Figures 4(a-d) show photographs of the bright and dark beams taken on the input [Fig. 4(a)] and output [Figs. 4(b,c,d)] faces of a 11.7-mm-long SBN crystal. Without a dc field, both beams merely diffract in a similar fashion, as shown in Fig. 4(b). When we launch only a dark-notch-bearing beam in the presence of an appropriate dc field in the x direction (||c| axis), it generates a Y-junction created by a pair of repelling dark solitons, as is seen in Fig. 4(d) (upper row) and earlier reported in Ref. [10]. On the other hand, the uncoupled bright beam diffracts even more strongly due to effectively defocusing nonlinearity, Fig. 4(d) (lower row).

Repulsion of a pair of dark solitons created by a wire is suppressed dramatically when we simultaneously launch a bright beam in the form of a second waveguide mode, thus creating two bright beams with opposite phases. As can be seen from Figs. 4(c) and 4(d), the output distance between the dark solitons is subsequently reduced from $53\mu m$ to $43\mu m$. In addition, we have verified that when the repulsion between the dark solitons is enhanced when the bright beams are in-phase.

In conclusion, we have suggested and verified experimentally a concept of solitonic gluons featuring a balance of attractive and repulsive forces acting between different components of compound solitons as a physical mechanism responsible for the formation of bound states of vector solitons. This novel type of interaction of compound solitons is a rather fundamental phenomenon which should also be observed for many other multicomponent nonlinear models, e.g. incoherent solitons.

The work has been supported by the Australian Photonics Cooperative Research Centre. The authors are indebted to D. Christodoulides, M. Mitchell, and A. Sheppard for useful discussions.

† Present address: Department of Physics and Astronomy, San Francisco State University, San Francisco, CA 94132, USA

‡ Present address: Physics Department, Technion-Israel Institute of Technology, Haifa, 3200, Israel

- See, e.g., Yu. S. Kivshar and B. A. Malomed, Rev. Mod. Phys. **61**, 763 (1989); M. Segev and G. I. Stegeman, Physics Today (1998) August.
- [2] See, e.g., C. Decem and P. L. Chu, In: Optical Solitons-Theory and Experiment, Ed. J. R. Taylor (Cambridge University Press, Cambridge, 1992), p. 107.
- [3] Yu. S. Kivshar and B. Luther-Davies, Phys. Rep. 298, 81 (1998), and references therein.
- [4] See, e.g., F. J. Ynduráin, "Quantum Chromodynamics: An Introduction to the Theory of Quarks and Gluons" (Springer-Verlag, New York, 1983).
- [5] D. N. Christodoulides, S. R. Singh, M. I. Carvalho, and M. Segev, Appl. Phys. Lett. 68, 1763 (1996); see also M. I. Carvalho, S. R. Singh, D. N. Christodoulides, and R. I. Joseph, Phys. Rev. E 53, R53 (1996).
- [6] Z. Chen, M. Segev, T. H. Coskun, D. N. Cristodoulides, Yu. S. Kivshar, and V. V. Afanasjev, Opt. Lett. **21**, 1821 (1996); J. Opt. Soc. Am. B **14**, 3066 (1997).
- [7] E. A. Ostrovskaya and Yu. S. Kivshar, Opt. Lett. 23, 1268 (1998).
- [8] See, e.g., M. Haelterman and A. P. Sheppard, Phys. Rev. E 49, 3376 (1994); A. W. Snyder, D. J. Mitchell, and M. Haelterman, Opt. Commun. 116, 365 (1995).
- [9] J. Yang, Stud. Appl. Math. 100, 127 (1997).
- [10] Z. Chen, M. Mitchell, and M. Segev, Opt. Lett. 21, 716 (1996).