COMMENT ON "DO ZERO-ENERGY SOLUIIONS OF MAXWELL EQUATIONS HAVE THE PHYSICAL ORIGIN SUGGESTED BY A. E. CHUBYKALO?" BY V.V. DVOEGLAZOV

## A. E. Chubykalo

Escuela de Física, Universidad Autónoma de Zacatecas Apartado Postal C-580 Zacatecas 98068, ZAC., México

Received October 6, 1999

Key words: electromagnetic field energy.

Dvoeglazov's paper is not trivial because we know the classical definition of any non-trivial truth: a truth is not trivial *if and only if* an *opposite* statement is not *false* (and vice versa, of course). So I strongly recommend his paper for a reader.

But I have a few notes:

1) V. Dvoeglazov writes in his footnote number 7: "In my opinion, equations (20,21) of ref. [1] are just another form of the Maxwell equations for this particular case, in the sense that there is no new physical content if one expects that the Maxwell electrodynamics describes also the Coulomb interaction."

The point is that precisely this ("another") form of the Maxwell equations (namely with total derivatives!) allows us to describe the Coulomb interaction as an action-at-a-distance (it was shown in [2]). It is well-known that generally accepted Maxwell equations (with partial derivatives only) describe just transverse<sup>1</sup> electromagnetic waves in vacuum which spread with a limited velocity

<sup>&</sup>quot;transverse" in this case means that vectors  ${\bf E}$  and  ${\bf B}$  are perpendicular with respect to wave vector  ${\bf k}$  in every point.

2 A. E. Chubykalo

(i.e. they describe so called *short-range* interaction only). That is why there is *new physical content*<sup>2</sup> in Eqs. (18-21) [1]. Note also that V. Dvoeglazov here erroneously mentioned just Eqs. (20,21). The point is that Eqs. (20,21) describe exclusively instantaneous (action-at-a-distance) interaction while the whole system of our equations (18-21) (Eqs. (1-4) in this Comment) describe both instantaneous and short-range. These equations are written as two *uncoupled* pairs of differential equations:

$$\nabla \times \mathbf{H}^* = \frac{1}{c} \frac{\partial \mathbf{E}^*}{\partial t},\tag{1}$$

$$\nabla \times \mathbf{E}^* = -\frac{1}{c} \frac{\partial \mathbf{B}^*}{\partial t} \tag{2}$$

and

$$\nabla \times \mathbf{H}_0 = \frac{4\pi}{c} \mathbf{j} - \frac{1}{c} \left( \sum_i \mathbf{V}_i \cdot \nabla \right) \mathbf{E}_0, \tag{3}$$

$$\nabla \times \mathbf{E}_0 = \frac{1}{c} \left( \sum_i \mathbf{V}_i \cdot \nabla \right) \mathbf{H}_0. \tag{4}$$

This system follows from:

$$\nabla \times \mathbf{H} = \frac{4\pi}{c} \mathbf{j} + \frac{1}{c} \frac{d\mathbf{E}}{dt}$$
 (5)

$$\nabla \times \mathbf{E} = -\frac{1}{c} \frac{d\mathbf{H}}{dt} \tag{6}$$

where

$$\mathbf{E} = \mathbf{E}_0 + \mathbf{E}^* = \mathbf{E}_0(\mathbf{r} - \mathbf{r}_q(t)) + \mathbf{E}^*(\mathbf{r}, t), \tag{7}$$

$$\mathbf{H} = \mathbf{H}_0 + \mathbf{H}^* = \mathbf{H}_0(\mathbf{r} - \mathbf{r}_q(t)) + \mathbf{H}^*(\mathbf{r}, t), \tag{8}$$

 $\mathbf{r}$  is the fixed point of observation,  $\mathbf{r}_q$  is the point of the location of a moving charge q at the instant t, the total time derivative of any vector field value  $\mathbf{E}$  (or  $\mathbf{H}$ ) can be calculated by the following rule:

$$\frac{d\mathbf{E}}{dt} = \frac{\partial \mathbf{E}^*}{\partial t} - \left(\sum_i \mathbf{V}_i \cdot \nabla\right) \mathbf{E}_0, \tag{9}$$

<sup>&</sup>lt;sup>2</sup>I should to note here that a considerable number of papers have recently been published which declare (prove?) an existence of so called longitudinal electromagnetic waves in vacuum but these ideas still are not generally accepted in classical electrodynamics (see, e.g., review [3]).

here  $\mathbf{V}_i$  are velocities of the particles at the same moment of time of observation.<sup>3</sup>

2) V. Dvoeglazov's phrase: "The main problem with the Chuby-kalo derivation is the following: the integrals are *divergent* when they extend over all the space. ... But, to the best of my knowledge, some persons claimed that such procedures do not have sound mathematical foundation."

I do not see any *mathematical* prohibition to use such procedures. See, e.g., [4] (Vol. 2, ch. 17, §1, point 282). Fihtengoltz here considers the following *definition* of the integral with infinite limits:

$$\int_{a}^{\infty} f(x)dx = \lim_{A \to \infty} \int_{a}^{A} f(x)dx.$$
 (10)

If this limit  $(rhs\ 10)$  does not exist or(attention!) is infinite, they say that this integral diverges. But then Fihtengoltz adds, however, that in the case when the limit  $(rhs\ 10)$  exists and it is equal to infinity one can consider the infinite limit  $(rhs\ 10)$  as a value of the integral  $(lhs\ 10)$ .

V. Dvoeglazov asks: "Furthermore, even if one accepts its validity the total energy resulting from integration of (28) over the whole space is to be infinite!?"

Why not? Many years ago, in Newton's times, for example, nobody (I mean, physicists) doubted that the Universe had an infinite number of stars. It meant that the mass of the Universe was infinite. And this mass is no more than  $\int\limits_{\infty} \varrho d\mathcal{V}$  over the infinite volume, where  $\varrho$  is a limited mass density of the Universe. Now many physicists believe that the mass (and, of course, the total energy) of our Universe is limited but their statements have no any evidence (however the statement that the Universe has an infinite mass also has no evidence). Unfortunately, it is still a matter of the belief.

At last V. Dvoeglazov cites: "It was noted by Barut [5, p.105] that in the case of non-vanishing fields at the spatial infinity 'we cannot expect to find globally conserved quantities'."

I have only one question here: Is it possible *experimentally* to find *globally* conserved quantities?

3) The final phrase of V. Dvoeglazov's paper is: "... we are not yet convinced in the necessity of correction of the formula for the

 $^4$ Recall that Eq. (28) in [1] is:

$$w = \frac{2\mathbf{E}^* \cdot \mathbf{E}_0 + 2\mathbf{H}^* \cdot \mathbf{H}_0 + E_0^2 + H_0^2}{8\pi}.$$

<sup>&</sup>lt;sup>3</sup>Note (see [2]) that unlike the fields  $\{\}^*$  the fields  $\{\}_0$  do not retard.

4 A. E. Chubykalo

energy density for radiation field because of the absence of firm experimental and mathematical bases in [1]."

In turn, I explained in this short comment that if we *suppose* that the radiation field *exists* in infinity (*mathematically* it is possible, see above, on the other hand the opposite statement has no evidences) we must correct the formula of the energy density for the electromagnetic field.

## REFERENCES

- 1. A. E. Chubykalo, Mod. Phys. Lett. A13, 2139 (1998).
- A. Chubykalo and R. Smirnov-Rueda, Phys Rev. E53, 5373 (1996); ibid. E55, 3793E (1997); Mod. Phys. Lett. A12, 1 (1997).
- 3. V. V. Dvoeglazov, *Hadronic J. Suppl.* **12**, 241 (1997).
- 4. G. M. Fihtengoltz, "Osnovy Matematicheskogo Analiza [Foundations of Mathematical Analysis]" (GI TTL, Moscow, 1956) (in Russian).
- 5. A. O. Barut, "Electrodynamics and classical theory of fields and particles" (Dover Publications, Inc., New York, 1979).