

# Finite nuclear size correction to the bound-electron $g$ factor in a hydrogenlike atom

D. A. Glazov and V. M. Shabaev

*Department of Physics, St.Petersburg State University, Oulianovskaya 1, Petrodvorets, St.Petersburg 198504, Russia*

The finite nuclear size correction to the bound-electron  $g$  factor in hydrogenlike atoms is investigated in the range  $Z=1-20$ . An analytical formula for this correction which includes the non-relativistic and dominant relativistic contributions is derived. In the case of the  $1s$  state, the results obtained by this formula are compared with previous non-relativistic analytical and relativistic numerical calculations.

PACS number(s): 31.30. Jv, 31.30. Gs

## I. INTRODUCTION

Recent experiments on measuring the bound-electron  $g$  factor in hydrogenlike carbon ( $C^{5+}$ ) reached an accuracy of about  $2 \cdot 10^{-9}$  [1,2]. The same accuracy is expected to be achieved soon for some other low- $Z$  ions. To obtain the corresponding precision in the theoretical predictions for the bound-electron  $g$  factor, the relativistic, QED, nuclear recoil and nuclear size corrections must be evaluated (see [3–6] and references therein). In the present paper we derive a relativistic formula for the finite nuclear size correction to the bound-electron  $g$  factor in the case of an arbitrary state of a hydrogenlike atom. This formula provides a sufficiently accurate evaluation of the correction under consideration in the range  $Z = 1-20$ . In the case of the  $1s$  state, the results obtained by this formula are compared with the results obtained by the related non-relativistic formula [7] and with the results of the relativistic numerical evaluation [4].

The relativistic units ( $\hbar = c = 1$ ) and the Heaviside charge unit ( $\alpha = \frac{e^2}{4\pi}, e < 0$ ) are used in the paper.

## II. BASIC FORMULAS

We consider a hydrogenlike atom, placed in a homogeneous magnetic field,  $\vec{A}(\vec{r}) = [\vec{B} \times \vec{r}]/2$ . We assume that the interaction of the electron with the magnetic field is much smaller than the fine structure splitting and much larger than the hyperfine structure splitting, if the nucleus has a nonzero spin. The energy shift of a level  $a$  to the first order in the magnetic field is

$$\Delta E_a = \langle \Psi_a | V_B | \Psi_a \rangle, \quad (2.1)$$

where

$$V_B = \frac{|e|}{2} (\vec{B} \cdot [\vec{r} \times \vec{\alpha}]). \quad (2.2)$$

Assuming that the vector  $\vec{B}$  is directed along the  $z$  axis, the energy shift reads

$$\Delta E_a = \mu_0 g B m_j, \quad (2.3)$$

where  $\mu_0 = \frac{|e|}{2m}$  is the Bohr magneton,  $m_j$  is the  $z$ -projection of the angular momentum, and  $g$  is the bound-electron  $g$  factor. In the case of a point-charge nucleus a simple calculation, based on the Dirac equation, yields [8]

$$g_D = \frac{\kappa}{j(j+1)} \left( \kappa \frac{E_{n\kappa}}{m} - \frac{1}{2} \right). \quad (2.4)$$

Here  $\kappa = (-1)^{j+l+\frac{1}{2}}(j + \frac{1}{2})$  is the relativistic angular quantum number,  $j$  is the total angular momentum of the electron,  $l = j \pm 1/2$  defines the parity of the state,  $E_{n\kappa}$  is the energy of the state

$$E_{n\kappa} = \frac{\gamma + n_r}{N} m, \quad (2.5)$$

$n_r = n - |\kappa|$  is the radial quantum number,  $n$  is the principal quantum number,  $\gamma = \sqrt{\kappa^2 - (\alpha Z)^2}$ , and  $N = \sqrt{(n_r + \gamma)^2 + (\alpha Z)^2}$ .

The finite nuclear size induces a deviation of the  $g$  factor from its Dirac value,

$$g = g_D + \Delta g. \quad (2.6)$$

To find the nuclear size correction by perturbation theory, we have to evaluate the expression

$$\Delta E_a = 2 \sum_n^{n \neq a} \frac{\langle \Psi_a | V_F | \Psi_n \rangle \langle \Psi_n | V_B | \Psi_a \rangle}{E_a - E_n}, \quad (2.7)$$

where  $V_F = V - V_C$  defines a deviation of the potential from the pure Coulomb one,  $V_C = -\alpha Z/r$ . After a simple integration over the angular variables, the  $\Delta g$  value reads

$$\Delta g = \frac{2\kappa m}{j(j+1)} \sum_{n'}^{n' \neq n} \frac{\langle n\kappa | V_F | n'\kappa \rangle \langle n'\kappa | r\sigma_x | n\kappa \rangle}{E_{n\kappa} - E_{n'\kappa}}, \quad (2.8)$$

where  $|n\kappa\rangle$  is a two-component vector defined by

$$|n\kappa\rangle = \begin{pmatrix} r g_{n\kappa}(r) \\ r f_{n\kappa}(r) \end{pmatrix}, \quad (2.9)$$

$g_{n\kappa}$  and  $f_{n\kappa}$  are the upper and lower radial components of the Dirac wave function defined as in [9],  $\sigma_x$  is the Pauli matrix acting in the space of the two-component vectors, and the scalar product of the two-component vectors is defined by

$$\langle a | b \rangle = \int_0^\infty dr r^2 (g_a g_b + f_a f_b). \quad (2.10)$$

The sum (2.8) can be evaluated analytically using the method of the generalized virial relations for the Dirac equation proposed in [10]. Equations (3.26)–(3.29) of Ref. [10] can be written as

$$(E_{n'\kappa} - E_{n\kappa}) \langle n'\kappa | r^s | n\kappa \rangle = -s \langle n'\kappa | i\sigma_y r^{s-1} | n\kappa \rangle, \quad (2.11)$$

$$(E_{n'\kappa} - E_{n\kappa}) \langle n'\kappa | i\sigma_y r^s | n\kappa \rangle = 2m \langle n'\kappa | \sigma_x r^s | n\kappa \rangle + s \langle n'\kappa | r^{s-1} | n\kappa \rangle - 2\kappa \langle n'\kappa | \sigma_z r^{s-1} | n\kappa \rangle, \quad (2.12)$$

$$(E_{n'\kappa} + E_{n\kappa}) \langle n'\kappa | \sigma_z r^s | n\kappa \rangle = 2m \langle n'\kappa | r^s | n\kappa \rangle + s \langle n'\kappa | \sigma_x r^{s-1} | n\kappa \rangle - 2\alpha Z \langle n'\kappa | \sigma_z r^{s-1} | n\kappa \rangle, \quad (2.13)$$

$$(E_{n'\kappa} + E_{n\kappa}) \langle n'\kappa | \sigma_x r^s | n\kappa \rangle = -s \langle n'\kappa | \sigma_z r^{s-1} | n\kappa \rangle + 2\kappa \langle n'\kappa | r^{s-1} | n\kappa \rangle - 2\alpha Z \langle n'\kappa | \sigma_x r^{s-1} | n\kappa \rangle, \quad (2.14)$$

where  $\sigma_x, \sigma_y, \sigma_z$  are the Pauli matrices. From these equations we obtain

$$\begin{aligned} \langle n'\kappa | r\sigma_x | n\kappa \rangle &= \left( \kappa \frac{E_{n\kappa}}{m^2} - \frac{1}{2m} \right) \langle n'\kappa | n\kappa \rangle + (E_{n\kappa} - E_{n'\kappa}) \\ &\times \left\langle n'\kappa \left| \frac{\kappa}{m^2} \left[ \left( E_{n\kappa} - \frac{m}{2\kappa} \right) r i\sigma_y + m r \sigma_x + \alpha Z i\sigma_y - \kappa \sigma_z \right] \right| n\kappa \right\rangle. \end{aligned} \quad (2.15)$$

From equation (2.15), taking into account that

$$\langle n'\kappa | n\kappa \rangle = \delta_{n'n} \quad (2.16)$$

and

$$\sum_{n'}^{n' \neq n} |n'\kappa\rangle \langle n'\kappa| = I - |n\kappa\rangle \langle n\kappa|, \quad (2.17)$$

we find

$$\sum_{n'}^{n' \neq n} \frac{|n'\kappa\rangle \langle n'\kappa| r \sigma_x |n\kappa\rangle}{E_{n\kappa} - E_{n'\kappa}} = (P - \langle n\kappa|P|n\kappa\rangle) |n\kappa\rangle, \quad (2.18)$$

where

$$P = \frac{\kappa}{m^2} \left[ \left( E_{n\kappa} - \frac{m}{2\kappa} \right) r i \sigma_y + m r \sigma_x + \alpha Z i \sigma_y - \kappa \sigma_z \right]. \quad (2.19)$$

Finally, we should evaluate the expression

$$\Delta g = \frac{2\kappa m}{j(j+1)} (\langle n\kappa|V_F P|n\kappa\rangle - \langle n\kappa|V_F|n\kappa\rangle \langle n\kappa|P|n\kappa\rangle). \quad (2.20)$$

We assume that the nuclear charge distribution is described by a spherically symmetric density  $\rho(\vec{r}) = \rho(r)$ , which is normalized by the equation

$$\int d\vec{r} \rho(\vec{r}) = 1. \quad (2.21)$$

The Poisson equation gives

$$\Delta V_F(\vec{r}) = 4\pi\alpha Z [\rho(\vec{r}) - \delta(\vec{r})], \quad (2.22)$$

where  $\Delta$  is the Laplace operator. When integrated with  $V_F$ , the radial functions  $g(r)$  and  $f(r)$  can be approximated by the lowest order term of the expansion in powers of  $r$ . It follows we have to evaluate the integral

$$I = \int_0^\infty dr r^2 r^{2\gamma-2} V_F. \quad (2.23)$$

Using the identity

$$r^\beta = \frac{1}{(\beta+2)(\beta+3)} \Delta r^{\beta+2} \quad (2.24)$$

and integrating by parts, we find

$$\begin{aligned} I &= \int_0^\infty dr r^2 \frac{1}{2\gamma(2\gamma+1)} \Delta r^{2\gamma} V_F = \int_0^\infty dr r^2 \frac{1}{2\gamma(2\gamma+1)} r^{2\gamma} \Delta V_F \\ &= \frac{4\pi\alpha Z}{2\gamma(2\gamma+1)} \int_0^\infty dr r^2 r^{2\gamma} \rho(r) = \frac{\alpha Z}{2\gamma(2\gamma+1)} \langle r^{2\gamma} \rangle, \end{aligned} \quad (2.25)$$

where

$$\langle r^{2\gamma} \rangle = \int d\vec{r} r^{2\gamma} \rho(r). \quad (2.26)$$

For the correction to the  $g$  factor we obtain

$$\begin{aligned} \Delta g &= \frac{\kappa^2}{j(j+1)} \cdot \frac{\Gamma(2\gamma+1+n_r) 2^{2\gamma-1}}{\gamma(2\gamma+1) \Gamma^2(2\gamma+1) n_r! (N-\kappa) N^{2\gamma+2}} \\ &\quad \times \left[ \left( n_r^2 + (N-\kappa)^2 \right) \left( 1 - 2\kappa \frac{E_{n\kappa}}{m} \right) - 2n_r(N-\kappa) \left( \frac{E_{n\kappa}}{m} - 2\kappa \right) \right] \\ &\quad \times (\alpha Z)^{2\gamma+2} m^{2\gamma} \langle r^{2\gamma} \rangle. \end{aligned} \quad (2.27)$$

For  $ns$ -states, which are of particular interest, the expansion of this expression to two lowest orders in  $\alpha Z$  yields

$$\Delta g = \frac{8}{3n^3}(\alpha Z)^4 m^2 \langle r^2 \rangle \left[ 1 + (\alpha Z)^2 \left( \frac{1}{4} + \frac{12n^2 - n - 9}{4n^2(n+1)} + 2\Psi(3) - \Psi(2+n) - \frac{\langle r^2 \log(2\alpha Zmr/n) \rangle}{\langle r^2 \rangle} \right) \right], \quad (2.28)$$

where  $\Psi(x) = \frac{d}{dx} \log \Gamma(x)$ . For the  $1s$  state, we have

$$\Delta g = \frac{8}{3}(\alpha Z)^4 m^2 \langle r^2 \rangle \left[ 1 + (\alpha Z)^2 \left( 2 - C - \frac{\langle r^2 \log(2\alpha Zmr) \rangle}{\langle r^2 \rangle} \right) \right], \quad (2.29)$$

where  $C=0.57721566490$  is the Euler constant. In the non-relativistic limit, we find

$$\Delta g = \frac{8}{3n^3}(\alpha Z)^4 m^2 \langle r^2 \rangle \quad (2.30)$$

for  $ns$  states and

$$\Delta g = \frac{2(n^2 - 1)}{3n^5}(\alpha Z)^6 m^2 \langle r^2 \rangle \quad (2.31)$$

for  $np_{\frac{1}{2}}$  states. In the case of the  $1s$  state, the expression (2.30) coincides with the related formula in [7].

### III. NUMERICAL RESULTS

In Table 1 we compare the  $\Delta g$  values for the  $1s$  state obtained by formula (2.29) with the non-relativistic results of Ref. [7] (it corresponds to equation (2.30) of the present paper) and with the relativistic numerical results of Ref. [4]. To calculate  $\langle r^2 \log r \rangle$  in equation (2.29), we considered the homogeneously charged sphere model for the nuclear charge distribution. As one can see from the table, the relativistic contribution to  $\Delta g$  becomes comparable with the current experimental accuracy for ions with  $Z \geq 12$ . It will be also important for lower  $Z$  ions, provided the experimental accuracy is improved by an order of magnitude.

### ACKNOWLEDGEMENTS

Valuable conversations with T. Beier, S. Karshenboim, J. Kluge, W. Quint, and V. Yerokhin are gratefully acknowledged. This work was supported in part by RFBR (Grant N. 01-02-17248) and by the program "Russian Universities - Basic Research" (project No. 3930).

- [1] N. Hermanspahn, H. Häffner, H.-J. Kluge, W. Quint, S. Stahl, J. Verdú and G. Werth, Phys. Rev. Lett. 84 (2000) 427.
- [2] H. Häffner, Th. Beier, N. Hermanspahn, H.-J. Kluge, W. Quint, S. Stahl, J. Verdú and G. Werth, Phys. Rev. Lett. 85 (2000) 5308.
- [3] T. Beier, I. Lindgren, H. Persson, S. Salomonson, P. Sunnergren, H. Häffner, N. Hermanspahn, Phys. Rev. A 62 (2000) 032510.
- [4] T. Beier, Phys. Rep. 339 (2000) 79.
- [5] A. Czarnecki, K. Melnikov, and A. Yelkhovsky, Phys. Rev. A 63 (2001) 012509.
- [6] S. G. Karshenboim, arXiv:hep-ph/0008227 (2000).
- [7] S. G. Karshenboim, Phys. Lett. A 266 (2000) 380.
- [8] S. A. Zapryagaev, Opt. Spectrosc. 47 (1979) 9.
- [9] A. I. Akhiezer and V. B. Berestetsky, Quantum Electrodynamics (Moscow, Nauka, 1969).
- [10] V. M. Shabaev, J. Phys. B 24 (1991) 4479.

TABLE I. The finite nuclear size correction  $\Delta g$  for the  $1s$  state.

$Z$	$\langle r^2 \rangle^{1/2}$ fm	Ref. [7] (= Eq. (2.30)) [ $10^{-9}$ ]	Eq. (2.29) [ $10^{-9}$ ]	Ref. [4] [ $10^{-9}$ ]
1	0.862	0.00003768	0.00003770	< 0.01
2	1.671	0.0022655	0.0022705	< 0.01
4	2.390	0.074154	0.074741	0.09
6	2.468	0.40031	0.40710	0.42
8	2.693	1.5064	1.5499	1.56
10	3.006	4.5822	4.7810	4.78
12	3.057	9.827	10.426	10.40
14	3.123	19.00	20.54	20.47
16	3.263	35.38	39.05	38.90
18	3.427	62.52	70.53	70.28
20	3.478	98.15	113.4	113.15