

Transparency of Magnetized Plasma at Cyclotron Frequency

G. Shvets

Princeton Plasma Physics Laboratory, Princeton University, Princeton, NJ 08543

J. S. Wurtele

University of California, Berkeley, CA 94720

Electromagnetic radiation is strongly absorbed by the magnetized plasma if its frequency equals the cyclotron frequency of plasma electrons. It is demonstrated that absorption can be completely canceled in the presence of a second radiation beam, or even a magnetostatic field of an undulator, resulting in plasma transparency at the cyclotron frequency. This effect is reminiscent of the electromagnetically-induced transparency (EIT) of the three-level atomic systems, except that it occurs in a completely *classical* plasma. Also, because of the complexity of the classical plasma, index of refraction at cyclotron frequency differs from unity. Potential applications of the EIT in plasma include selective plasma heating, electromagnetic control of the index of refraction, and electron/ion acceleration.

Electromagnetically induced transparency (EIT) in quantum-mechanical atomic systems is a well understood and thoroughly studied [1] subject. EIT is the basis of several very important applications, such as slow light [3], information transfer between matter and light [4,5], sound wave generation [6], or even testing of the black-hole physics [7]. Several recent reviews [2] illucidated the quantum mechanical mechanism of EIT which relies on the destructive interference between several pathways which connect the ground and excited states of the atom. The purpose of this Letter is to describe EIT in a classical plasma.

We consider an externally magnetized plasma with $\vec{B} = B_0 \vec{e}_z$ and density n_0 . A right-hand polarized electromagnetic wave (which we refer to as the probe) at the frequency ω_1 equal to cyclotron frequency $\Omega_0 = eB_0/mc$ cannot propagate in the plasma because it undergoes resonant cyclotron absorption [8]. The cold magnetized plasma dispersion relation ω_1 v. $s.k_1$ for the right-hand polarized probe, plotted in Fig. 1, is given by $\omega_1^2 = k_1^2 c^2 + \frac{\omega_p^2 \omega_1}{\omega_1 - \Omega_0}$. Plasma current and the wavenumber k_1 become infinite for $\omega_1 \rightarrow \Omega_0$, and a forbidden bandgap develops between Ω_0 and $\Omega_c = \sqrt{\Omega_0^2/4 + \omega_p^2} + \Omega_0/2$, where $\omega_p = (4\pi e^2 n_0/m)^{1/2}$ is the plasma frequency. This Letter demonstrates that by adding a second intense electromagnetic wave (pump) with frequency $\omega_0 = \Omega_0 - \omega_p$ create a transparency near the cyclotron frequency. Moreover, if $\omega_p = \Omega_c$, transparency can be created by a magnetostatic undulator with arbitrary wavenumber k_0 .

The classical mechanism of the electromagnetically induced transparency is the destructive interference between the electric field of the probe $\vec{E}_{1\perp}$ and the sidebands of the electric $\vec{E}_{0\perp}$ and magnetic $\vec{B}_{0\perp}$ fields of the pump which are produced by the collective electron plasma oscillation with frequency ω_p along the magnetic field. Qualitatively, the total force at the cyclotron fre-

quency experienced by a plasma electron is given by $\vec{F}_{\text{tot}} \approx -e(\vec{E}_{1\perp} + \zeta_z \partial_z \vec{E}_{0\perp} + \zeta_z \vec{e}_z \times \vec{B}_{0\perp})$, where ζ_z is the electron displacement in the plasma wave. If the pump, probe, and plasma waves are properly phased, then Therefore, if the amplitudes and phases of the pump and the plasma wave are properly correlated, then $\vec{F}_{\text{tot}} = 0$. Consequently, the plasma current at the cyclotron frequency is small (or even vanishing), and the probe propagates as if in vacuum. Our numerical simulation below demonstrates that this correlation is naturally achieved in a collisionless plasma.




FIG. 1. Dispersion curve for a right-hand polarized wave propagating along magnetic field. Forbidden gap exists between cyclotron frequency $\Omega_0 = eB_0/mc$ and cutoff frequency $\Omega_c = \Omega_0/2 + \sqrt{\Omega_0^2/4 + \omega_p^2}$

We assume two right-hand polarized EM waves propagating along z -direction, with their electric and magnetic fields given by $2e\vec{E}_{0\perp}/mc\omega_0 = a_{\text{pump}}\vec{e}_+ \exp(i\theta_0) + c. c.$, $2e\vec{E}_{1\perp}/mc\omega_1 = a_{\text{probe}}\vec{e}_+ \exp(i\theta_1) + c. c.$, and $\vec{B}_{0,1\perp} = (c\vec{k}_{0,1}/\omega_{0,1}) \times \vec{E}_{0,1}$, where $\vec{e}_+ = \vec{e}_x + i\vec{e}_y$, $\vec{e}_- = \vec{e}_x - i\vec{e}_y$, $\theta_0 = k_0 z - \omega_0 t$, and $\theta_1 = k_1 z - \omega_1 t$. Non-relativistic equation of motion of a plasma electron in the combined field is given by

$$\frac{d^2 \vec{x}}{dt^2} + \Omega_0 \vec{v} \times \vec{e}_z + \omega_p^2 \zeta_z \vec{e}_z = -\frac{e}{m} \sum_{m=0,1} \left(\vec{E}_m + \frac{\vec{v} \times \vec{B}_{\perp m}}{c} \right), \quad (1)$$

where $\vec{x} \equiv (z_0 + \zeta_z) \vec{e}_z + \vec{x}_\perp$ and $\vec{v} = d\vec{x}/dt$ are the particle position and velocity, and the initial conditions are $\vec{v} = 0$ and $\vec{x} = z_0 \vec{e}_z$. The third term in the lhs of Eq. (1) is the restoring force of the ions [9].

Equation (1) was integrated for two cases: (a) when only a probe field is turned on, and (b,c) when both the pump and the probe are turned on. The pump and the probe amplitudes were increased adiabatically in time, up to their respective peak amplitudes of a_0 and a_1 , according to

$$\begin{aligned} a_{\text{pump}} &= \frac{a_0}{2} (1 + \tanh[(\Omega_0 t - 160)/40]), \\ a_{\text{probe}} &= \frac{a_1}{2} (1 + \tanh[(\Omega_0 t - 320)/40]), \end{aligned} \quad (2)$$

enabling the pump to turn on first, followed by the probe.

Simulation results for $\omega_p/\Omega_0 = 0.3$ ($\omega_0 = 0.7\Omega_0$) are shown in Fig. 2. Without the pump electron is resonantly driven by the probe as shown in Fig. 2(a). In the plasma, this growth manifests itself in a large electron current and probe absorption because, time-averaged, $\vec{E}_\perp \cdot \vec{v}_\perp < 0$. Adding a strong pump with $a_0 = 0.1$ and $k_0 \approx 0.83\Omega_0/c$ dramatically changes electron motion, see Fig. 2(b). After the pump is turned on but before the turning on of the probe, electron oscillates in the field of the pump according to $\beta_{x0} = \omega_0 a_{\text{pump}}/(\omega_0 - \Omega_0) \sin(k_0 z_0 - \omega_0 t)$. Switching on the probe does not significantly alter electron motion: $\beta_x - \beta_{x0}$ is shown as a barely visible dashed line in Fig. 2(b)). Comparing Figs. 2(a) and (b), observe that the pump suppressed electron response at the cyclotron frequency, making the plasma transparent to the probe.




FIG. 2. Numerical simulation of the single particle motion in the combined field of two EM waves with ($\omega_1 = \Omega_0, k_1 = \omega_1/c$) and ($\omega_0 = \Omega_0 - \omega_p, k_0 \approx 0.83\Omega_0/c$). Both pump and probe are slowly turned on according to Eq. (2). (a) Without the pump electron is resonantly driven by probe: β_x growth indefinitely; (b) With the pump, electron motion is almost unaffected by the probe. Solid line – total β_x ; barely visible dashed line – $(\beta_x - \beta_{x0})$, where $\beta_{x0} = \omega_0 a_{\text{pump}}/(\omega_0 - \Omega_0) \sin(k_0 z_0 - \omega_0 t)$. Since $\beta_x - \beta_{x0}$. (c) Solid line: longitudinal displacement $\Omega_0 \zeta_z/c$; dashed line: $\Omega_0(\zeta_z - \zeta_0)/c$, where $\zeta_0 = 2a_{\text{pump}}/a_{\text{probe}} \sin \omega_p t$ from Eq. (3)

Suppression is caused by the excitation of a strong plasma [shown in Fig. 2(c)] which produced a sideband of the pump at the cyclotron frequency. This sideband canceled the electric field of the probe. An approximate analytic formula for the steady-state amplitude of the plasma oscillation,

$$\zeta_0 = \frac{2a_{\text{probe}}}{k_0 a_{\text{pump}}} \sin \omega_p t, \quad (3)$$

is derived below by requiring that the sideband cancels the probe. which is in good agreement with the simulation result. Simulation results demonstrate *stability* of the steady-state values of β_x and ζ_z which are naturally reached in a collisionless plasma. Note that the pump has to be switched on prior to the arrival of the probe. In atomic physics, this pulse sequence is referred to as “counter-intuitive” [2].

Maintaining high-power pumping waves in the plasma may prove challenging in practice. For example, supporting $a_0 = 0.01$ over an area $A = (2\pi c/\omega_0)^2$ requires microwave power of 3 megawatts. Fortunately, for $\omega_p = \Omega_0$, a magnetostatic helical undulator can replace a microwave beam. We simulated electron motion in the combined field of an undulator with $a_0 = 0.1$ and $k_0 = 2\Omega_0/c$, and a probe which is switched on according to $a_{\text{probe}} = 0.5a_1 (1 + \tanh[(\Omega_0 t - 270)/60])$, where $a_1 = 0.01$. Suppression of the electron response at the cyclotron frequency is apparent from Fig. 3(a). Electric field of the probe is canceled by the $(\dot{\zeta}_z/c) \vec{e}_z \times \vec{B}_{0\perp}$ force which is exerted on a longitudinal plasma wave by the helical magnetic field of the undulator.

Steady-state values of $\beta_+ = \beta_-^* = \beta_x - i\beta_y$ and ζ_z can be analytically obtained by linearizing Eq. (1) in the weak probe $a_1 \ll a_0$ limit.

$$\begin{aligned} \dot{\beta}_+ + i\Omega_0 \beta_+ = & \\ - \left(\omega_0 a_0 e^{i\bar{\theta}_0} + \omega_1 a_1 e^{i\bar{\theta}_1} - k_0 a_0 \dot{\zeta}_z e^{i\bar{\theta}_0} - k_1 a_1 \dot{\zeta}_z e^{i\bar{\theta}_1} \right). \end{aligned} \quad (4)$$

Introducing $\theta_{0,1} = k_{0,1} z_0 - \omega_{0,1} t$ and assuming that $k_{0,1} \zeta_z < 1$, exponentials in Eq. (4) are expanded as $e^{i\bar{\theta}_{0,1}} \approx e^{i\bar{\theta}_{0,1}} (1 + ik_{0,1} \zeta_z)$, yielding

$$\begin{aligned} \dot{\beta}_+ + i\Omega_0 \beta_+ = & -\omega_0 a_0 e^{i\bar{\theta}_0} (1 + ik_0 \zeta_z - k_0 \dot{\zeta}_z/\omega_0) \\ & -\omega_1 a_1 e^{i\bar{\theta}_1} (1 + ik_1 \zeta_z - k_1 \dot{\zeta}_z/\omega_1). \end{aligned} \quad (5)$$

Longitudinal equation of motion is given by

$$\ddot{\zeta}_z + \omega_p^2 \zeta_z \approx -\frac{e}{mc} \left(\vec{v}_\perp \times \vec{B}_\perp + \zeta_z \vec{v}_\perp \times \frac{\partial \vec{B}_\perp}{\partial z} \right),$$

where $\vec{B}_\perp(z, t)$ was expanded as $\vec{B}_\perp(z_0 + \zeta_z) \approx \vec{B}_\perp(z_0) + \zeta_z \partial_{z_0} \vec{B}_\perp(z_0)$ to first order in ζ_z . Inserting the expression for \vec{B}_\perp , obtain

$$\ddot{\zeta}_z + \omega_p^2 \zeta_z = -\frac{c^2}{2} (k_0 a_0 \beta_- e^{i\theta_0} + k_1 a_1 \beta_- e^{i\theta_1} - i k_0^2 \zeta_z \beta_- a_0 e^{i\theta_0} - i k_1^2 \zeta_z a_1 e^{i\theta_1}) + c. c. \quad (6)$$

The last term in the RHS of Eq. (6) will be later dropped because it is proportional to the product of two small quantities, ζ_z and a_1 . Note that, unlike the transverse velocity β_+ which is excited directly by each of the two lasers according to Eq. (5), plasma waves are excited only in the presense of *two* lasers via the beatwave mechanisms.

The physical reason for EIT in plasma is the strong coupling between longitudinal and transverse degrees of freedom of the plasma electrons. The steady-state solution of Eq. (6) $\zeta_z = 0.5\tilde{\zeta} \exp i(\Delta k z - \Delta \omega t) + c. c.$, where $\Delta \omega = \omega_1 - \omega_0$ and $\Delta k = k_1 - k_0$, is substituted into the transverse equation of motion (5). Retaining the terms with $\exp -i\omega_0 t$ and $\exp -i\omega_1 t$ dependence results in

$$\beta_+ = -\frac{i\omega_0 a_0}{\omega_0 - \Omega_0} e^{i\theta_0} - \frac{i\omega_1}{\omega_1 - \Omega_0} \left(a_1 + \frac{ik_0 \tilde{\zeta}}{2} a_0 \right) e^{i\theta_1}. \quad (7)$$

Applying Eq. (7) to the simulated earlier case of $\omega_1 = \Omega_0$ and $\Delta \omega = \omega_p$ yields the steady-state amplitude of the plasma wave given by Eq. (3).

In the general case of $\omega_1 \neq \Omega_0$ we insert β_+ and β_- into Eq. (6) yielding

$$(\omega_p^2 - \Delta \omega^2) \tilde{\zeta} = ic^2 \left[\frac{k_0 a_0^* \omega_1}{\omega_1 - \Omega_0} (a_1 + ik_0 \tilde{\zeta} a_0 / 2) - \frac{k_1 a_1 \omega_0}{\omega_0 - \Omega_0} a_0^* - i \frac{k_0^2 \tilde{\zeta} \omega_0}{\omega_0 - \Omega_0} |a_0|^2 \right], \quad (8)$$

where $\theta_1 - \theta_0 = (k_1 - k_0)z_0 - \Delta \omega t$. Equation (8) is then solved for $\tilde{\zeta}$ which is substituted into Eq. (7) yielding the steady-state value of β_+ :

$$\beta_{+s} = -\frac{i\omega_0 a_0}{\omega_0 - \Omega_0} e^{i\theta_0} - i \frac{\omega_1 a_1}{\omega_0 - \Omega_0} e^{i\theta_1} \times \frac{c^2 k_0^2 \omega_0 |a_0|^2 (k_1/k_0 - 2) + 2(\omega_p^2 - \Delta \omega^2)(\omega_0 - \Omega_0)}{c^2 k_0^2 |a_0|^2 \omega_1 + 2(\omega_p^2 - \Delta \omega^2)(\omega_1 - \Omega_0)}, \quad (9)$$

where we have neglected terms proportional to the product of laser detuning $\delta \Omega = \omega_1 - \Omega_0$ from resonance and the pump intensity a_0^2 . Qualitatively, the pump influence is strong only close to the cyclotron resonance,

and is negligible far from $\omega_1 = \Omega_0$. From Eq. (9), plasma is resonantly driven when the denominator $D = 2(\omega_p^2 - \Delta \omega^2)(\omega_1 - \Omega_0) + c^2 k_0^2 |a_0|^2 \omega_1$ vanishes. Close to cyclotron resonance $D \approx 4\omega_p(\Omega_R^2 - \delta \Omega^2)$, where $\Omega_R = ck_0 a_0 (\Omega_0 / 4\omega_p)^{1/2}$ is the effective Rabi frequency. Hence, the modified plasma resonances are shifted from $\omega_1 = \Omega_0$ to $\omega_1 = \Omega_0 \pm \Omega_R$.




FIG. 3. Same as Fig. 2, except $\omega_p = \Omega_0$, $\omega_0 = 0$, $k_0 \approx 2\Omega_0/c$ (static helical undulator is switched on from the start). (a) Transverse velocity β_x and (b) longitudinal displacement $\Omega_0 \zeta_z / c$ during and after the turning on of the probe.

Fluid velocity component $b_+ \approx \beta_{+s} - \partial_z(\zeta_z \beta_{+s}) = \beta_+ - ik_1 \zeta_z \beta_+$ proportional to $\exp i\theta_1$ is given by $b_+ = ia_1 \omega_1 (\delta \Omega + \delta \Omega_0) / (\Omega_R^2 - \delta \Omega^2)$ where $\delta \Omega_0(k_1) = (2\Omega_R^2 \omega_0 / \omega_p \Omega_0)(k_1/k_0 - 1)$.

Dispersion relation for classical EIT in magnetized plasma is derived from the wave equation for the probe $-(c^2 \partial_z^2 - \partial_t^2) \vec{E} = 4\pi \partial_t \vec{J}$, where the rhs is equal to $-4\pi ic \omega_1 n_0 b_+ + c. c.$:

$$\omega_1^2 = c^2 k_1^2 - \omega_p^2 \omega_1 \frac{\delta \Omega + \delta \Omega_0(k_1)}{\Omega_R^2 - (\delta \Omega)^2}, \quad (10)$$

where it was assumed that the frequency of the pump is fixed at $\omega_0 = \Omega_0 - \omega_p$. Complete transparency ($\omega_1 = k_1^2 c^2$) is achieved at $\omega_1 = \Omega_0 - \delta \bar{\Omega}_0$, where $\delta \bar{\Omega}_0 \approx (2\omega_0 \Omega_R^2 / \omega_p \Omega_0)(\Omega_0/k_0 c - 1)$. Note that this frequency shift is in general very small in the most interesting regime of $\Omega_R \ll \omega_p$: $|\delta \Omega_0| < 4\Omega_R^2 / \omega_p \ll \Omega_R$, and can be even smaller near cyclotron resonance when pump and probe co-propagate. Equation (10) reduces to the dispersion relation for a single probe in magnetized plasma for large detunings $(\delta \Omega)^2 \gg \Omega_R^2$. The influence of the pump is significant only in the vicinity of $\delta \Omega = 0$.

Note that index of refraction is not identically equal to unity at the cyclotron resonance. This is different from the quantum-mechanical result for a three-level system [11], where $\omega_1 = k_1 c$ on resonance. It can be demonstrated that this difference occurs because multiple Landau levels $E_n = n\hbar \Omega_0$ and corresponding Raman-shifted

levels $E_{Rn} = E_n + \hbar\omega_p$ participate in the classical EIT.

Dispersion relation given by Eq. (10) is plotted in Fig. (4) for the same plasma parameters as in Fig. (1), plus a co-propagating pump with $\Omega_R = 0.5\omega_p$. The flat band between the $\Omega_0 \pm \Omega_R$ resonant frequencies is a novel feature which is not present without the pump (compare with Fig. (1)). The width of this EIT band proportional to $\Omega_R \propto a_0$ can become very narrow for low pump amplitude. The corresponding “group velocity” (understood in a strictly geometrical sense explained below) $v_g = \partial\omega_1/\partial k_1 \approx 2c\Omega_R^2/\omega_p^2$ can also be made arbitrarily small. Slowly propagating wavepacket of electromagnetic waves is a classical analog of the “slow light” in atomic systems [3].

Qualitatively, the spectacular slowing down of EM waves in the EIT plasma can be understood by considering the entrance of a probe beam of duration L_0 into the plasma. In steady state inside the plasma, the “slow light” wavepacket of length L_f consists of the transversely polarized field of the probe $|\vec{E}_1| = |\vec{B}_1| = a_1 mc\omega_1/e$ and the longitudinal electric field of the plasma wave $E_z = 4\pi en_0(2a_1/k_0 a_0)$. As the pulse enters the plasma, it loses photons to the pump at the same rate as new plasmons are created (according to the Manley-Rowe relation). Classical photon density of a field with frequency ω is proportional to the action density $\propto U/\omega$, where U is the energy density. We calculate that the ratio of the plasmon to photon density inside the “slow light” pulse,

$$\frac{U_{\text{plas}}/\omega_p}{U_{\text{phot}}/\omega_1} = \frac{\Omega_0}{\omega_p} \frac{E_z^2}{2E_1^2} = \frac{\omega_p^2}{2\Omega_R^2}, \quad (11)$$

is $\gg 1$ if $\Omega_R \ll \omega_p$. Thus, most photons of the original pulse are lost to the pump. Since the index of refraction remains close to unity, so is the photon energy density. Therefore, the loss of photons is due to the spatial shortening of the pulse from L_0 to $L_f = L_0 \times (2\Omega_R^2/\omega_p^2)$. Because temporal pulse duration does not change, we recover the previously calculated $v_g/c = 2\Omega_R^2/\omega_p^2$. It is precisely in this geometric sense of $v_g/c = L_f/L_0$ that the group velocity of the slow light is interpreted. v_g is not related to the speed of individual photons since their number is not conserved during the pulse transition into the plasma.




FIG. 4. EIT dispersion curve, $\Omega_0/\omega_p = 3$ and $\Omega_R/\omega_p = 1/2$. Flat band above $\Omega_0 - \Omega_R$ up to $\Omega_0 + \Omega_R$ labeled “EIT Band” corresponds to “slow light” and appears only in the presence of a pump.

One interesting application of EIT in magnetized plasma is ion acceleration. While laser-plasma accelerators of electrons [12] have long been considered as a long-term alternative to conventional rf cavity-based linacs, the field of plasma-based ion accelerators is still in its infancy [13]. EIT enables one to conceive a short-pulse ion accelerator which consists of a “slow light” pulse in plasma with approximately equal group and phase velocities. Acceleration is accomplished by the longitudinal electric field of the plasma wave. Counter-propagating geometry is chosen to match the phase and group velocities because $v_{\text{ph}} = \omega_p/|k_0| + k_1 \approx 0.5c\omega_p/\Omega_0$. Matching $v_{\text{ph}} = v_g$ yields $a_0 \approx \omega_p^2/\Omega_0^2 \ll 1$. Other types of accelerators based on the “slow light” which rely on the ponderomotive force also appear attractive because the ponderomotive force, which scales as the gradient of the energy density $E_z^2/L_f \propto (\omega_p/\Omega_0)U_0/v_g^2$, increases rapidly with decreasing group velocity of the probe.

-
- [1] K. J. Boller, A. Imamoglu, and S. E. Harris, Phys. Rev. Lett. **66**, 2593 (1991); S. E. Harris, Phys. Rev. Lett. **70**, 552 (1993).
 - [2] S. E. Harris, Physics Today **7**, 36 (1997); J. P. Marangos, Journ. Modern Optics **45**, 471 (1998); A. B. Matsko *et. al.*, *Advances in Atomic, Molecular, and Optical Physics* **46**, 191 (2001).
 - [3] L. V. Hau *et. al.*, Nature **397**, 594 (2001).
 - [4] M. Fleischhauer, S. F. Yelin, and M. D. Lukin, Opt. Commun. **179**, 395 (2000).
 - [5] M. D. Lukin, S. F. Yelin, and M. Fleischhauer, Phys. Rev. Lett. **84**, 4232 (2000).
 - [6] A. B. Matsko, Y. Rostovtsev, M. Fleischhauer, and M. O. Scully, Phys. Rev. Lett. **86**, 2006 (2001).
 - [7] U. Leonhard and P. Piwnicki, Phys. Rev. Lett. **84**, 822 (2000).
 - [8] N. A. Krall and A. W. Trivelpiece, *Principles of Plasma Physics*, Chapt. 4 (McGraw-Hill, New York, 1973).
 - [9] J. Dawson, Phys. Rev. **113**, 383 (1959).
 - [10] M. N. Rosenbluth, C. S. Liu, Phys. Rev. Lett. **29**, 701 (1972).
 - [11] M. O. Scully, Phys. Rev. Lett. **67**, 1855 (1991); M. Fleischhauer, C. H. Keitel, and M. O. Scully, Phys. Rev. A **46**, 1468 (1992).
 - [12] T. Tajima and J. M. Dawson, Phys. Rev. Lett. **43**, 267 (1979).
 - [13] R. A. Snavely *et. al.*, Phys. Rev. Lett. **85**, 2945 (2000).