The configuration of a topological current and physical structure: an application and paradigmatic evidence

Li-Bin Fu¹, Jie Liu¹, Shi-Gang Chen¹, and Yi-Shi Duan²

¹ Institute of Applied Physics and Computational Mathematics, P.O. Box 8009(28), Beijing 100088, P.R. China

² Institute of Theoretical Physics, Department of Physics, Lanzhou University, 730000, P. R. China

In the ϕ -mapping theory, the topological current constructed by the order parameters can possess different inner structure. The difference in topology must correspond to the difference in physical structure. The transition between different structures happens at the bifurcation point of the topological current. In a self-interaction two-level system, the change of topological particles corresponds to change of energy levels.

In recent years, topology has established itself as an important part of the physicist's mathematical arsenal [1]. The concepts of the topological particle and its current have been widely used in particle physics [2,3] and topological defect theory [4]. Here, the topological particles are regarded as abstract particles, such as monopoles and the points defects.

In this paper, we give a new understanding in topology and physics. Many physics system can be described by employing the order parameters. By making use of the ϕ -mapping theory, we find that the topological current constructed by the order parameters can possess different inner structure. The topological properties are basic properties for a physics system, so the difference in configuration of the topological current must correspond to the difference in physical structure.

Considering a (n+1)-dimensional system with *n*-component vector order parameter field $\vec{\phi}(\mathbf{x})$, where $\mathbf{x} = (x^0, x^1, x^2, \cdots x^n)$ correspond to local coordinates. The direction unit field of $\vec{\phi}$ is defined by

$$n^a = \frac{\phi^a}{||\phi||}, \quad a = 1, 2, \cdots n \tag{1}$$

where

$$||\phi|| = (\phi^a \phi^a)^{1/2}.$$

The topological current of this system is defined by

$$j^{\mu}(x) = \frac{\in^{\mu\mu_1\cdots\mu_n}}{A(S^{n-1})(d-1)!} \in_{a_1\cdots a_n} \partial_{\mu_1} n^{a_1}\cdots \partial_{\mu_n} n^{a_n}$$
(2)

where $A(S^{n-1})$ is the surface area of (n-1)-dimensional unit sphere S^{n-1} . Obviously, the current is identically conserved,

$$\partial_{\mu}j^{\mu} = 0$$

If we define a Jacobians by

$$\in^{a_1\cdots a_n} D^{\mu}(\frac{\phi}{x}) = \in^{\mu\mu_1\cdots\mu_n} \partial_{\mu_1}\phi^{a_1}\cdots\partial_{\mu_n}\phi^{a_n}, \qquad (3)$$

as has been proved before [5], this current takes the form as

$$j^{\mu} = \delta(\vec{\phi}) D^{\mu}(\frac{\phi}{x}). \tag{4}$$

Then, we can obtain

$$j^{\mu} = \sum_{i=1}^{l} \beta_{i} \eta_{i} \delta(\vec{x} - \vec{z}_{i}(x^{0})) \frac{dz_{i}^{\mu}}{dx^{0}}, \qquad (5)$$

in which $z_i(x^0)$ are the zero lines where $\vec{\phi}(\mathbf{x}) = 0$, the positive integer β_i and $\eta_i = sgnD(\frac{\phi}{\vec{x}})$ are the Hopf index and Brouwer degree of ϕ -mapping [7] respectively, and l is the total number of the zero lines. This current is similar to a current of point particles and the *i*-th one with the charge $\beta_i \eta_i$, and the zero lines $z_i(x)$ are just the trajectories of the particles, for convenience we define these point particles as topological particles. Then the total topological charge of the system is

$$Q = \int_M j^0 d^n x = \sum_{i=1}^l \beta_i \eta_i,$$

here M is a *n*-dimensional spatial space for a given x^0 . This is a topological invariant and corresponds to some basic conditions of this physical system. However, it is important that the inner structure of the topological invariant can be constructed in different configurations, i.e., the number of the topological particles and their charge can be changed. This change in configuration of the topological current must correspond to some change in physical structure.

All of the above discussion is based on the condition that

$$D\left(\frac{\phi}{x}\right) = \left.D^0\left(\frac{\phi}{x}\right)\right|_{z_i} \neq 0.$$

When $D\left(\frac{\phi}{x}\right)\Big|_{z_i} = 0$ at some points $p_i^* = z^*(x_c^0)$ at $x^0 = x_c^0$ along the zero line $z_i(x^0)$, it is shown that there exist several crucial cases of branch process, which correspond to the topological particle generating or annihilating at limit points and splitting, encountering or merging at the bifurcation points. A vast amount of literature has been devoted to discussing these features of the evolution of the topological particles [8]. Here, we will not spend more attention on describing these evolution, but put our attention on the physical substance of these processes.

As we have known before, all of these branch processes keep the total topological charge conserved, but it is very important that these branch processes change the number and the charge of the topological particles. i.e. change the inner structure of the topological current. In our point of view, the different configuration of topological current corresponds to the different physical structure.

We consider x^0 as a parameter λ of a physics system. Let us define

$$f_i(\lambda) = D^0 \left(\frac{\phi}{x}\right)\Big|_{z_i} \tag{6}$$

As λ changing, the value of $f_i(\lambda)$ changes along the zero lines $z_i(\lambda)$. At a critical point $\lambda = \lambda_c$, when $f_i(\lambda_c) = 0$, we know that the inner structure of the topological current will be changed in some way, at the same time the physical structure will also be changed, i.e., the physical structure when $\lambda < \lambda_c$ will be different from the one when $\lambda > \lambda_c$. The transition between these structures occurs at the bifurcation points where $f_i(\lambda) = 0$. As an application and example, let us consider a self-interacting two-level model introduced in Ref. [9]. The nonlinear two-level system is described by the dimensionless Schrödinger equation

$$i\frac{\partial}{\partial t} \begin{pmatrix} a\\ b \end{pmatrix} = H(\gamma) \begin{pmatrix} a\\ b \end{pmatrix}$$
(7)

with the Hamiltonian given by

$$H(\gamma) = \begin{pmatrix} \frac{\gamma}{2} + \frac{C}{2}(|b|^2 - |a|^2) & \frac{V}{2} \\ \frac{V}{2} & -\frac{\gamma}{2} - \frac{C}{2}(|b|^2 - |a|^2) \end{pmatrix},$$
(8)

in which γ is the level separation, V is the coupling constant between the two levels, and C is the nonlinear parameter describing the interaction. The total probability $|a|^2 + |b|^2$ is conserved and is set to be 1.

We assume $a = |a|e^{i\varphi_1(t)}$, $b = |b|e^{i\varphi_2(t)}$, the fractional population imbalance and relative phase can be defined by

$$z(t) = |b|^2 - |a|^2, \quad \varphi(t) = \varphi_2(t) - \varphi_1(t).$$
 (9)

From Eqs. (7) and (8), we obtain

$$\frac{d}{dt}z(t) = -V\sqrt{1-z^2(t)}\sin[\varphi(t)]$$
(10)

$$\frac{d}{dt}\varphi(t) = \gamma + Cz(t) + \frac{Vz(t)}{\sqrt{1 - z^2(t)}}\cos[\varphi(t)].$$
(11)

If we chose $x = 2|a||b|\cos(\varphi)$, $y = 2|a||b|\sin(\varphi)$, it is easy to see that $x^2+y^2+z^2 = 1$ by considering $|a|^2 + |b|^2 = 1$, which describes a unit sphere S^2 with z and φ a pair of co-ordinates. We define a vector field on this unit sphere:

$$\phi^1 = -V\sqrt{1-z^2}\sin(\varphi),\tag{12}$$

$$\phi^2 = \gamma + Cz + \frac{Vz}{\sqrt{1 - z^2}}\cos(\varphi) \tag{13}$$

Apparently, there are singularities at the two pole points $z = \pm 1$, which make the vector $\vec{\phi}$ is discontinuous at these points. However, the direction unit vector \vec{n} is

continuous. In the ϕ -mapping theory, we only need the unit vector \vec{n} is continuous and differentiable on the whole sphere S^2 (at the zero points of $\vec{\phi}$, the differential of \vec{n} is a general function), and the vector $\vec{\phi}$ is continuous and differentiable at the neighborhoods of its zero points. Then from ϕ -mapping theory, we can obtain a topological current as

$$\vec{j} = \sum_{i=1}^{l} \beta_i \eta_i \delta(\varphi - \varphi_i(\gamma)) \delta(z - z_i(\gamma)) \frac{dz_i}{d\gamma} |_{p_i} , \qquad (14)$$

in which $p_i = p_i(z_i, \varphi_i)$ is the trajectory of the *i*-th topological particle P_i and

$$\eta_{i} = sgn\left(D(\gamma)\right) = sgn\left(\det\left(\begin{array}{cc}\partial\phi^{1}/\partial\varphi & \partial\phi^{2}/\partial\varphi\\\partial\phi^{1}/\partialz & \partial\phi^{2}/\partialz\end{array}\right)\Big|_{p_{i}}\right).$$
(15)

From Eqs. (12) and (13), it is easy to see that ϕ is single-valued on S^2 , which states that Hopf index $\beta_i = 1$ $(i = 1, 2, \dots, l)$ here. It can be proved that the total charge of this system is just the Euler number of S^2 which is 2 [6]. This is a topological invariant of S^2 which corresponds to the basic condition of this system: $|a|^2 + |b|^2 = 1$. The following discussion can show that this topological invariant can possess different configuration when γ changes. This difference in topology corresponds to the change in adiabatic levels of this nonlinear system.

We can prove that every topological particle corresponds to an eigenstate of the nonlinear two-level system. By solving $\vec{\phi} = 0$ form (12) and (13), we find there are two different cases for discussing.

Case 1. For $|C/V| \leq 1$, there only two topological particles P_1 and P_2 , which locate on the line $\varphi = 0$ and $\varphi = \pi$ respectively as shown in the upper panel of Fig.1. All of them with topological charge +1, and $D(\gamma)|_{P_{1,2}} > 0$ for any γ . Correspondingly, in this case, there are only two adiabatic energy levels in this nonlinear two-level mode for various γ [9], as shown in the lower panel of Fig.1, P_1 corresponds to the upper level and P_2 corresponds to the lower level.

Case 2, For C/V > 1, two more topological particles can appear when γ lies in a window $-\gamma_c < \gamma < \gamma_c$. The boundary of the window can be obtained by assuming $D(\gamma)|_{P_i} = 0$, yielding

$$\gamma_c = (C^{2/3} - V^{2/3})^{3/2}.$$
(16)

The striking feature happens at $\gamma = -\gamma_c$, there exists a critical point $T_1^*(z_c, \pi)$ with $D(\gamma_c)|_{T_1^*} = 0$, as we have shown in Ref. [8], we can prove that this point is a limit point, and a pair of topological particles P_3 and P_4 generating with opposite charge -1 and +1 respectively, both of the new topological particles lie on the line $\varphi = \pi$. One of the original topological particle, P_2 with charge +1 on the line $\varphi = \pi$, moves smoothly up to $\gamma = \gamma_c$, where it collides with P_3 and annihilates with it at another limit point $T_2^*(-z_c, \pi)$. The other, P_1 , which lies on the line $\varphi = 0$, still moves safely with γ .

As pointed out by B. Wu and Q. Niu [9], when the interaction is strong enough (C/V > 1), a loop appears at the tip of the lower adiabatic level when C/V > 1 while $-\gamma_c \leq \gamma \leq \gamma_c$. We show the interesting structure in Fig. 2 in which C/V = 2. For $\gamma < -\gamma_c$, there are two adiabatic levels, the upper level corresponds to the topological particle P_1 , the lower one corresponds to the topological particle P_2 ; for $\gamma > \gamma_c$, there are also only two adiabatical levels, but at this time the lower one corresponds to P_4 while the upper one still corresponds to P_1 . The arc part of the loop on the tip of lower level when $-\gamma_c < \gamma < \gamma_c$ just corresponds to P_3 , which merges with the level corresponding to P_4 at the point M on the left and with the one corresponding to P_1 at the point T on the right.

From the above discussion, one finds that when the structure of the topological current is changed by generating or annihilating a pair of topological particles (the upper panel of Fig. 2), at the same time, the physical structure is changed by adding two energy levels or subtracting two energy levels respectively (the lower panel of Fig.2). The critical behaviors happen at the limit points where $D(\gamma_c)|_{T_{1,2}^*} = 0$.

In fact, this nonlinear two-level model is of comprehensive interest for it associates with a wide range of concrete physical systems, e.g., BEC in an optical lattice [10] or in a double-well potential [11], and the motion of small polarons [12]. So, the relation between topological particles and physical inner structure can be observed in experimental methods. Here we propose a perspective system for observing the striking phenomenon: a Bose-Einstein condensate in a double-well potential [11,13]. The amplitudes of general occupations $N_{1,2}(t)$ and phases $\varphi_{1,2}$ obey the nonlinear two-mode Schrödinger equations, approximately [13],

$$i\hbar \frac{\partial \phi_1}{\partial t} = (E_1^0 + U_1 N_1) \phi_1 - K \phi_2$$

$$i\hbar \frac{\partial \phi_2}{\partial t} = (E_2^0 + U_2 N_2) \phi_2 - K \phi_1$$
(17)

with $\phi_{1,2} = \sqrt{N_{1,2}}exp(i\varphi_{1,2})$, and total number of atoms $N_1 + N_2 = N_T$, is conserved. Here $E_{1,2}^0$ are zero-point energy in each well, $U_{1,2}N_{1,2}$ are proportional to the atomic self-interaction energy, and K describes the amplitude of the tunnelling between the condensates. After introducing the new variables $z(t) = (N_2(t) - N_1(t))/N_T$ and $\varphi = \varphi_2 - \varphi_1$, one also obtain an equations having the same form as Eqs. (7) and (8) except for the parameters replaced by

$$\gamma = -[(E_1^0 - E_2^0) - (U_1 - U_2)N_T/2]/\hbar, \qquad (18)$$

$$V = 2K/\hbar, \ C = (U_1 + U_2)N_T/2\hbar.$$
 (19)

With these explicit expressions, our theory and results can be directly applied to this system without intrinsic difficulty. In this system the topological particles can be located by the stable occupation and relative phase $\varphi = 0$ or $\varphi = \pi$ for a give parameter γ . And, one can draw the zero line of each topological particle by giving different γ . We hope our discussions will stimulate the experimental works in the direction.

We note that for a system the global property (topology) is given, the interesting feature is that under the same topology the topological configuration can be different, this difference must correspond the different physical structure. This relation between topological configuration and physical structure gives an important property to classify some physical system which contains many different structures.

We thank Prof. B.Y. Ou for useful discussions. This project was supported by Fundamental Research Project of China.

- S. Nash and S. Sen, Topology and Geometry for Physicists (Academic, London, 1983);
 A.S. Schwarz, Topology for Physicists (Springer-Verlag, Berlin, 1994). G. Morandi, The Role of Topology in Classical and Quantum Physics, Lecture Note in Physics (Springer-Verlag, Berlin, 1992) Vol. M7.
- [2] G. 't Hooft, Nucl. Phys. B 79, 276 (1974); J. Wess and B. Zumino, Phys. Lett. B 37, 95 (1971)
- [3] L. Faddeev and A. Niemi, Nature (London) 387, 58 (1997); R.A. Battye and P.M.
 Sutcliffe. Phys. Rev. Lett. 81, 4798 (1998); H. Aratyn, L.A. Ferreira and A. H. Zimerman, Phys. Rev. Lett. 83, 1723 (1999)
- [4] N.D. Mermin, Rev. Mod. Phys. 51, 591 (1979); A.J. Bray, Adv. Phys. 43, 375 (1994);
 Phys. Rev. E 55. 5297 (1997); P.W. Anderson, *Basic Notations of Condensed Matter Physics* (Benjamin, London, 1984)
- [5] Y.S. Duan, T. Xu and L.B. Fu, Progress of Theoretical Physics **101**(2), 467 (1999).
- [6] Y.S. Duan, S. Li and G.H. Yang, Nucl. Phys. B 514, 705 (1998).
- [7] H. Hopf, Math. Ann. 96, 209 (1929); Y.S. Duan, X.H. Meng, J. Math. Phys. 34, 1149 (1993).
- [8] L.B. Fu, Y.S. Duan and H. Zhang, Phys. Rev. D 62 (2000); Y.S. Duan, H. Zhang and L.B. Fu, Phys. Rev. E 58 528 (1999); Y.S. Duan, H. Zhang and G. Jian, Phys. Lett. A 253, 57 (1999).
- [9] B. Wu and Q. Niu, Phys. Rev. A **61** 023402 (2000).
- [10] M.H. Anderson et al., Science 269, 198 (1995); Dae-I1 Choi and Q. Niu, Phys. Rev. Lett. 82, 2022 (1999); W. Zhang and D.F. Walls, Phys. Rev. A 49, 3799 (1994); W. Zhang and D.F. Walls, ibid. 52, 4696 (1995)

- [11] M.R. Andrews et al., Science 275, 637 (1997); G.J. Milburn et al. Phys. Rev. A 55, 4318 (1997); M.O. Mewes et al., Phys. Rev. Lett. 78, 582 (1997)
- [12] J.C. Eilbeck, P.S. Lomdahl, and A.C. Scott, Physica D 16, 318 (1985); V.M. Kenkre and D.K. Campbell, Phys. Rev. B 34, 4959 (1986); P.K. Datta and K. Kundu, ibid. 53, 14929 (1996)
- [13] S.Raghavan, A.Smerzi, S.Fantoni, and S.R.Shenoy, Phys. Rev. A 59, 620 (1999)

I. FIGURE CAPTION

Fig. 1. (a) The projecting of trajectory of topological particles on $(z - \gamma)$ plane for C/V = 0. P_i denotes the *i*-th topological particle. (b). The energy levels for C/V = 0. Each level is labelled by the topological particle which corresponds to it.

Fig. 2. (a) The projecting of trajectory of topological particles on $(z - \gamma)$ plane for C/V = 2. P_i denotes the *i*-th topological particle. (b). The energy levels for C/V = 2. Each level is labelled by the topological particle which corresponds to it.





Fig. 2