Comment: Anomalous Anisotropic Light Scattering in Ge-Doped Silica Glass

Saulius Juodkazis,* Shigeki Matsuo, and Hiroaki Misawa

Faculty of Engineering, The University of Tokushima,

2-1 Minamijyosanjima, Tokushima 770-8506, Japan

(Dated: November 20, 2018)

An anomalous anisotropic light scattering which peaks in the plane of light polarization, the "propeller effect" named by authors, was observed in glass pumped by intense femtosecond laser radiation [1]. A quiver movement of photoelectrons in the light field was the key to an explanation of the effect. A blue emission pattern of a propeller shape was attributed to a defect-related photoluminescence (PL) and explained by its scattering. The emission follows the angular distribution of the photoelectrons given by $\frac{d\sigma}{d\Omega} \propto (1 + \cos^2 \varphi)$, where the $\frac{d\sigma}{d\Omega}$ is the differential scattering cross section of electrons, φ is the angle between the field amplitude vector **E** and the electron momentum \mathbf{k}_e .

This is an interesting phenomenon relevant to a fast growing research on the interaction on ultra-short pulses with materials, in particular, with transparent solids. This is a field where some long-lasting controversies over generation of white-light continuum (super-continuum (SC)), self-focusing and dielectric breakdown (DB) mechanisms should be resolved. By this comment we intend to draw attention to the phenomenon of *bremsstrahlung* (German word for "braking radiation"), which escapes proper consideration and can explain, for example, emission spectra of dielectric breakdown and the above mentioned "propeller effect" [1]. In our judgement, the explanation of the "propeller effect" should take into account *bremsstrahlung*, transmission function of a measurement setup, and optical aberrations.

Once the optical excitation generates free electrons, this not only changes the dielectric function, which is usually considered in the SC and light-induced DB mechanisms from the point of view of nonlinear optics, but also inherently causes the light emission due to the *bremsstrahlung*. In a plasma state, after the dielectric breakdown, when free charges are available, the radiated energy, W, per unit angular frequency, ω , in the nonrelativistic case is given by: [2]

$$\frac{dW}{d\omega} = \frac{e^2}{6\pi^2\varepsilon_0 c^3} \left| \int_{-\infty}^{\infty} \dot{v} e^{i\omega t} dt \right|^2, \tag{1}$$

where \dot{v} is the acceleration of an electron, which has a charge of e, c - speed of light, and ε_0 - permittivity of vacuum. Spectral extent of *bremsstrahlung* depends on the type of electron-ion collisions (Fig. 1) and can be defined as a reciprocal of the collision time τ . In the case of straight-line collisions $\tau^{-1} \simeq v/(2b)$ while for the parabolic ones $\tau^{-1} \simeq (v/(2b))(2b_{90}/b)^2$ [2], v is the velocity of incoming electron. Emission spectrum



FIG. 1: (a) Geometry of electron-ion collision. Time dependence of acceleration for straight-line, $b \ll b_{90}$, (b) and parabolic $b \gg b_{90}$ (c) collisions (b is the impact parameter; b_{90} is the electron-ion distance at right angle scattering). At t = 0 the particles are at the closest.

of *bremsstrahlung* is continuous and spreads from Xrays-to-IR. For an external observer (on z-axis) its spectral content depends on the bandgap of the host material, defects' absorption, PL, fractoluminescence, sonoluminescence, and a transmission function of a measurement setup. Time resolved spectroscopy could, e.g., distinguish among those different contributions. For an isotropic photoelectron distribution in plasma the eqn. 1 defines an emission spectrum integrated over all solid angles.

The actual polarization and angular dependence of the radiation of quiver photoelectrons is defined by $\mathbf{R} \times (\mathbf{R} \times \dot{\mathbf{v}})$ [2], here \mathbf{R} is the vector from the charge to the field point (only a far-field contribution is considered). Figure 1(b-c) qualitatively shows the acceleration projections in the quiver plane of photoelectron oscillating, e.g., along \mathbf{E}_x . Obviously, there are considerable accelerations in the perpendicular directions to the \mathbf{E}_x for straight-line and parabolic encounters. Bremsstrahlung due to the $\dot{\mathbf{v}}_x$ will be projected into the $\mathbf{E}_x \propto \mathbf{R}_Z \times (\mathbf{R}_Z \times \dot{\mathbf{v}}_x)$, the quiver direction, for an observer on z-axis (\vec{z} is usually a direction along the illumination/observation). Thus, the xy-projection of bremsstrahlung exactly follows the angular distibution of photoelectrons and explains the "propeller effect".

^{*} Corresponding author; Electronic address: saulius@eco.tokushima-u.ac.jp

- P. G. Kazansky, H. Inouye, T. Mitsuyu, K. Miura, J. Qui, K. Hirao, and F. Starrost, Phys. Rev. Lett. 82, 2199 (1999).
- [2] I. H. Hutchinson, Principles of Plasma Diagnostics (Cambridge University Press, New York, 1987).