

Equilibrium Distribution and Tune Shift of Beams in a Linear Collider

Bjoern S. Schmekel and Joseph T. Rogers
Cornell University, Department of Physics, Ithaca, New York 14853, USA

The evolution of two colliding beams in a linear accelerator can be described by two coupled Vlasov equations. In [1] the case without external focusing was considered. In this paper we derive the equilibrium distribution and the tune shift in the presence of external focusing. Motion is considered only in the vertical direction and the beams are presumed to be one-dimensional.

I. BEAM EVOLUTION

The beam-beam force due to the first (second) beam on the second (first) one is given by

$$\frac{dy'_{1,2}}{ds} = -\frac{4\pi Nr_e}{L_x \gamma} \int_{-\infty}^{\infty} d\bar{y} \operatorname{sgn}(y - \bar{y}) \int_{-\infty}^{\infty} d\bar{y}' \psi_{2,1}(\bar{y}, \bar{y}') \quad (1)$$

where N is the particle density in a bunch and r_e the classical radius of the electron. The distributions of the beams ψ_1 and ψ_2 are normalized to unity, i.e.

$$\int_{-\infty}^{\infty} d\bar{y} \int_{-\infty}^{\infty} d\bar{y}' \psi_{1,2}(\bar{y}, \bar{y}') = 1 \quad (2)$$

and are assumed to be one-dimensional with horizontal width L_x . Motion is considered only in the vertical direction. Then $\psi_{1,2}$ satisfy the Vlasov equations

$$\frac{\partial \psi_{1,2}}{\partial s} + y' \frac{\partial \psi_{1,2}}{\partial y} - \left(K(s)y + \frac{4\pi Nr_e}{L_x \gamma} \int_{-\infty}^{\infty} d\bar{y} \operatorname{sgn}(y - \bar{y}) \int_{-\infty}^{\infty} d\bar{y}' \psi_{2,1}(\bar{y}, \bar{y}', s) \right) \frac{\partial \psi_{1,2}}{\partial y'} = 0 \quad (3)$$

II. EQUILIBRIUM DISTRIBUTION

We are interested in finding an equilibrium (i.e. time-independent) distribution ψ_0 . The ansatz

$$\psi_1(y, y', s) = \psi_2(y, y', s) = \psi_0(y, y') \quad (4)$$

simplifies eqn. 3 significantly.

$$y' \frac{\partial \psi_0}{\partial y} - \left(K(s)y + \frac{8\pi Nr_e}{L_x \gamma} \int_{-\infty}^y d\bar{y} \int_{-\infty}^{\infty} d\bar{y}' \psi_0(\bar{y}, \bar{y}') \right) \frac{\partial \psi_0}{\partial y'} = 0 \quad (5)$$

This equation can be solved easily for a constant focusing function $K(s) = K$:

$$\psi_0(y, y') = f \left(y'^2 + Ky^2 + \frac{16\pi Nr_e}{L_x \gamma} \int dy \int_{-\infty}^y d\bar{y} \int_{-\infty}^{\infty} d\bar{y}' \psi_0(\bar{y}, \bar{y}') \right) \quad (6)$$

where f is any arbitrary function whose derivative exists. This equilibrium distribution ψ_0 must satisfy the normalization condition eqn. 2. We choose an exponential function because the beams going into the collision from a damping ring have an approximately Gaussian distribution. Dividing both sides by the derivative with respect to y' and taking the derivative with respect to y we obtain the following ordinary differential equation

$$\frac{\partial^2 Y}{\partial y^2} Y - \left(\frac{\partial Y}{\partial y} \right)^2 = -2aY^2 \left[K + \sqrt{\frac{\pi}{a}} \xi Y \right] \quad (7)$$

where $Y \equiv Y(y)$, $\xi \equiv \frac{8\pi Nr_e}{L_x \gamma}$ and the ansatz

$$\psi_0(y, y') = Y(y) e^{-ay'^2} \quad (8)$$

has been used. The function $Y(y)$ satisfies the integral relation

$$\int^{Y(y)} \frac{1}{\zeta \sqrt{-4\xi\zeta\sqrt{a\pi} - 4aK \ln(\zeta) + c_1}} d\zeta = y + c_2 \quad (9)$$

Depending on the constants $Y(y)$ describes single Gaussian-like functions, a nearly periodic sum of Gaussian-like functions or oscillating functions. Because Gaussian beams enter the collision, we are interested in the case of a Gaussian-like $Y(y)$. Thus, we choose the ansatz

$$\psi_0(y, y') = \frac{1}{2\pi\sigma} e^{-\frac{J}{\sigma}} \quad (10)$$

where J is defined for a constant focusing function $K = \frac{1}{\beta_0^2}$ as

$$J = \frac{1}{2}y'^2\beta + \frac{y^2}{2\beta} \quad (11)$$

This leads to the identification $\sigma = \frac{\beta}{2a}$. ψ_0 satisfies the normalization condition eqn. 2. Substituting eqn. 10 into eqn. 7 yields a relation for the tune shift

$$\beta_0 = \frac{\beta^{3/2}(2\pi\sigma)^{1/4} \sqrt{\sqrt{2\pi\sigma} - \xi\beta^{3/2}e^{-\frac{y^2}{2\sigma\beta}}}}{\sqrt{2\pi\beta\sigma} - \xi\beta^2e^{-\frac{y^2}{2\sigma\beta}}} = \beta \sqrt{\frac{\sqrt{2\pi\sigma}}{\sqrt{2\pi\sigma} - \xi\beta^{3/2}e^{-\frac{y^2}{2\sigma\beta}}}} \quad (12)$$

If ξ is sufficiently high the expression inside the root will become negative or - at a particular ξ - the denominator will vanish. Therefore, ξ has to obey the inequality

$$\xi < \sqrt{\frac{2\pi\sigma}{\beta^3}} \quad (13)$$

since the design tune has to be finite and real.

III. ACKNOWLEDGMENTS

As always we would like to express our gratitude to Alex Chao from SLAC for helpful discussions. This work was supported by the National Science Foundation.

[1] S. Kheifets and A. W. Chao, "Steady State Distribution for Unbunched Beams Colliding in a Drift Space", PEP-NOTE-325 (1979)