

Foldy-Wouthuysen transformation and a quasiparaxial approximation scheme for the scalar wave theory of light beams

Sameen Ahmed Khan,^{a*} Ramaswamy Jagannathan,^{b†}
Rajiah Simon^{b‡}

^a*Centro de Ciencias Físicas, Universidad Nacional Autónoma de México,
Apartado Postal 48-3, Cuernavaca 62251, Morelos, MÉXICO*

<http://www.pd.infn.it/~khan/>

^b*The Institute of Mathematical Sciences,
Central Institutes of Technology Campus, Tharamani, Chennai - 600113,
INDIA*

<http://www.imsc.ernet.in/~jagan/>

<http://www.imsc.ernet.in/~simon/>

<http://www.imsc.ernet.in/~optics/>

Abstract

The Foldy-Wouthuysen iterative diagonalization technique is applied to the Helmholtz equation to obtain a Hamiltonian description of the propagation of a monochromatic quasiparaxial light beam through a system in which the refractive index $n(x, y, z)$ varies about a background value n_0 such that $|n(x, y, z) - n_0| \ll n_0$. This technique is presented as an alternative to the conventional method of series expansion of the radical. Besides reproducing all the traditional quasiparaxial terms, this method leads to additional terms in the optical Hamiltonian.

PACS: 42.25.-p, 42.25.Bs

Keywords: Scalar wave optics, Helmholtz equation, Beam propagation, Quasiparaxial beams, Hamiltonian description, Mathematical methods of optics, Foldy-Wouthuysen transformation.

**E-mail address:* khan@ce.fis.unam.mx

†*E-mail address:* jagan@imsc.ernet.in

‡*E-mail address:* simon@imsc.ernet.in

1 Introduction

In the traditional scalar wave theory for treating monochromatic quasiparaxial light beam propagating along the positive z -axis, the z -evolution of the optical wave function $\psi(\mathbf{r})$ is taken to obey the Schrödinger-like equation

$$i\bar{\lambda}\frac{\partial\psi(\mathbf{r})}{\partial z} = \hat{H}\psi(\mathbf{r}), \quad (1)$$

where $\bar{\lambda} = \lambda/2\pi = c/\omega$. The optical Hamiltonian \hat{H} is formally given by the radical

$$\hat{H} = - \left(n^2(\mathbf{r}) - \hat{\mathbf{p}}_{\perp}^2 \right)^{1/2}, \quad (2)$$

where $\hat{\mathbf{p}} = -i\bar{\lambda}\nabla$. Then, one expands the square root in a series, with suitable ordering, or symmetrization, of the resulting polynomials in the components of \mathbf{r}_{\perp} and $\hat{\mathbf{p}}_{\perp}$, if necessary, to get a Hermitian \hat{H} [1]-[5]. Note that $n(\mathbf{r})$ and $\hat{\mathbf{p}}_{\perp} = -i\bar{\lambda}\nabla_{\perp}$ do not commute. Here we shall use the Foldy-Wouthuysen (FW) iterative diagonalization procedure, well known in the treatment of the Dirac theory of electron [6] to develop an alternative to this technique of obtaining the optical Hamiltonian.

2 The traditional paraxial formalism

Let the system we are considering have its optic axis along the z -direction and let the refractive index of the medium, $n(\mathbf{r})$, vary by a small amount around a background, or mean, value n_0 . That is, $n(\mathbf{r}) = n_0 - \varepsilon(\mathbf{r})$, $|\varepsilon(\mathbf{r})| \ll n_0$, and $n_0^2 - n^2(\mathbf{r}) \approx 2n_0\varepsilon(\mathbf{r}) \ll n_0^2$. For a monochromatic quasiparaxial beam, with leading z -dependence $\psi(\mathbf{r}) \sim \exp(in_0z/\bar{\lambda})$, we have $|\mathbf{p}_{\perp}| \ll n_0$. This means that $p_z \approx n_0$, and that all rays propagate almost parallel to the positive z -axis. Thus, the expansion of \hat{H} in the small parameters $|\mathbf{p}_{\perp}|/n_0$ and $\varepsilon(\mathbf{r})/n_0$ is basically an expansion in $1/n_0$. We have

$$\begin{aligned} \hat{H} &= - \left(n^2(\mathbf{r}) - \hat{\mathbf{p}}_{\perp}^2 \right)^{1/2} \\ &\approx - \left\{ n_0^2 - \left(\hat{\mathbf{p}}_{\perp}^2 + 2n_0\varepsilon \right) \right\}^{1/2} \\ &= -n_0 + \left(\frac{1}{2n_0}\hat{\mathbf{p}}_{\perp}^2 + \varepsilon \right) + \frac{1}{2n_0} \left(\frac{1}{2n_0}\hat{\mathbf{p}}_{\perp}^2 + \varepsilon \right)^2 \\ &\quad + \frac{1}{2n_0^2} \left(\frac{1}{2n_0}\hat{\mathbf{p}}_{\perp}^2 + \varepsilon \right)^3 + \frac{5}{8n_0^3} \left(\frac{1}{2n_0}\hat{\mathbf{p}}_{\perp}^2 + \varepsilon \right)^4 + \dots \end{aligned} \quad (3)$$

If the terms up to first order in $1/n_0$ only are retained in \hat{H} , dropping the terms proportional to the second and higher powers of $1/n_0$, one gets the paraxial theory applicable to an ideal first order, or linear, system without any aberrations, or nonlinearities. To treat a nonlinear or aberrating system, to a given order, one has to keep in \hat{H} the terms proportional to powers of $1/n_0$ up to the desired order of approximation. To treat a given system the corresponding Hamiltonian is chosen, up to desired order of accuracy, and the integration of the optical Schrödinger equation (1) leads to the required transfer (or Green's) function, for the evolution of $\psi(\mathbf{r})$ across the system, from the input transverse plane to the output transverse plane.

To arrive at the above formalism one usually starts with the scalar optical wave equation

$$\left(\nabla^2 - \frac{n^2(\mathbf{r})}{c^2} \frac{\partial^2}{\partial t^2} \right) \Psi(\mathbf{r}, t) = 0, \quad (4)$$

and specializes to the monochromatic case $\Psi(\mathbf{r}, t) = \psi(\mathbf{r}) \exp(-i\omega t)$. Then, $\psi(\mathbf{r})$ satisfies the Helmholtz equation

$$\left(\nabla^2 + \frac{n^2(\mathbf{r})}{\lambda^2} \right) \psi(\mathbf{r}) = 0. \quad (5)$$

Now, the optical Schrödinger equation (1) follows from rewriting (5) as

$$-\left(\bar{\lambda} \frac{\partial}{\partial z} \right)^2 \psi(\mathbf{r}) = \left(n^2(\mathbf{r}) - \hat{\mathbf{p}}_{\perp}^2 \right) \psi(\mathbf{r}), \quad (6)$$

and then choosing the 'square root' as

$$\left(i\bar{\lambda} \frac{\partial}{\partial z} \right) \psi(\mathbf{r}) = - \left(n^2(\mathbf{r}) - \hat{\mathbf{p}}_{\perp}^2 \right)^{1/2} \psi(\mathbf{r}), \quad (7)$$

corresponding to the requirement that the propagation be entirely in the positive z -direction; if the propagation is in the negative z -direction, with $\psi(\mathbf{r}) \sim \exp(-in_0 z/\bar{\lambda})$, the right hand side of (7) will have the opposite sign. As seen from (3), to first order in $1/n_0$, (7) becomes

$$i\bar{\lambda} \frac{\partial \psi(\mathbf{r})}{\partial z} \approx \left(-n_0 + \frac{1}{2n_0} \hat{\mathbf{p}}_{\perp}^2 + \varepsilon \right) \psi(\mathbf{r}), \quad (8)$$

the paraxial, or the parabolic, approximation to the Helmholtz equation.

It should be noted that the passage from (5) to (7) reduces the original boundary value problem to a first order initial value problem in z . This reduction is of great practical value since it leads to the powerful system or the Fourier optic approach [7] However, the above reduction process itself can never be claimed to be rigorous or exact. Hence there is room for alternative procedures for the reduction. Of course, any such reduction scheme is bound to lack in rigor to some extent, and the ultimate justification lies only in the results the scheme leads to. The purpose of this note is to explore one such possibility based on the analogy between (4) and the Klein-Gordon equation for a spin-0 particle. Before beginning this exploration, it is useful to recount briefly some of the other attempts to go beyond the paraxial regime.

3 Beyond the paraxial approximation

There have been several notable attempts to go beyond the paraxial approximation and, in that process, to obtain a precise knowledge of the meaning and accuracy of paraxial wave optics itself. We highlight here only some of these, and the reader may consult these works for further references.

A significant early attempt in this regard is due to Lax *et al.* [8]. These authors pointed out that the process of neglecting $\text{grad div } \vec{E}$ and seeking a solution that is plane polarized in the same sense everywhere is simply incompatible with the exact Maxwell equations. They developed an expansion procedure in powers of the beam parameter (w_0/ℓ) , where w_0 is the waist size and $\ell = 2\pi w_0^2/\lambda$ is the diffraction length or (twice the) Rayleigh range of the beam under consideration. In addition to showing that the zero-order field obeyed the Maxwell system of equations, they developed the equations obeyed by the higher-order corrections. The first-order correction was shown to be longitudinal.

Agarwal and Pattanayak [9] studied the propagation of Gaussian beam in a simple [linear, homogeneous, and isotropic] dielectric using the angular spectrum representation for the electric field, and showed that the exact solution consisted of the paraxial result plus higher-order non-Gaussian correction terms. They demonstrated, in particular, that the second-order correction term satisfied an equation consistent with the work of Lax *et al.* cited above. In another paper Agarwal and Lax [10] examined the role of the boundary condition in respect of the corrections to the paraxial approximation of Gaussian beams. This work resolved the controversy between the

work of Couture and Belanger [11] and that of Agarwal and Pattanayak [9], by tracing it to the fact that the two works had used qualitatively different boundary conditions for the correction terms: while Agarwal and Pattanayak had made the more natural demand that the field distribution in the waist plane be strictly Gaussian in the exact treatment, as in the paraxial case, Couture and Belanger had demanded the on-axis field to be the same in both treatments.

A major step in directly connecting solutions of the paraxial equation to those of the exact treatment was taken by Wünsche [12], who showed that it is possible to construct a linear *transition operator* which transforms arbitrary solutions of the paraxial equation into exact (monochromatic) solutions of the scalar wave equation (Helmholtz equation). Indeed, Wünsche constructed two such operators T_1 , T_2 corresponding to two different boundary conditions and noted, moreover, that the transition operator method is equivalent to the *complete integration of the system of coupled differential equations of Lax et al.*, restricted to the scalar case. Cao and Deng [13] derived a simpler transform operator under the condition that the evanescent waves can be ignored, and used this transform to study the corrections to the paraxial approximation of *arbitrary* freely propagating beam. They verified the consistency of their conclusions with those of the perturbation approach of Lax *et al.* Subsequently, Cao [14] applied the method of Lax et al to nonparaxial light beam propagating in a *transversely nonuniform* refractive index medium, computed the correction terms in detail, and specialized the results to the case of Gaussian beam propagating in transversely quadratic refractive index media.

The transition operator method has been further extended by Alonso *et al.* [15]. The uncertainty product has played a fundamental role in the analysis of paraxial (both coherent and partially coherent) beams. Alonso and Forbes [16] have recently generalized the uncertainty product to the case of nonparaxial fields.

There are two aspects to the issue of going beyond paraxial optics. The first one is to do with the spatial dependence alone, and hence applies to ‘scalar’ optics. The second one is to do with the vectorial nature of the light field and, more specifically, with the fact that Maxwell’s is a *constrained* system of equations. The restriction $\text{div } \vec{E} = 0$ (in free space and in homogeneous dielectrics) demands that the spatial dependence of the field, even in a transverse plane like the waist plane, cannot be chosen independent of polarization. [Thus, an input plane-polarized plane wave going through a

thin spherical lens results, in the exit plane of the lens, in a wave which is spherical and hence necessarily has spatially-varying polarization so that the Poynting vector at all points in the exit plane points to the focus of the lens.] Though the work of Lax *et al.* pointed to both these aspects, the subsequent ones noted above largely concentrated on the first.

Examining the fundamental Poincaré symmetry of the Maxwell system in the front-form of Dirac, Mukunda *et al.* [17] developed a formalism which converts a solution of the scalar wave equation into a corresponding (approximate) solution of the Maxwell system, resulting in a generalization of Fourier optics for vector electromagnetic beams [18]. This formalism leads to simple-looking electromagnetic Gaussian beams [19], and predicts a cross-polarization term [20] which is consistent with experimental observations [21]. Further analysis of electromagnetic Gaussian beams has been presented by Sheppard and Saghafi [22], and by Chen *et al.* [23].

We describe below some preliminary results of an ongoing research on the use of the FW transformations to study nonparaxial beams and the passage through optical systems which are not necessarily paraxial (Gaussian). There are two primary reasons for our believing that this approach may have advantage over the ones noted above. First, the FW technique iteratively takes the field to a new representation where the forward propagating components get progressively *decoupled* from the backward propagating components. Secondly, the FW method appears ideally suited for the Lie algebraic approach [5]. Finally, the FW technique generalizes to the vector case with very little extra effort, as will be shown in a subsequent report.

4 The Foldy-Wouthuysen formalism

In the traditional scheme the purpose of expanding the Hamiltonian $\hat{H} = -\left(n^2(\mathbf{r}) - \hat{\mathbf{p}}_{\perp}^2\right)^{1/2}$ in a series using $1/n_0$ as the expansion parameter is to understand the propagation of the quasiparaxial beam in terms of a series of approximations (paraxial + nonparaxial). Let us recall that in relativistic quantum mechanics too one has a similar problem of understanding the relativistic wave equation as the nonrelativistic approximation plus the relativistic correction terms in the quasirelativistic regime. For the Dirac equation (which is first order in time) this is done using the FW transformation leading to an iterative diagonalization technique. For the Klein-Gordon equation (which is second order in time) this is done using the same FW technique

after linearizing it with respect to time, and thus bringing it to a Dirac-like form, following the Feschbach-Villars method [6]. Nonrelativistic Schrödinger equation and the Klein-Gordon equation applicable in the cases of ion and electron optics, when the spin is disregarded, have also been treated in a similar way using the FW technique [24, 25]. The analogy between the optical wave equation (4) and the Klein-Gordon equation suggests naturally a similar technique for treating the scalar wave theory of light beams. Though the suggestion to employ the FW technique in the case of the Helmholtz equation exists in the literature as a remark [26] it has not so far been exploited to analyze the quasiparaxial approximations for any specific beam optical system.

Written as a first order system, the Helmholtz equation reads

$$i\bar{\lambda}\frac{\partial}{\partial z}\begin{bmatrix}\psi(\mathbf{r}) \\ i\bar{\lambda}\frac{\partial}{\partial z}\psi(\mathbf{r})\end{bmatrix}=\begin{bmatrix}0 & 1 \\ n^2(\mathbf{r})-\hat{\mathbf{p}}_{\perp}^2 & 0\end{bmatrix}\begin{bmatrix}\psi(\mathbf{r}) \\ i\bar{\lambda}\frac{\partial}{\partial z}\psi(\mathbf{r})\end{bmatrix}. \quad (9)$$

Now, let us define

$$\mathbf{\Psi}^{(1)}=\begin{bmatrix}\mathbf{\Psi}_+^{(1)} \\ \mathbf{\Psi}_-^{(1)}\end{bmatrix}=\frac{1}{2}\begin{bmatrix}\psi-i\frac{\bar{\lambda}}{n_0}\frac{\partial}{\partial z}\psi \\ \psi+i\frac{\bar{\lambda}}{n_0}\frac{\partial}{\partial z}\psi\end{bmatrix}. \quad (10)$$

The effect of the transformation in (10) for a quasiparaxial beam moving in the forward z -direction is to separate the component which is ‘fast’ in z from the one which is ‘slow’: there exists one linear combination of ψ and $\partial\psi/\partial z$ which varies rapidly in z and another which varies slowly in z , and the above transformation picks precisely these components. In our case of forward propagation, since $\psi(\mathbf{r})\sim\exp(in_0z/\bar{\lambda})$, we have $\mathbf{\Psi}_+^{(1)}\approx\psi(\mathbf{r})$ and $\mathbf{\Psi}_+^{(1)}\gg\mathbf{\Psi}_-^{(1)}$. In other words, $\mathbf{\Psi}_+^{(1)}$ and $\mathbf{\Psi}_-^{(1)}$ are, respectively, the *large* and the *small* components of $\mathbf{\Psi}^{(1)}$. Let us now rewrite the Helmholtz equation (5), or (9), as

$$i\bar{\lambda}\frac{\partial\mathbf{\Psi}^{(1)}}{\partial z}=\hat{\mathbf{H}}^{(1)}\mathbf{\Psi}^{(1)}, \quad (11)$$

with

$$\hat{\mathbf{H}}^{(1)}=-n_0\sigma_z+\hat{\mathcal{E}}^{(1)}+\hat{\mathcal{O}}^{(1)},$$

$$\begin{aligned}
\hat{\mathcal{E}}^{(1)} &= \left(\frac{1}{2n_0} \hat{\mathbf{p}}_{\perp}^2 + \varepsilon \right) \sigma_z, \\
\hat{\mathcal{O}}^{(1)} &= \left(\frac{1}{2n_0} \hat{\mathbf{p}}_{\perp}^2 + \varepsilon \right) (i\sigma_y),
\end{aligned} \tag{12}$$

where σ_y and σ_z are, respectively, the y and z components of the triplet of Pauli matrices, $\boldsymbol{\sigma}$, namely,

$$\sigma_x = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \sigma_y = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \sigma_z = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}. \tag{13}$$

This form of the Helmholtz equation is analogous to the Feschbach-Villars form of the Klein-Gordon equation. Note that equation (11) thus derived is algebraically very similar to the Dirac equation. Like in the Dirac equation, $\hat{\mathbf{H}}^{(1)}$ is the sum of a leading diagonal term $-n_0\sigma_z$ (analogous to $mc^2\beta$), a diagonal ‘even’ term $\hat{\mathcal{E}}^{(1)}$ which does not couple the large and the small components of $\boldsymbol{\Psi}^{(1)}$, and the off-diagonal ‘odd’ term $\hat{\mathcal{O}}^{(1)}$ which couples them. The even term $\hat{\mathcal{E}}^{(1)}$ commutes with σ_z and the odd term $\hat{\mathcal{O}}^{(1)}$ anticommutes with σ_z : $\hat{\mathcal{E}}^{(1)}\sigma_z = \sigma_z\hat{\mathcal{E}}^{(1)}$ and $\hat{\mathcal{O}}^{(1)}\sigma_z = -\sigma_z\hat{\mathcal{O}}^{(1)}$. This perfect analogy between the Dirac equation and (11) enables us to use the standard FW transformation technique to analyze (11) in terms of paraxial and higher order expansions with $1/n_0$ as the expansion parameter. This technique works as follows.

In $\hat{\mathbf{H}}^{(1)}$ the off-diagonal odd term is small compared to the diagonal part $-n_0\sigma_z + \hat{\mathcal{E}}^{(1)}$ in which the leading term is of order n_0 . A series of successive transformations of $\boldsymbol{\Psi}^{(1)}$, following a fixed recipe described below, is applied to (11) such that after each transformation the off-diagonal odd term of the resulting equation becomes weaker, being of higher order in $1/n_0$. If at any stage the odd term is considered weak enough to be neglected then one can approximate the corresponding iterated Hamiltonian by keeping only its diagonal part. Thus, this iterative diagonalization scheme can be carried out systematically up to any desired order in the expansion parameter $1/n_0$. It is interesting to note that (11) corresponds already to the paraxial approximation (8) of the Helmholtz equation if the odd term $\hat{\mathcal{O}}^{(1)}$ is dropped from $\hat{\mathbf{H}}^{(1)}$, retaining only the diagonal part $-n_0\sigma_z + \hat{\mathcal{E}}^{(1)}$.

The first FW transformation is

$$\begin{aligned}
\boldsymbol{\Psi}^{(2)} &= \exp\left(-\sigma_z\hat{\mathcal{O}}^{(1)}/2n_0\right) \boldsymbol{\Psi}^{(1)} \\
&= \exp\left[-\frac{1}{2n_0} \left(\frac{1}{2n_0} \hat{\mathbf{p}}_{\perp}^2 + \varepsilon \right) \sigma_x\right] \boldsymbol{\Psi}^{(1)}.
\end{aligned} \tag{14}$$

The result of this transformation is to turn (11) into

$$i\bar{\lambda}\frac{\partial\Psi^{(2)}}{\partial z} = \hat{H}^{(2)}\Psi^{(2)}, \quad (15)$$

with

$$\begin{aligned} \hat{H}^{(2)} &= -n_0\sigma_z + \hat{\mathcal{E}}^{(2)} + \hat{\mathcal{O}}^{(2)}, \\ \hat{\mathcal{E}}^{(2)} &= \hat{\mathcal{E}}^{(1)} - \frac{1}{2n_0}\sigma_z (\hat{\mathcal{O}}^{(1)})^2 - \frac{1}{8n_0^2} \left[\hat{\mathcal{O}}^{(1)}, [\hat{\mathcal{O}}^{(1)}, \hat{\mathcal{E}}^{(1)}] + i\bar{\lambda}\frac{\partial\hat{\mathcal{O}}^{(1)}}{\partial z} \right] + \dots, \\ \hat{\mathcal{O}}^{(2)} &= -\frac{1}{2n_0}\sigma_z \left(i\bar{\lambda}\frac{\partial\hat{\mathcal{O}}^{(1)}}{\partial z} \right) - \frac{1}{2n_0}\sigma_z [\hat{\mathcal{O}}^{(1)}, \hat{\mathcal{E}}^{(1)}] + \dots. \end{aligned} \quad (16)$$

Note that in $\hat{H}^{(1)}$ the odd term $\hat{\mathcal{O}}^{(1)}$ is of order $(1/n_0)$. In $\hat{H}^{(2)}$ the odd term $\hat{\mathcal{O}}^{(2)}$ is of order $(1/n_0)^2$. Hence in (15) the odd part is weaker than the diagonal part, by one more order, compared to (11). Further, note that the basic algebraic structure, $\hat{\mathcal{E}}^{(2)}\sigma_z = \sigma_z\hat{\mathcal{E}}^{(2)}$ and $\hat{\mathcal{O}}^{(2)}\sigma_z = -\sigma_z\hat{\mathcal{O}}^{(2)}$, is preserved by the iteration.

The second FW transformation is

$$\Psi^{(3)} = \exp\left(-\sigma_z\hat{\mathcal{O}}^{(2)}/2n_0\right)\Psi^{(2)}. \quad (17)$$

The result of this transformation is to turn (15) into

$$i\bar{\lambda}\frac{\partial\Psi^{(3)}}{\partial z} = \hat{H}^{(3)}\Psi^{(3)}, \quad (18)$$

with

$$\hat{H}^{(3)} = -n_0\sigma_z + \hat{\mathcal{E}}^{(3)} + \hat{\mathcal{O}}^{(3)}, \quad (19)$$

where $\hat{\mathcal{E}}^{(3)}$ and $\hat{\mathcal{O}}^{(3)}$ are obtained by the replacements

$$\begin{aligned} \hat{\mathcal{E}}^{(3)} &= \hat{\mathcal{E}}^{(2)} \left(\hat{\mathcal{E}}^{(1)} \rightarrow \hat{\mathcal{E}}^{(2)}, \hat{\mathcal{O}}^{(1)} \rightarrow \hat{\mathcal{O}}^{(2)} \right), \\ \hat{\mathcal{O}}^{(3)} &= \hat{\mathcal{O}}^{(2)} \left(\hat{\mathcal{E}}^{(1)} \rightarrow \hat{\mathcal{E}}^{(2)}, \hat{\mathcal{O}}^{(1)} \rightarrow \hat{\mathcal{O}}^{(2)} \right). \end{aligned} \quad (20)$$

This happens because the expressions in (16) follow as a result of the general algebraic properties of the even and odd operators, independent of the specific expressions for $\hat{\mathcal{E}}^{(1)}$ and $\hat{\mathcal{O}}^{(1)}$ given in (12). Further, $\hat{\mathcal{E}}^{(3)}\sigma_z = \sigma_z\hat{\mathcal{E}}^{(3)}$ and

$\hat{\mathcal{O}}^{(3)}\sigma_z = -\sigma_z\hat{\mathcal{O}}^{(3)}$, and $\hat{\mathcal{O}}^{(3)}$ is of order $(1/n_0)^3$. Thus the odd part of $\hat{\mathcal{H}}^{(3)}$ is weaker than the odd part of $\hat{\mathcal{H}}^{(2)}$ by one more order. Now, it is straightforward to see how this sequence of FW transformations can be continued up to any desired stage, making the odd part of the resulting $\hat{\mathcal{H}}^{(j)}$ weaker by one more order at each stage.

If the FW transform process is stopped at, say, the j -th stage then one would have arrived at

$$i\bar{\lambda}\frac{\partial\Psi^{(j)}}{\partial z} = \hat{\mathcal{H}}^{(j)}\Psi^{(j)}, \quad (21)$$

with

$$\Psi^{(j)} = \begin{pmatrix} \Psi_+^{(j)} \\ \Psi_-^{(j)} \end{pmatrix}, \quad \hat{\mathcal{H}}^{(j)} = -n_0\sigma_z + \hat{\mathcal{E}}^{(j)} + \hat{\mathcal{O}}^{(j)}, \quad (22)$$

where $\hat{\mathcal{O}}^{(j)}$ is of order $(1/n_0)^j$. It is important to note that each FW transformation preserves and improves the property that the upper component of $\Psi^{(j)}$ is large compared to its lower component: $\Psi_+^{(j)} \gg \Psi_-^{(j)}$. In view of this, we can drop the odd part $\hat{\mathcal{O}}^{(j)}$ from $\hat{\mathcal{H}}^{(j)}$, as negligible compared to the diagonal part $-n_0\sigma_z + \hat{\mathcal{E}}^{(j)}$, and write

$$i\bar{\lambda}\frac{\partial\psi(\mathbf{r})}{\partial z} \approx \hat{\mathcal{H}}\psi(\mathbf{r}), \quad \hat{\mathcal{H}} = -n_0 + \hat{\mathcal{E}}_{11}^{(j)}, \quad (23)$$

where $\hat{\mathcal{E}}_{11}^{(j)}$ is the 11 matrix element of $\hat{\mathcal{E}}^{(j)}$ and $\Psi_+^{(j)}$ has been simply relabeled ψ . Equation (23) is the j -th order approximation to the Helmholtz equation in this approach.

As already noted the first order approximation corresponds to the usual paraxial theory. In this case, equation (23) becomes

$$\begin{aligned} i\bar{\lambda}\frac{\partial\psi(\mathbf{r})}{\partial z} &\approx \hat{\mathcal{H}}\psi(\mathbf{r}), \\ \hat{\mathcal{H}} &= -n_0 + \hat{\mathcal{E}}_{11}^{(1)} = -n_0 + \frac{1}{2n_0}\hat{\mathbf{p}}_{\perp}^2 + \varepsilon. \end{aligned} \quad (24)$$

Let us now look at the second order approximation. From (12), (15), (16), and (23), we have, keeping terms up to order $(1/n_0)^5$,

$$i\bar{\lambda}\frac{\partial\psi(\mathbf{r})}{\partial z} \approx \hat{\mathcal{H}}\psi(\mathbf{r}),$$

$$\begin{aligned}
\hat{\mathcal{H}} &= -n_0 + \hat{\mathcal{E}}_{11}^{(2)} \\
&= -n_0 + \left(\frac{1}{2n_0} \hat{\mathbf{p}}_{\perp}^2 + \varepsilon \right) + \frac{i\bar{\lambda}}{16n_0^3} \left[\hat{\mathbf{p}}_{\perp}^2, \frac{\partial \varepsilon}{\partial z} \right] \\
&\quad + \frac{1}{2n_0} \left(\frac{1}{2n_0} \hat{\mathbf{p}}_{\perp}^2 + \varepsilon \right)^2 + \frac{1}{2n_0^2} \left(\frac{1}{2n_0} \hat{\mathbf{p}}_{\perp}^2 + \varepsilon \right)^3 + \dots \\
&= -n_0 + \left(\frac{1}{2n_0} \hat{\mathbf{p}}_{\perp}^2 + \varepsilon \right) \\
&\quad + \frac{\bar{\lambda}^2}{16n_0^3} \frac{\partial}{\partial z} (\hat{\mathbf{p}}_{\perp} \cdot \nabla_{\perp} \varepsilon + \nabla_{\perp} \varepsilon \cdot \hat{\mathbf{p}}_{\perp}) + \dots \quad (25)
\end{aligned}$$

Comparing (25) with the traditional expansion in (3) it is clear that $\hat{\mathcal{H}}$ has the first few terms of the traditional Hamiltonian \hat{H} correctly, plus an extra term (the commutator term). To get the higher order terms of \hat{H} in $\hat{\mathcal{H}}$ with the correct coefficients one will have to consider approximations beyond the second order. We assume, for consistency, that the derivatives of $\varepsilon(\mathbf{r})$ are also small compared to n_0 .

One may note that the FW scheme automatically leads to a Hermitian Hamiltonian without need for any further ordering of its noncommuting components.

5 Concluding remarks

It is interesting that the extra commutator term $\frac{i\bar{\lambda}}{16n_0^3} \left[\hat{\mathbf{p}}_{\perp}^2, \frac{\partial \varepsilon}{\partial z} \right]$ in (25) contributes a correction to the optical Hamiltonian, even at the ‘paraxial level’, when the refractive index of the medium suffers both longitudinal and transverse inhomogeneities. Such a z -derivative term is not natural to the traditional power series expansion. This commutator term is what survives in the commutator term $-\frac{1}{8n_0^2} \left[\hat{\mathcal{O}}^{(1)}, \left[\hat{\mathcal{O}}^{(1)}, \hat{\mathcal{E}}^{(1)} \right] + i\bar{\lambda} \frac{\partial \hat{\mathcal{O}}^{(1)}}{\partial z} \right]$ in the expression for $\hat{\mathcal{E}}^{(2)}$ in (16). In the Foldy-Wouthuysen formalism of the Dirac theory the corresponding commutator term is responsible for the correct explanation of the spin-orbit energy (including the Thomas precession effect) and the Darwin term (attributed to the zitterbewegung) (see Section 4.3 of [6]). Similarly in the nonrelativistic reduction and interpretation of the Klein-Gordon equation using the Foldy-Wouthuysen transformation theory such a commutator term corresponds to the Darwin term correcting the classical electrostatic interaction of a point charge in analogy to the zitterbewegung of the Dirac

electron (see Section 9.7 of [6]). In the quantum theory of beam optics of charged Klein-Gordon and Dirac particles [24, 25] the corresponding terms add to the Hamiltonian the lowest order quantum corrections to the classical aberration terms. In view of this analogy it should be of interest to study the effect of this correction term to the optical Hamiltonian, $\frac{i\lambda}{16n_0^3} \left[\hat{\mathbf{p}}_{\perp}^2, \frac{\partial \varepsilon}{\partial z} \right]$, on the propagation of Gaussian beams in a parabolic index medium whose focusing strength is modulated in the axial variable z , however tiny it may be compared to the classical terms.

Other questions naturally suggested by the above preliminary report are: the issue of convergence in respect of the series expansion resulting from the FW method and the boundary condition the series satisfies in relation to the paraxial result in any special plane like the waist plane of the beam. The precise relation between the FW series and the results of the other approaches recounted in Section 3 are not immediately clear at the moment, but it is an important issue worth investigating. We hope to return to these issues elsewhere.

The authors would like to thank the Referees for some insightful comments leading to improved clarity of presentation.

References

- [1] D. Gloge and D. Marcuse, J. Opt. Soc. 59 (1969) 1629.
- [2] J. R. Klauder, in: J. S. Mondragón and K. B. Wolf (Eds.), Lie Methods in Optics, Lecture Notes in Physics vol. 250, Springer, Heidelberg, 1986, p.183.
- [3] V. I. Man'ko and K. B. Wolf, in: J. S. Mondragón and K. B. Wolf (Eds.), Lie Methods in Optics, Lecture Notes in Physics vol. 250, Springer, Heidelberg, 1986, p.207.
- [4] K. B. Wolf, in: K. B. Wolf (Ed.), Lie Methods in Optics II, Lecture Notes in Physics vol. 352, Springer, Heidelberg, 1989, p.115.
- [5] A. J. Dragt, J. Opt. Soc. Am. 72 (1982) 37; Lie Algebraic Method for Ray and Wave Optics, University of Maryland Physics Department Report, 1998.

- [6] J. D. Bjorken and S. D. Drell, *Relativistic Quantum Mechanics*, McGraw-Hill, New York, 1964.
- [7] J. W. Goodman, *Introduction to Fourier Optics*, Second Ed., McGraw-Hill, New York, 1996.
- [8] M. Lax, W. H. Louisell, and W. B. Knight, *Phys. Rev. A* **11**, 1365 (1975).
- [9] G. P. Agarwal and D. N. Pattanayak, *J. Opt. Soc. Am.* **69**, 575 (1979).
- [10] G. P. Agarwal and M. Lax, *Phys. Rev. A* **27**, 1693 (1983).
- [11] M. Couture and P. A. Belanger, *Phys. Rev. A* **24**, 355 (1981).
- [12] A. Wünsche, *J. Opt. Soc. Am. A* **9**, 765 (1992).
- [13] Q. Cao and X. Deng, *J. Opt. Soc. Am. A* **15**, 1144 (1998).
- [14] Q. Cao, *J. Opt. Soc. Am. A* **16**, 2494 (1999).
- [15] M. A. Alonso, A. A. Asatryan, and G. W. Forbes, *J. Opt. Soc. Am. A* **16**, 1958 (1999).
- [16] M. A. Alonso and G. W. Forbes, *J. Opt. Soc. Am. A* **17**, 2391 (2000).
- [17] N. Mukunda, R. Simon, and E. C. G. Sudarshan, *Phys. Rev. A* **28**, 2933 (1983); See also, E. C. G. Sudarshan, R. Simon, and N. Mukunda, *Phys. Rev. A* **28**, 2921 (1983).
- [18] N. Mukunda, R. Simon, and E. C. G. Sudarshan, *J. Opt. Soc. Am. A* **2**, 416 (1985).
- [19] R. Simon, E. C. G. Sudarshan, and N. Mukunda, *J. Opt. Soc. Am. A* **3**, 416 (1986).
- [20] R. Simon, E. C. G. Sudarshan, and N. Mukunda, *Appl. Opt.* **26**, 1589 (1987).
- [21] Y. Feinman and J. Shamir, *Appl. Opt.* **23**, 3188 (1984).
- [22] C. J. R. Sheppard and S. Saghafi, *J. Opt. Soc. Am. A* **16**, 1381 (1999).

- [23] C. G. Chen, P. T. Konkola, J. Ferrera, R. K. Heilmann, and M. L. Schattenburg, *J. Opt. Soc. Am. A* **19**, 404 (2002).
- [24] S. A. Khan and R. Jagannathan, *Phys. Rev. E* 51 (1995) 2510.
- [25] R. Jagannathan and S. A. Khan, in: P. W. Hawkes (Ed.), *Advances in Imaging and Electron Physics* vol.97, Academic Press, San Diego, 1996, p.257; M. Conte, R. Jagannathan, S. A. Khan and M. Pusterla, Preprint: IMSc/96/03/07, INFN/AE-96/08, *Particle Accelerators*. **56**, (1996) 99; physics/9803042; physics/0101060; physics/9809032; physics/9904063; physics/0112085; physics/0112086.
- [26] L. Fishman and J. J. McCoy, *J. Math. Phys.* **25** (1984) 285.